

## On the existence of steady-state solutions to the Navier-Stokes system for large fluxes

ANTONIO RUSSO AND GIULIO STARITA

**Abstract.** In this paper we deal with the stationary Navier-Stokes problem in a domain  $\Omega$  with compact Lipschitz boundary  $\partial\Omega$  and datum  $\mathbf{a}$  in Lebesgue spaces. We prove existence of a solution for arbitrary values of the fluxes through the connected components of  $\partial\Omega$ , with possible countable exceptional set, provided  $\mathbf{a}$  is the sum of the gradient of a harmonic function and a sufficiently small field, with zero total flux for  $\Omega$  bounded.

**Mathematics Subject Classification (2000):** 76D05 (primary); 31B10, 35Q30, 42B20 (secondary).

### 1. Introduction

The boundary value problem associated with the Navier-Stokes equations is to find a solution to the system

$$\nu \Delta \mathbf{u} - \mathbf{u} \cdot \nabla \mathbf{u} = \nabla p \quad \text{in } \Omega, \quad (1.1)$$

$$\operatorname{div} \mathbf{u} = 0 \quad \text{in } \Omega, \quad (1.2)$$

$$\mathbf{u} = \mathbf{a} \quad \text{on } \partial\Omega, \quad (1.3)$$

where  $\Omega$  is a bounded domain (open connected set) of  $\mathbb{R}^n$  ( $n = 2, 3$ ),  $\mathbf{u}$ ,  $p$  the unknown kinetic and pressure fields,  $\nu$  the kinematical viscosity and  $\mathbf{a}$  the boundary datum which must satisfy the condition

$$\int_{\partial\Omega} \mathbf{a} \cdot \mathbf{n} = 0,$$

where  $\mathbf{n}$  is the outward unit normal to  $\partial\Omega$  (see [6]). Existence of a variational solution

$$(\mathbf{u}, p) \in [W_{\sigma}^{1,2}(\Omega) \cap C^{\infty}(\Omega)] \times [L^2(\Omega) \cap C^{\infty}(\Omega)] \quad (1.4)$$

Received January 29, 2007; accepted in revised form December 13, 2007.

to system (1.1)-(1.3) is known under the hypothesis of smallness of the fluxes

$$\Phi_i = \int_{\partial\Omega_i} \mathbf{a} \cdot \mathbf{n}$$

through the connected components  $\partial\Omega_i$  of  $\partial\Omega$ , provided  $\mathbf{a} \in W^{1/2,2}(\partial\Omega)$  and  $\partial\Omega$  is Lipschitz. Precisely, in [1, 5] (see also [6, Chapter VIII]) it is proved that there is a positive constant depending on  $\Omega$  such that if  $\sum |\Phi_i|$  is suitably small, then (1.1)-(1.3) has a variational solution. These results have been extended for  $\mathbf{a}$  in Lebesgue's spaces in [10]. In particular, it is proved that if  $\mathbf{a} \in L^q(\partial\Omega)$  ( $q \geq 8/3$  for  $n = 3$  and  $q = 2$  for  $n = 2$ ), then system (1.1)-(1.3) has a  $C^\infty$  solution which for  $q > 4$  takes the boundary datum in the sense of nontangential convergence. Moreover, making use of some regularity results in [2, 12], it is showed that if  $\mathbf{a}$  is more regular (say Hölder continuous, with Hölder's coefficient depending on the Lipschitz character of  $\partial\Omega$ ) then so does  $\mathbf{u}$ .

In [3] H. Fujita and H. Morimoto considered problem (1.1)-(1.3) in a regular domain with

$$\mathbf{a} = \mu \mathbf{u}_0|_{\partial\Omega} + \boldsymbol{\gamma}, \tag{1.5}$$

$\mu \in \mathbb{R}$ ,  $\mathbf{u}_0 = \nabla\beta$  and  $\beta$  harmonic function. They proved that if  $\Omega$  is regular,  $\beta|_{\partial\Omega} \in W^{2,2}(\partial\Omega)$ ,

$$\int_{\partial\Omega} \boldsymbol{\gamma} \cdot \mathbf{n} = 0 \tag{1.6}$$

and  $\|\boldsymbol{\gamma}\|_{W^{1/2,2}(\partial\Omega)}$  is less than a suitable constant depending on  $v, \mu, \Omega$  and  $\mathbf{u}_0$ , then system (1.1)-(1.3), (1.5) admits a solution (1.4) for any  $\mu \in \mathbb{R} \setminus G$ , with  $G$  countable subset of  $\mathbb{R}$ . Moreover, if  $\beta \in W^{3,2}(\Omega)$  and  $\|\boldsymbol{\gamma}\|_{W^{3/2,2}(\partial\Omega)}$  is sufficiently small, then  $(\mathbf{u}, p) \in W_{\sigma}^{2,2}(\Omega) \times W^{1,2}(\Omega)$ . This result is remarkable in view of the fact that, even though for special boundary data, it assures the existence of a solution to system (1.1)-(1.3) in arbitrary bounded regular domains for large fluxes. It is worth to mention that for the annulus  $\{x \in \mathbb{R}^2 : R_1 < |x| < R_2\}$  and  $\beta = \nabla \log |x|$ , H. Morimoto proved that  $G = \emptyset$  [8] (see also [4, 9]).

The aim of the present paper is twofold:

- (i) to extend the results of [3] under more general assumptions on the domains and on the data and for any  $n \geq 2$ ; in particular, for  $n = 3$  we show that, if  $\Omega$  is a bounded Lipschitz domain,  $\mathbf{a}$  is given by (1.5) with  $\mathbf{u}_0 \in W^{1,q}(\Omega)$ ,  $q > 3/2$ ,  $\boldsymbol{\gamma} \in L^2(\partial\Omega)$  satisfies (1.6) and  $\|\boldsymbol{\gamma}\|_{L^2(\partial\Omega)}$  is sufficiently small, then system (1.1)-(1.3) has a solution  $(\mathbf{u}, p) \in [W_{\sigma}^{1/2,2}(\Omega) \cap C^\infty(\Omega)] \times C^\infty(\Omega)$  for any  $\mu \in \mathbb{R} \setminus G$ , with  $G$  countable subset of  $\mathbb{R}$ ;
- (ii) to prove existence of a solution for system (1.1)-(1.3), (1.5) in a Lipschitz exterior domain in the class  $[L_{\sigma}^{\infty}(\Omega, r) \cap C^\infty(\Omega)] \times [L^{\infty}(\mathbb{C}S_R, r^2 \log r) \cap C^\infty(\Omega)]$ , provided  $\mathbf{u}_0 \in L^{\infty}(\Omega, r^2)$  and  $\|\boldsymbol{\gamma}\|_{L^{\infty}(\partial\Omega)}$  is sufficiently small.

NOTATION – We use a standard vector notation, as in [6]. Let  $\Omega$  be a domain of  $\mathbb{R}^n$ ,  $n \geq 2$ , and let  $\{\boldsymbol{\gamma}(\xi)\}_{\xi \in \partial\Omega}$  be a family of circular finite (not empty) cones with vertex

at  $\xi$  such that  $\gamma(\xi) \setminus \{\xi\} \subset \Omega$  (as well-known, if  $\Omega$  is Lipschitz, such a family of cones certainly exists). Let  $\chi$  be a function in  $\Omega$ ;  $\chi(x)$  is said to converge nontangentially at the boundary if  $\chi(\xi) = \lim_{x \rightarrow \xi} \chi(x) \Leftrightarrow \chi(x) \xrightarrow{\text{nt}} \chi(\xi)$ , for almost all  $\xi \in \partial\Omega$ . As customary,  $L^q(\Omega)$ ,  $W^{s,q}(\Omega)$  and  $L^q(\partial\Omega)$ ,  $W^{s,q}(\partial\Omega)$  ( $q \in [1, +\infty]$ ,  $s \geq 0$ ) denote respectively the Lebesgue and the Sobolev-Besov spaces of (scalar, vector and tensor) fields in  $\Omega$  and  $\partial\Omega$  endowed with their natural norms;  $W^{-s,q}(\partial\Omega)$  is the dual space of  $W^{s,q}(\partial\Omega)$  and  $L^\infty(\Omega, f(r))$ , with  $f(r)$  positive function of  $r = |x|$ , is the Banach space of all measurable fields  $\chi$  in  $\Omega$  such that  $\|f(r)\chi\|_{L^\infty(\Omega)} < +\infty$ ; if  $V \subset L^1_{\text{loc}}(\Omega)$  is a function space,  $V_\sigma$  stands for the subspace of  $V$  of all (weakly) divergence free vector fields; also, the subscript  $\phi$  in the symbol  $W_\phi$ , where  $W \subset W^{s,q}(\partial\Omega)$ ,  $s \in \mathbb{R}$ , denotes the set of all fields  $\mathbf{a} \in W$  such that  $\int_{\partial\Omega} \mathbf{a} \cdot \mathbf{n} \, d\sigma = 0$  or  $\langle \mathbf{a}, \mathbf{n} \rangle = 0$  respectively for  $s \geq 0$  or  $s < 0$ , where  $\langle \cdot, \cdot \rangle$  stands for the duality pairing between  $W^{s,q}(\partial\Omega)$  and its dual.

## 2. Some results for the Stokes system

The boundary-value problem associated with the Stokes system is to find a solution to the problem

$$\begin{aligned} \nu \Delta \mathbf{u} &= \nabla p && \text{in } \Omega, \\ \operatorname{div} \mathbf{u} &= 0 && \text{in } \Omega, \\ \mathbf{u} &= \mathbf{a} && \text{on } \partial\Omega. \end{aligned} \tag{2.1}$$

If  $\Omega$  is exterior we require that  $\mathbf{u}$  tends to zero at infinity.

The following theorems are proved in [2, 7, 10, 12].

**Theorem 2.1.** *Let  $\Omega$  be a Lipschitz bounded domain of  $\mathbb{R}^n$  and let  $\mathbf{a} \in L^q_\phi(\partial\Omega)$ ,  $q \geq 2$ . There exists a positive constant  $\epsilon$  depending on  $\Omega$  such that if  $q \in [2, 2 + \epsilon)$ , then system (2.1), admits a solution  $(\mathbf{u}, p) \in [W^{1/q,q}_\sigma(\Omega) \cap C^\infty(\Omega)] \times C^\infty(\Omega)$ ,  $\mathbf{u} \xrightarrow{\text{nt}} \mathbf{a}$  and*

$$\|\mathbf{u}\|_{W^{1/q,q}(\Omega)} \leq c \|\mathbf{a}\|_{L^q(\partial\Omega)}. \tag{2.2}$$

Moreover:

- (i) if  $\mathbf{a} \in W^{1,q}(\partial\Omega)$ ,  $q \in [2, 2 + \epsilon)$ , then  $\mathbf{u} \in W^{1+1/q,q}(\Omega)$ .
- For  $n = 3$  there are two positive constants  $\epsilon$  and  $\alpha_0$ , depending on the Lipschitz character of  $\partial\Omega$  such that:

(ii) if  $\mathbf{a} \in L^q(\partial\Omega)$ ,  $q \in [2, +\infty]$ , then  $\mathbf{u} \in W^{1/q,q}(\Omega)$  and (2.2) holds;

(iii) if  $\mathbf{a} \in W^{1-1/q,q}(\partial\Omega)$ ,  $q \in [2, 3 + \epsilon)$ , then

$$\|\mathbf{u}\|_{W^{1,q}(\Omega)} \leq c \|\mathbf{a}\|_{W^{1-1/q,q}(\partial\Omega)}; \tag{2.3}$$

(iv) if  $\mathbf{a} \in C^{0,\alpha}(\partial\Omega)$ ,  $\alpha \in [0, \alpha_0)$ , then

$$\|\mathbf{u}\|_{C^{0,\alpha}(\overline{\Omega})} \leq c \|\mathbf{a}\|_{C^{0,\alpha}(\partial\Omega)}.$$

- If  $n = 4$  and  $\mathbf{a} \in L^3(\partial\Omega)$ , then

$$\|\mathbf{u}\|_{L^4(\Omega)} \leq c \|\mathbf{a}\|_{L^3(\partial\Omega)}. \tag{2.4}$$

If  $\Omega$  is of class  $C^1$ , then properties (i)-(iv) are satisfied for all  $n \geq 2$  with  $q \in (1, +\infty)$ ,  $\epsilon = +\infty$  and  $\alpha_0 = 1$ . In particular, if  $n \geq 5$  and  $\mathbf{a} \in L^{n-1}(\partial\Omega)$ , then

$$\|\mathbf{u}\|_{L^n(\Omega)} \leq c \|\mathbf{a}\|_{L^{n-1}(\partial\Omega)}. \tag{2.5}$$

**Theorem 2.2.** Let  $\Omega$  be a bounded domain of  $\mathbb{R}^n$  of class  $C^{2,1}$  and let  $\mathbf{a} \in W_{\phi}^{-1/q,q}(\partial\Omega)$ , with  $q \in (1, +\infty)$ . Then system (2.1), admits a solution  $(\mathbf{u}, p) \in C_{\sigma}^{\infty}(\Omega) \times C^{\infty}(\Omega)$  such that  $\mathbf{u}$  takes the boundary value  $\mathbf{a}$  in the sense of the space  $W^{-1/q,q}(\partial\Omega)^1$  and

$$\|\mathbf{u}\|_{L^q(\Omega)} \leq c \|\mathbf{a}\|_{W^{-1/q,q}(\partial\Omega)}. \tag{2.6}$$

**Theorem 2.3.** Let  $\Omega$  be an exterior domain of  $\mathbb{R}^3$ . If  $\mathbf{a} \in L^{\infty}(\partial\Omega)$ , then system (2.1) admits a solution  $(\mathbf{u}, p) \in [L_{\sigma}^{\infty}(\Omega, r) \cap C^{\infty}(\Omega)] \times [L^{\infty}(\mathbb{B}_{SR}, r^2) \cap C^{\infty}(\Omega)]$ ,  $\mathbf{u} \xrightarrow{\text{nt}} \mathbf{a}$  and

$$\|\mathbf{u}\|_{L^{\infty}(\Omega, r)} \leq c \|\mathbf{a}\|_{L^{\infty}(\partial\Omega)}. \tag{2.7}$$

Moreover, property (iv) in Theorem 2.1 holds unchanged.

### 3. Existence theorems for the Navier-Stokes system

Thanks to the results just recalled concerning the Stokes problem (2.1), we are in a position to extend the existence results of Fujita-Morimoto [3] to data in  $L_{\phi}^q(\partial\Omega)$  for Lipschitz domain and in  $W_{\phi}^{-1/q,q}(\partial\Omega)$  for domains of class  $C^{2,1}$ , for suitable  $q$ . Moreover, we also prove an existence theorem in Lipschitz exterior domains with data in  $L^{\infty}(\partial\Omega)$ .

**Theorem 3.1.** Let  $\Omega$  be a Lipschitz bounded domain of  $\mathbb{R}^3$  and let

$$\mathbf{a} = \mu \mathbf{u}_0|_{\partial\Omega} + \boldsymbol{\gamma}, \tag{3.1}$$

where  $\mu \in \mathbb{R}$ ,  $\mathbf{u}_0 = \nabla\beta$ , with  $\beta \in W^{2,q}(\Omega)$  ( $q > 3/2$ ) harmonic function, and  $\boldsymbol{\gamma} \in L_{\phi}^2(\partial\Omega)$ . Then, for every  $\mu/\nu \in \mathbb{R} \setminus G$ , with  $G$  countable subset of  $\mathbb{R}$ , there exists a constant  $\kappa = \kappa(\Omega, \nu, \mathbf{u}_0, \mu)$  such that, if

$$\|\boldsymbol{\gamma}\|_{L^2(\partial\Omega)} \leq \kappa$$

then system (1.1)-(1.3) has a solution

$$(\mathbf{u}, p) \in [W_{\sigma}^{1/2,2}(\Omega) \cap C^{\infty}(\Omega)] \times C^{\infty}(\Omega).$$

<sup>1</sup> See [10].

*Proof.* Recall that the fundamental solution to the Stokes equation is expressed by

$$\begin{aligned} \mathcal{U}(x - y) &= \frac{1}{8\pi\nu} \left[ \frac{\mathbf{1}}{|x - y|} + \frac{(x - y) \otimes (x - y)}{|x - y|^3} \right], \\ \mathcal{P}(x - y) &= \frac{1}{4\pi} \frac{x - y}{|x - y|^3}, \end{aligned}$$

where  $\mathbf{1}$  denotes the unit second-order tensor.

Let  $\mathbf{u} \in L^3_\sigma(\Omega)$ . By classical results the linear operator

$$\hat{\mathcal{L}}[\mathbf{u}](x) = - \int_{\Omega} \mathcal{U}(x - y) [\mathbf{u}_0 \cdot \nabla \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u}_0](y)$$

maps  $L^3_\sigma(\Omega)$  into  $W^{1,t}(\Omega)$  for some  $t > 3/2$ . Let  $\mathcal{L}_0[\mathbf{u}]$  be the solution to the Stokes problem with boundary datum  $-\text{tr} \hat{\mathcal{L}}[\mathbf{u}] \in W^{1-1/t,t}(\partial\Omega)$ . Since by (2.3) and the trace theorem

$$\|\mathcal{L}_0[\mathbf{u}]\|_{W^{1,t}(\Omega)} \leq c \|\text{tr} \hat{\mathcal{L}}[\mathbf{u}]\|_{W^{1-1/t,t}(\partial\Omega)} \leq \|\hat{\mathcal{L}}[\mathbf{u}]\|_{W^{1,t}(\Omega)}$$

we see that also the linear operator  $\mathcal{L}_0$  maps  $L^3_\sigma(\Omega)$  into  $W^{1,t}(\Omega)$ . Therefore, by Rellich's compactness theorem, the operator

$$\mathcal{L} = \hat{\mathcal{L}} + \mathcal{L}_0$$

maps compactly  $L^3_\sigma(\Omega)$  into itself and  $\text{tr} \mathcal{L}[\mathbf{u}] = \mathbf{0}$  on  $\partial\Omega$ .

Set

$$\mathcal{F} = \mathcal{J} - \frac{\mu}{\nu} \mathcal{L},$$

where  $\mathcal{J}$  denotes the identity map. By classical results there is a countable subset  $G$  of  $\mathbb{R}$ , with a possible accumulation at 0, such that  $\mathcal{F}$  is invertible for all  $\mu/\nu \notin G$ .

The nonlinear operator

$$\hat{\mathcal{N}}[\mathbf{u}](x) = -\frac{1}{\nu} \int_{\Omega} \mathcal{U}(x - y) (\mathbf{u} \cdot \nabla \mathbf{u})(y)$$

maps  $L^3_\sigma(\Omega)$  into  $W^{1,3/2}_\sigma(\Omega)$  and it holds

$$\|\hat{\mathcal{N}}[\mathbf{u}]\|_{W^{1,3/2}(\Omega)} \leq c \|\mathbf{u}\|_{L^3(\Omega)}^2. \tag{3.2}$$

Let  $\mathcal{N}_0[\mathbf{u}]$  be the solution to the Stokes problem with boundary datum  $-\text{tr} \hat{\mathcal{N}}[\mathbf{u}]$ . By the trace theorem and (3.2) we have

$$\|\mathcal{N}_0[\mathbf{u}]\|_{L^3(\Omega)} \leq c \|\text{tr} \hat{\mathcal{N}}[\mathbf{u}]\|_{W^{1/3,3/2}(\partial\Omega)} \leq \|\hat{\mathcal{N}}[\mathbf{u}]\|_{W^{1,3/2}(\Omega)} \leq c \|\mathbf{u}\|_{L^3(\Omega)}^2.$$

Of course, the operator

$$\mathcal{N} = \hat{\mathcal{N}} + \mathcal{N}_0$$

maps  $L^3_\sigma(\Omega)$  into itself and  $\text{tr } \mathcal{N}[\mathbf{u}] = \mathbf{0}$  on  $\partial\Omega$ .

For  $\mu/\nu \notin G$  consider the map

$$\mathbf{u}' = \mathcal{F}^{-1}[\mathbf{u}_\gamma + \mathcal{N}[\mathbf{u}]] \tag{3.3}$$

from  $L^3_\sigma(\Omega)$  into itself, where  $\mathbf{u}_\gamma$  is the solution to the Stokes problem with boundary datum  $\boldsymbol{\gamma}$ . Since

$$\|\mathcal{F}^{-1}[\mathcal{N}[\mathbf{u}]]\|_{L^3(\Omega)} \leq c_0 \|\mathbf{u}\|_{L^3(\Omega)}^2,$$

taking into account (2.2), if  $\|\boldsymbol{\gamma}\|_{L^2(\partial\Omega)}$  is chosen such that

$$\|\mathcal{F}^{-1}[\mathbf{u}_\gamma]\|_{L^3(\Omega)} < \frac{1}{4c_0},$$

then (3.3) is a contraction in the ball

$$\mathcal{S} = \left\{ \mathbf{u} \in L^3_\sigma(\Omega) : \|\mathbf{u}\|_{L^3(\Omega)} \leq \frac{1}{2c_0} \right\}.$$

Therefore, by a classical theorem of S. Banach, there is a unique field  $\mathbf{u} \in \mathcal{S}$  such that

$$\mathbf{u} = \mathcal{F}^{-1}[\mathbf{u}_\gamma + \mathcal{N}[\mathbf{u}]].$$

Hence it follows that  $\mathbf{u}$  is a solution to the equation

$$\mathbf{u} = \mathbf{u}_\gamma + \frac{\mu}{\nu} \mathcal{L}[\mathbf{u}] + \mathcal{N}[\mathbf{u}].$$

Since  $\mathbf{u}_0$  is a solution to both Stokes and Navier-Stokes equations, by taking also into account standard regularity theory we see that the field  $\mu\mathbf{u}_0 + \mathbf{u} \in C^\infty(\Omega)$  is a solution to equations (1.1)-(1.2) for a suitable pressure field  $p \in C^\infty(\Omega)$ . This solution assumes the boundary datum in the sense that  $\mathbf{u}_0 \xrightarrow{\text{nt}} \mathbf{u}_0|_{\partial\Omega}$ ,  $\mathbf{u}_\gamma \xrightarrow{\text{nt}} \boldsymbol{\gamma}$  and  $\mathcal{N}[\mathbf{u}], \mathcal{L}[\mathbf{u}]$  have zero trace on  $\partial\Omega$  as elements of the Sobolev space  $W_0^{1,3/2}(\Omega)$  (see also Remark 3.2). □

**Remark 3.2.** Assume for simplicity that  $\beta$  is a regular harmonic function. By the regularity results for the Stokes problem we have, in particular, that if the norm of  $\boldsymbol{\gamma}$  is small in the corresponding function space, then

- if  $\boldsymbol{\gamma} \in L^q(\partial\Omega)$ ,  $q > 4$ , and  $\mathbf{u}_0 \in W^{1,t}(\Omega)$ ,  $t > 3$ , then  $\mathbf{u} \xrightarrow{\text{nt}} \mathbf{a}$ ;
- if  $\boldsymbol{\gamma} \in L^\infty(\partial\Omega)$ , then  $\mathbf{u} \in L^\infty(\Omega)$

and there are two positive constants  $\epsilon$  and  $\alpha_0 (< 1)$  depending on  $\Omega$  such that

- if  $\boldsymbol{\gamma} \in L^s(\partial\Omega)$ ,  $s \in [2, 2 + \epsilon)$ , then  $\mathbf{u} \in W^{1/s,s}(\Omega)$ ,
- if  $\boldsymbol{\gamma} \in W^{1-1/s,s}(\partial\Omega)$ ,  $s \in [3/2, 3 + \epsilon)$ , then  $\mathbf{u} \in W^{1,s}(\Omega)$ ,
- if  $\boldsymbol{\gamma} \in C^{0,\alpha}(\partial\Omega)$ ,  $\alpha \in [0, \alpha_0)$ , then  $\mathbf{u} \in C^{0,\alpha}(\overline{\Omega})$ .

If  $\Omega$  is of class  $C^1$ , then the above constants  $\epsilon$  and  $\alpha_0$  can be taken arbitrarily large and equal to 1 respectively. Standard regularity results also hold for the pressure field  $p$ .

**Remark 3.3.** In virtue of Theorem 2.1 and estimates (2.3), (2.4), (2.5), existence theorems like the above one can also be established for all  $n \geq 2$ . If  $n = 2$  we can take  $\mathbf{a} \in L^{2-\epsilon}(\partial\Omega)$  with  $\epsilon$  depending on  $\Omega$  ( $\mathbf{a} \in L^q(\partial\Omega)$ ,  $q > 1$ , for  $\Omega$  of class  $C^1$ ); if  $n = 4$ , we have to require  $\mathbf{a} \in L^3(\partial\Omega)$ ; if  $n > 4$ ,  $\Omega$  must be of class  $C^1$  and  $\mathbf{a} \in L^{n-1}(\partial\Omega)$ .

Taking into account Theorem 2.2 and estimate (2.2), following the argument in the proof of the above Theorem it is not difficult to get:

**Theorem 3.4.** *Let  $\Omega$  be a bounded domain of  $\mathbb{R}^n$  of class  $C^{2,1}$  and let  $\mathbf{a}$  be given by (3.1) with  $\mu \in \mathbb{R}$ ,  $\mathbf{u}_0 = \nabla\beta$ ,  $\beta \in W^{2,q}(\Omega)$  ( $q > n$ ) harmonic function and  $\boldsymbol{\gamma} \in W_\phi^{-1/n,n}(\partial\Omega)$ . There is a countable subset  $G \subset \mathbb{R}$  such that, for  $\mu/\nu \notin G$ , there is a constant  $\kappa = \kappa(\Omega, \nu, \mathbf{u}_0, \mu)$  such that, if*

$$\|\boldsymbol{\gamma}\|_{W^{-1/n,n}(\partial\Omega)} \leq \kappa$$

then (1.1)-(1.3) has a solution

$$(\mathbf{u}, p) \in [L^n_\sigma(\Omega) \cap C^\infty(\Omega)] \times C^\infty(\Omega).$$

Taking into account the result of H. Morimoto recalled in the introduction, we also have:

**Theorem 3.5.** *Let  $\Omega$  be the annulus*

$$\Omega = \{x \in \mathbb{R}^2 : R_1 < |x| < R_2\},$$

and let  $\mathbf{a}$  be given by (3.1), with  $\mathbf{u}_0 \in W^{1,q}(\Omega)$ ,  $q > 1$ ,  $\boldsymbol{\gamma} \in L^q_\phi(\Omega)$ . If  $\|\boldsymbol{\gamma}\|_{L^q(\partial\Omega)}$  is sufficiently small, then (1.1)-(1.3) has a solution

$$(\mathbf{u}, p) \in [W_\sigma^{1/q,q}(\Omega) \cap C^\infty(\Omega)] \times C^\infty(\Omega).$$

Let us pass to treat the case of an exterior domain. The following theorem holds.

**Theorem 3.6.** *Let  $\Omega$  be an exterior Lipschitz domain of  $\mathbb{R}^3$  and let  $\mathbf{a}$  be expressed by (3.1) where  $\mathbf{u}_0|_{\partial\Omega}$ ,  $\boldsymbol{\gamma} \in L^\infty(\partial\Omega)$  and  $\mathbf{u}_0 = O(r^{-2})$ . There is a countable subset  $G \subset \mathbb{R}$  such that, for  $\mu/\nu \notin G$ , there is a constant  $\xi = \xi(\Omega, \nu, \mathbf{u}_0, \mu)$  such that, if  $\|\boldsymbol{\gamma}\|_{L^\infty(\partial\Omega)} \leq \xi$ , then the Navier-Stokes problem admits a solution*

$$(\mathbf{u}, p) \in [L^\infty_\sigma(\Omega, r) \cap C^\infty(\Omega)] \times [L^\infty(\mathbb{C}S_R, r^2 \log r) \cap C^\infty(\Omega)]$$

and  $\mathbf{u} \xrightarrow{nt} \mathbf{a}$ .

*Proof.* By well-known results about the behavior at infinity of volume potential (see, e.g., [6] Lemma II.7.2), the operator

$$\hat{\mathcal{V}}[\mathbf{u}](x) = -\nabla \int_{\Omega} \mathcal{U}(x - y)(\mathbf{u} \otimes \mathbf{u}_0 + \mathbf{u}_0 \otimes \mathbf{u})(y)$$

maps boundedly  $L^\infty(\Omega, r)$  into  $L^\infty(\Omega, r^{2-\eta})$  for every  $\eta \in (0, 1)$ . Hence  $\hat{\mathcal{V}}[\mathbf{u}] \in L^q(\Omega)$ , for every  $q > 3/2$ . Moreover, by Calderón-Zygmund's theorem  $\nabla \hat{\mathcal{V}}$  maps boundedly  $L^\infty(\Omega, r)$  into  $L^q(\Omega)$ , for every  $q > 3/2$ .

Let  $\{\mathbf{u}_k\}_{k \in \mathbb{N}}$  be a bounded sequence in  $L^\infty_\sigma(\Omega, r)$ . By what we said above  $\hat{\mathcal{V}}[\mathbf{u}_k]$  is bounded in  $W^{1,q}(\Omega)$  for every  $q > 3/2$  so that we can extract from it a subsequence, we denote by the same symbol, which converges uniformly to a field  $\mathbf{u}$  in  $C_{\text{loc}}(\bar{\Omega})$ . On the other hand

$$\begin{aligned} & \|\hat{\mathcal{V}}[\mathbf{u}_k - \mathbf{u}_h]\|_{L^\infty(\Omega, r)} \\ & \leq R \|\hat{\mathcal{V}}[\mathbf{u}_k - \mathbf{u}_h]\|_{C(\bar{\Omega} \cap S_R)} + R^{\eta-1} \|r^{2-\eta} \hat{\mathcal{V}}[\mathbf{u}_k - \mathbf{u}_h]\|_{L^\infty(\mathbb{C}S_R)} \\ & \leq R \|\hat{\mathcal{V}}[\mathbf{u}_k - \mathbf{u}_h]\|_{C(\bar{\Omega} \cap S_R)} + cR^{\eta-1}. \end{aligned}$$

Let  $\epsilon > 0$  and let  $m$  be such that for every  $h, k > m$ ,  $\|\hat{\mathcal{V}}[\mathbf{u}_k - \mathbf{u}_h]\|_{C(\bar{\Omega} \cap S_R)} < \epsilon/(2R)$ , with  $R^{1-\eta} > 2c/\epsilon$ . Therefore, from the above relation it follows that  $\|\hat{\mathcal{V}}[\mathbf{u}_k - \mathbf{u}_h]\|_{L^\infty(\Omega, r)} < \epsilon$  for all  $h, k > m$  so that  $\hat{\mathcal{V}}[\mathbf{u}_k]$  is a Cauchy sequence in  $L^\infty_\sigma(\Omega, r)$  and the operator  $\hat{\mathcal{V}}$  is compact from  $L^\infty_\sigma(\Omega, r)$  into itself. Let  $\mathcal{V}_0[\mathbf{u}]$  be the solution to the Stokes problem with boundary datum  $-\text{tr} \hat{\mathcal{V}}[\mathbf{u}]$ . It is not difficult to see that  $\mathcal{V}_0$  maps compactly  $L^\infty(\Omega, r)$  into itself. Set

$$\mathcal{G} = \mathcal{J} - \frac{\mu}{\nu} \mathcal{V}$$

with  $\mathcal{V} = \hat{\mathcal{V}} + \mathcal{V}_0$ . Since  $\mathcal{G}$  is compact, there is a countable subset  $G \subset \mathbb{R}$  such that  $\mathcal{G}$  is invertible for all  $\mu/\nu \notin G$ .

The operators

$$\hat{\mathcal{W}}[\mathbf{u}] = -\frac{1}{\nu} \nabla \int_{\Omega} \mathcal{U}(x - y)(\mathbf{u} \otimes \mathbf{u})(y)$$

and  $\mathcal{W}_0[\mathbf{u}]$ , solution to the Stokes problem with datum  $-\text{tr} \hat{\mathcal{W}}[\mathbf{u}]$ , map  $L^\infty(\Omega, r)$  into itself. Consider the map

$$\mathbf{u}' = \mathcal{G}^{-1}[\mathbf{u}_\gamma + \mathcal{W}[\mathbf{u}]] \tag{3.4}$$

for  $\mu \notin G$ , where  $\mathcal{W} = \hat{\mathcal{W}} + \mathcal{W}_0$  and  $\mathbf{u}_\gamma$  is the solution to the Stokes problem with boundary datum  $\gamma$ . It is not difficult to see that

$$\|\mathcal{G}^{-1}[\mathcal{W}[\mathbf{u}]]\|_{L^\infty(\Omega, r)} \leq c'_0 \|\mathbf{u}\|_{L^\infty(\Omega, r)}^2.$$



Therefore, taking into account (2.7), if  $\|\boldsymbol{\gamma}\|_{L^\infty(\partial\Omega)}$  is chosen such that

$$\|\mathcal{G}^{-1}[\mathbf{u}_\gamma]\|_{L^3(\Omega)} < \frac{1}{4c'_0},$$

then the map (3.4) has a fixed point  $\mathbf{u}$  and the field  $\mathbf{u} + \mu\mathbf{u}_0$  is a solution to the Navier-Stokes problem.  $\square$

**Remark 3.7.** Of course, if  $\mathbf{u}_0$  and  $\boldsymbol{\gamma}$  are more regular, then so does the solution  $(\mathbf{u}, p)$ . In particular if  $\mathbf{u}_0, \boldsymbol{\gamma} \in C(\partial\Omega)$ , then is  $\mathbf{u} \in C(\overline{\Omega})$ .

**Remark 3.8.** In virtue of the result in [13] the derivatives of  $\mathbf{u}$  have the following behavior at infinity

$$\underbrace{\nabla \dots \nabla}_{k \text{ times}} \mathbf{u} = O(r^{-1-k})$$

and

$$p = O(r^{-2} \log r), \quad \underbrace{\nabla \dots \nabla}_{k \text{ times}} p = O(r^{-2-k}),$$

with  $k \in \mathbb{N}$ .

**Remark 3.9.** As far as uniqueness of the solutions in the above theorems are concerned, we quote [10], and [6, Chapter IX]. The existence of a solution to system (1.1)-(1.3) in a Lipschitz exterior domain, with  $\mathbf{a} \in L^\infty(\partial\Omega)$ , which converges at infinity to an assigned nonzero constant vector has been recently proved in [11].

## References

- [1] W. BORCHERS and K. PILECKAS, *Note on the flux problem for stationary incompressible Navier-Stokes equations in domains with a multiply connected boundary*, Acta Appl. Math. **37** (1994), 21–30.
- [2] R.M. BROWN and Z. SHEN, *Estimates for the Stokes operator in Lipschitz domains*, Indiana Univ. Math. J. **44** (1995), 1183–1206.
- [3] H. FUJITA and H. MORIMOTO, *A remark on the existence of the Navier-Stokes flow with non-vanishing outflow condition*, GAKUTO Internat. Ser. Math. Sci. Appl. **10** (1997), 53–61.
- [4] H. FUJITA, H. MORIMOTO and H. OKAMOTO, *Stability analysis of Navier–Stokes flows in annuli*, Math. Methods Appl. Sci. **20** (1997), 959–978.
- [5] G. P. GALDI, *On the existence of steady motions of a viscous flow with nonhomogeneous boundary conditions*, Matematiche (Catania) **46** (1991), 503–524.
- [6] G. P. GALDI, “An Introduction to the Mathematical Theory of the Navier–Stokes Equations”, Vol. I, II, revised edition, Springer Tracts in Natural Philosophy, C. Truesdell (ed.), Vol. 38, Springer–Verlag, 1998.
- [7] M. MITREA and M. TAYLOR, *Navier–Stokes equations on Lipschitz domains in Riemannian manifolds*, Math. Ann. **321** (2001), 955–987.
- [8] H. MORIMOTO, *Note on the boundary value problem for the Navier–Stokes equations in 2–D domain with general outflow condition* (in Japanese), Memoirs of the Institute of Science and Technology, Meiji University **35** (1997), 95–102.

- [9] H. MORIMOTO, *General outflow condition for Navier–Stokes system*, In: “Recent Topics on Mathematical Theory of Viscous Incompressible fluid”, Lectures Notes in Num. Appl. Anal., Vol. 16, 1998, 209–224.
- [10] R. RUSSO, *On the existence of solutions to the stationary Navier–Stokes equations*, Ricerche Mat. **52** (2003), 285–348.
- [11] R. RUSSO and C. G. SIMADER, *A note on the existence of solutions to the Oseen system in Lipschitz domains*, J. Math. Fluid Mech. **8** (2006), 64–76.
- [12] Z. SHEN, *A note on the Dirichlet problem for the Stokes system in Lipschitz domains*, Proc. Amer. Math. Soc. **123** (1995), 801–811.
- [13] V. SVERÁK and T-P TSAI, *On the spatial decay of 3-D steady-state Navier-Stokes flows*, Comm. Partial Differential Equations **25** (2000), 2107–2117.

Dipartimento di Matematica  
Seconda Università degli Studi di Napoli  
Via Vivaldi, 43  
I-81100 Caserta, Italy  
a.russo@unina2.it  
giulio.starita@unina2.it