

## On flops and canonical metrics

IVAN A. CHELTSOV AND YANIR A. RUBINSTEIN

**Abstract.** This article is concerned with an observation for proving non-existence of canonical Kähler metrics. The idea is to use a rather explicit type of degeneration that applies in many situations. Namely, in a variation on a theme introduced by Ross-Thomas, we consider flops of the deformation to the normal cone. This yields a rather widely applicable notion of stability that is still completely explicit and readily computable, but with wider scope. We describe some applications in dimension two, among them, a proof of one direction of the Calabi conjecture for asymptotically logarithmic Del Pezzo surfaces.

**Mathematics Subject Classification (2010):** 14J45 (primary); 14E30, 32Q20 (secondary).

### 1. Motivation and results

A variety is slope stable in the sense of Ross-Thomas if, roughly, it is K-stable with respect to degenerations to the normal cone of its subvarieties. This notion has been studied extensively by a number of authors and has yielded many non-existence results for canonical metrics on projective Kähler manifolds. Our main purpose in this article is to introduce a slight variation on this theme by considering a somewhat more involved notion of stability that involves additional flops on the degeneration to the normal cone but that is still geometric and computable, and is partly inspired by the work of Arezzo-Della Vedova-La Nave and Li-Xu.

In this article we only develop the details of this idea in the two-dimensional case.

This already gives many new non-existence results, and most notably allows us to resolve one direction of the Calabi conjecture for asymptotically logarithmic Del Pezzo surfaces.

The NSF supported this research through grants DMS-1206284,1515703. IAC was also supported by grant 15-11-20036 of the Russian Science Foundation. YAR was also supported by a Sloan Research Fellowship.

Received September 30, 2015; accepted in revised form October 18, 2016.  
Published online March 2018.

### 1.1. Existence theorem for KEE metrics

Kähler-Einstein edge (KEE) metrics are a natural generalization of Kähler-Einstein metrics: they are smooth metrics on the complement of a divisor, and have a conical singularity of angle  $2\pi\beta$  transverse to that “complex edge”. When  $\beta = 1$ , of course, this is just an ordinary Kähler-Einstein metric, that extends smoothly across the divisor. One can think of the metric as being “bent” at an angle  $2\pi\beta$  along the divisor. In the case of Riemann surfaces, KEE metrics are just the familiar constant curvature metrics with isolated cone singularities, that have been studied since the late 19th century, *e.g.*, by Picard [26].

A basic question, whose origins trace back to Tian’s 1994 lectures in the setting of nonpositive curvature [33], extended in Donaldson’s 2009 lectures to the setting of anticanonical divisors on Fano manifolds [10], and further extended in our previous work [6] (see also the survey [28, Section 8]), is the following:

**Problem 1.1.** Under what analytic conditions on the triple  $(X, D, \beta)$  does a KEE metric exist on the Kähler manifold  $X$  bent at an angle  $2\pi\beta$  along the divisor  $D \subset X$ ?

This is partly motivated by Troyanov’s solution in the Riemann surface case [36], and was settled by Jeffres-Mazzeo-Rubinstein (sufficient condition) and Darvas-Rubinstein (sufficient and necessary conditions) in higher dimensions for smooth  $D$  [7, 18]. These results give an analytic criterion characterizing existence, once the cohomological condition

$$-K_X - (1 - \beta)D \text{ is } \mu \text{ times an ample class, for some } \mu \in \mathbb{R}, \quad (1.1)$$

is satisfied.

**Remark 1.2.** The analytic condition of [7, Theorem 9.1] is optimal and in particular improves on that of [18, Theorem 2] in the presence of automorphisms. An alternative proof of the sufficient condition of [18] was later also given by Guenancia-Paun [14], who treated the more general case of a simple normal crossing (snc)  $D$ , based on work of Berman *et al.* [4]; we refer to the survey [28] for a thorough discussion and many more references.

### 1.2. Angle increasing to $2\pi$

The existence theorem of [18] coupled with Berman’s work [3] showed that KEE metrics always exist for  $(X, D, \beta)$  when  $X$  is a Fano manifold admitting a smooth anticanonical divisor  $D$  and  $\beta$  is small [18, Corollary 1].

Following these results, considerable amount of work about KEE metrics in recent years has concerned the behavior of such metrics when the cone angle *increases towards*  $2\pi$ , the two main issues being to show that when  $X$  is Fano admitting a smooth anticanonical divisor  $D$ , then:

- (a)  $X$  admits KEE metrics with angle  $2\pi\beta$  along a smooth anticanonical divisor for all angles  $\beta < 1$  sufficiently close to 1 iff  $X$  is K-semistable;

- (b) the limit of these KEE metrics as  $\beta$  tends to 1 is a smooth KE metric iff  $X$  is K-stable.

Problems (a)-(b) attracted a good deal of work building on combined efforts of many researchers in the past two decades, culminating in a solution (see, *e.g.*, the survey [28]).

### 1.3. Angle decreasing to 0

In [6], we initiated a systematic study of the behavior in the other extreme when the *cone angle  $\beta$  goes to zero*. In partial analogy with the previous paragraph, the program initiated in [6] concerns:

- (a) Determining all triples  $(X, D, \beta)$  satisfying (1.1) with *sufficiently small  $\beta$* ;
- (b) Obtaining a condition equivalent to existence of KEE metrics for such triples;
- (c) Understanding the limit, when such exists, of these KEE metrics as  $\beta$  tends to zero.

This program is largely open. In [6] we established (a) in dimension two under the technical assumption that the pair  $(X, D)$  is strongly asymptotically log Fano (see Definition 1.3; this is satisfied, *e.g.*, when  $D$  is smooth), and made some initial progress towards (b). One of our goals in the present article is to establish one direction of the equivalence in part (b) in dimension two.

To make the notion of “sufficiently small  $\beta$ ” more precise, we introduce some terminology. Consider a pair  $(X, D)$  where  $D = \sum_{i=1}^m D_i$  is a snc divisor. Denote

$$\text{Amp}(X, D) := \left\{ \beta \in \mathbb{R}_+^m \text{ s.t. } -K_X - \sum_{i=1}^m (1 - \beta_i) D_i \text{ is ample} \right\}. \quad (1.2)$$

**Definition 1.3** ([6, Definition 1.1]). We say  $(X, D)$  is asymptotically log Fano if  $0 \in \text{Amp}(X, D)$ , and strongly asymptotically log Fano if  $\text{Amp}(X, D)$  contains a punctured neighborhood of 0 in  $\mathbb{R}_+^m \setminus \{0\}$ .

When  $m = 1$  these two notions coincide. Understanding which pairs  $(X, D)$  admit a KEE metric with a small angle along  $D$  requires understanding the class of asymptotically log Fano varieties. Recall that by Kawamata-Shokurov’s Basepoint-free Theorem if  $(X, D)$  is asymptotically log Fano, then  $|a(K_X + D)|$  (for some  $a \in \mathbb{N}$ ) is free from base points and gives a morphism

$$\eta: X \rightarrow Z,$$

so that  $Z$  is a point if and only if  $D \sim -K_X$  [31, Theorem 2.1] (see also [6, Theorem 1.9]), since  $\text{Pic}(X)$  has no torsion [17, Proposition 2.1.2]. The following conjecture, posed in our earlier work, gives a rather complete picture concerning (b)-(c) when  $D$  is smooth:

**Conjecture 1.4 ([6, Conjecture 1.11]).** Suppose that  $(X, D)$  is asymptotically log Fano manifold with  $D$  smooth and irreducible.

- (i) If  $\eta$  is birational, there exist no KEE metrics for sufficiently small  $\beta$ ;
- (ii) If  $\eta$  is not birational, then there exist KEE metrics  $\omega_\beta$  with angle  $2\pi\beta$  along  $D$  for all sufficiently small  $\beta > 0$ . Moreover, as  $\beta$  tends to zero  $(X, D, \omega_\beta)$  converges in an appropriate sense to a generalized KE metric  $\omega_\infty$  on  $X \setminus D$  that is Calabi-Yau along generic fibers of  $\eta$ .

This conjecture suggests that the existence problem for KEE metrics in the small angle regime boils down to computing a single intersection number; namely, checking whether

$$(K_X + D)^n = 0.$$

This is a rather far-reaching simplification as compared to checking the much harder condition of K-stability. Indeed, the easier direction of the Yau-Tian-Donaldson conjecture implies that a KEE metric exists only if the pair  $(X, D)$  is log K-stable [3]. However, even in dimension two, it is a very difficult problem to check (log) K-stability as it involves, in theory, computing the Futaki invariant of an infinite number of test configurations.

#### 1.4. Flop-slope stability and non-existence

When  $n = 2$ , Conjecture 1.4 (i) amounts to:

**Conjecture 1.5 ([6, Conjecture 1.6]).** Let  $S$  be a smooth surface, and let  $C$  be a smooth irreducible curve on  $S$ . Suppose that  $(S, C)$  is asymptotically log Del Pezzo. Then  $S$  admits KEE metrics with angle  $\beta$  along  $C$  for all sufficiently small  $\beta$  only if  $(K_S + C)^2 = 0$ .

Our main result is a verification of Conjecture 1.5.

**Theorem 1.6.** *Let  $S$  be a smooth surface, and let  $C$  be a smooth irreducible curve on  $S$ . Suppose that  $(S, C)$  is asymptotically log Del Pezzo and  $(K_S + C)^2 \neq 0$ . Then  $S$  does not admit KEE metrics with angle  $\beta$  along  $C$  for all sufficiently small  $\beta$ .*

**Remark 1.7.** In fact, the proof of Theorem 1.6 gives a quantitative estimate. More precisely,  $S$  does not admit KEE metrics with angle  $\beta$  along  $C$  for all  $\beta \in \text{Amp}(S, C)$  for which (5.3) is negative.

In other words, we give a completely elementary and verifiable criterion that is equivalent to log K-unstability in the small angle regime. The proof involves a modification of the notion of slope stability due to Ross and Thomas [27], where we additionally perform flops on the deformation to the normal cone. After posting our paper, Fujita was able to generalize Theorem 1.6 to any dimension for rational values of  $\beta$  using a non-explicit construction of flag ideals [11]. It would be interesting to compare our work with his. We note that we believe our methods do generalize to

higher dimensions, but such an extension is certainly highly non-trivial. We believe the interest of the notion of flop-slope stability goes beyond its application to the study of Conjecture 1.5. In Arezzo-Della Vedova-La Nave and Li-Xu [1, 22] it was shown that the generalized Futaki invariant decreases under certain modifications and our construction is partly inspired by those general results, although we do not make use of them. This construction using flops occupies most of this article, and we believe it is of independent interest. Indeed, following Ross-Thomas' definition of a slope destabilizing subvariety [27, Definition 3.8], we can define flop-slope stability in a similar manner.

Let  $D$  be a divisor in  $X$ ,  $L$  an ample line bundle over  $X$ , and  $\beta \in (0, 1]$ . We note that destabilizing test configurations for  $(X, L, D, \beta)$  are defined in Section 2.

**Definition 1.8.** We say  $(X, L, D, \beta)$  is *flop-slope unstable* with respect to a subscheme  $Z$  in  $X$  if there exists a destabilizing test configuration for  $(X, L, D, \beta)$  obtained by the degeneration to the normal cone of  $Z$  composed with a sequence of flops of subvarieties of the central fiber.

This flop-slope construction is essential to the proof of Theorem 1.6 since for asymptotically logarithmic Del Pezzo surfaces the more traditional obstructions of Matsushima, Futaki, and Ross-Thomas [12, 23, 27] are not sufficient, as examples in this article and in [6] show. We expect the method developed in this article to yield many more new examples of non-existence in different settings and in higher dimensions.

We remark that the converse to Conjecture 1.5 is open: we refer to [28, Section 9] for a discussion of partial results.

## 1.5. Organization

In Section 2 we review some preliminaries: the intersection-theoretic formula for the generalized Futaki invariant, (log) slope stability, and also derive some related useful formulas for asymptotically log Del Pezzo surfaces. In Section 3 we apply these formulas to prove Theorem 1.6 for the simplest subclass of asymptotically log Del Pezzo surfaces: the Maeda class for which  $0 \in \text{Amp}(M, D)$ . Section 4 is the heart of the article, and contains our modification of slope stability, which we call flop-slope stability. The main result here is Proposition 4.9 that gives a formula for the Futaki invariant for the flopped test configuration. Some technical intersection-theoretic result needed here is proved in Appendix A. The proof of Theorem 1.6 is then carried out in Section 5. In Section 6 we collect some further examples.

**ACKNOWLEDGEMENTS.** The authors thank J. Martínez-García and R. Thomas for comments. The authors are grateful to R. J. Berman and Chalmers University of Technology for the hospitality and financial support during Summer 2014.

## 2. Preliminaries

### 2.1. Generalized Futaki invariant

Let  $\beta = (\beta_1, \dots, \beta_m) \in (0, 1]^m$  be a vector. Let  $X$  be a normal  $\mathbb{Q}$ -factorial variety (of complex dimension  $n$ ), let  $D = \sum_{i=1}^m D_i$  be a divisor (where the  $D_i$  are distinct  $\mathbb{Q}$ -Cartier prime Weil divisors) on  $X$ , and let  $L_\beta$  be an ample  $\mathbb{R}$ -divisor on  $X$  (that a priori may depend on  $\beta$ ). Put

$$D_\beta = \sum_{i=1}^m (1 - \beta_i) D_i.$$

Let  $(\mathcal{X}, \mathcal{L}_\beta, \mathcal{D}, \beta)$  be a quadruple consisting of a normal  $\mathbb{Q}$ -factorial variety  $\mathcal{X}$  of dimension  $n + 1$ , equipped with a flat surjective map  $p: \mathcal{X} \rightarrow \mathbb{P}^1$ ,  $\mathbb{R}$ -divisor  $\mathcal{L}_\beta$ <sup>1</sup>; a divisor  $\mathcal{D} = \sum_{i=1}^m \mathcal{D}_i$  (where the  $\mathcal{D}_i$  are distinct  $\mathbb{Q}$ -Cartier prime Weil divisors) on  $\mathcal{X}$ . Suppose that all fibers of  $p$  except the fiber over  $[0 : 1]$  (which we call the central fiber) are isomorphic to  $X$ , and the divisors  $\mathcal{L}_\beta$  and  $\mathcal{D}_i$  restricted to these fibers are  $L_\beta$  and  $D_i$ , respectively. Thus,  $\text{Supp}(\mathcal{D})$  does not contain components of the fibers of  $p$  (if it did,  $\mathcal{D}_i$  restricted to different fibers would be different, but we assume the restriction is always the same, namely,  $D_i$ ), and so in particular it does not contain components of the central fiber. The generalized Futaki invariant is

$$\begin{aligned} F(\mathcal{X}, \mathcal{L}_\beta, \mathcal{D}, \beta) := & n \frac{-\left(K_{\mathcal{X}} + \sum_{i=1}^m (1 - \beta_i) \mathcal{D}_i\right) \cdot \mathcal{L}_\beta^{n-1}}{L_\beta^n} \mathcal{L}_\beta^{n+1} \\ & + (n + 1) \left(K_{\mathcal{X}} - p^*(K_{\mathbb{P}^1}) + \sum_{i=1}^m (1 - \beta_i) \mathcal{D}_i\right) \cdot \mathcal{L}_\beta^n. \end{aligned} \tag{2.1}$$

Whenever the triple  $(\mathcal{X}, \mathcal{L}_\beta, \mathcal{D})$  is a test configuration in the sense of Tian [34] and Donaldson [9], then  $F(\mathcal{X}, \mathcal{L}_\beta, \mathcal{D})$  equals its Futaki invariant in the sense of Ding-Tian or Donaldson [3, 8, 9, 22, 24, 35, 37]. If

$$L_\beta \sim_{\mathbb{R}} -\left(K_X + \sum_{i=1}^m (1 - \beta_i) D_i\right) \tag{2.2}$$

(so that  $(X, \sum_{i=1}^m (1 - \beta_i) D_i)$  is a log Fano variety), the formula for  $F(\mathcal{X}, \mathcal{L}_\beta, \mathcal{D}, \beta)$  simplifies to

$$F(\mathcal{X}, \mathcal{L}_\beta, \mathcal{D}, \beta) = n \mathcal{L}_\beta^{n+1} + (n + 1) \left(K_{\mathcal{X}} - p^* K_{\mathbb{P}^1} + \sum_{i=1}^m (1 - \beta_i) \mathcal{D}_i\right) \cdot \mathcal{L}_\beta^n. \tag{2.3}$$

<sup>1</sup> We do *not* assume  $\mathcal{L}_\beta$  is  $p$ -ample in the definition of  $F(\mathcal{X}, \mathcal{L}_\beta, \mathcal{D}, \beta)$ . An  $\mathbb{R}$ -Cartier divisor  $A$  on  $\mathcal{X}$  is  $p$ -ample (respectively,  $p$ -big) if  $A + p^* B$  is ample (respectively, big) for some divisor  $B$  on  $\mathbb{P}^1$ .

We recall the following result [3, Theorem 4.8].

**Theorem 2.1 ([3, Theorem 4.8]).** *Suppose that  $X$  is smooth,  $D = \sum_{i=1}^m D_i$  is a simple normal crossing divisor, and (2.2) holds. Let  $(\mathcal{X}, \mathcal{L}_\beta, \mathcal{D}, \beta)$  be a test configuration for  $(X, L_\beta, D_\beta)$ . Assume that  $\mathcal{L}_\beta$  is  $p$ -ample. If  $F(\mathcal{X}, \mathcal{L}_\beta, \mathcal{D}, \beta) < 0$ , then  $(X, D, \beta)$  does not admit a KEE metric.*

Test configurations for which  $\mathcal{L}_\beta$  is  $p$ -ample and  $F(\mathcal{X}, \mathcal{L}_\beta, \mathcal{D}, \beta) < 0$  are called destabilizing test configurations for  $(X, L, D, \beta)$ .

The simplest possible case (beyond a product configuration) when we can effectively apply this theorem is when  $X$  is smooth and the triple  $(\mathcal{X}, \mathcal{L}_\beta, \mathcal{D})$  is a very particular test configuration obtained via deformation to the normal cone of a smooth subvariety in  $X$ . This construction is originally due to Ross-Thomas [27]. We now turn to describe it.

### 2.2. Slope stability

Let  $X$  be a smooth variety, and let  $Z$  be a smooth subvariety in  $X$ . Consider the blow-up of  $Z \times \{[0 : 1]\}$  in  $X \times \mathbb{P}^1$ . We denote the resulting space (of complex dimension  $n + 1$ ) by  $\mathcal{X}$  and denote the blow-down map by  $\pi_Z$ . Denote the  $\pi_Z$ -exceptional divisor by  $E_Z$ . Let  $p_{\mathbb{P}^1} : X \times \mathbb{P}^1 \rightarrow \mathbb{P}^1$  and  $p_X : X \times \mathbb{P}^1 \rightarrow X$  denote the natural projections.

Put

$$p := p_{\mathbb{P}^1} \circ \pi_Z.$$

The morphism  $p : \mathcal{X} \rightarrow \mathbb{P}^1$  is flat, see [15, Proposition 9.7]. Its fibers over every point that is different from  $[0 : 1]$  are isomorphic to  $X$ . The fiber  $\mathcal{X}_0$  over  $[0 : 1] \in \mathbb{P}^1$  is the union  $E_Z \cup X_0$ , where  $X_0$  is the proper transform of  $X \times \{[0 : 1]\}$ , and

$$E_Z = \mathbb{P}(v_Z \oplus \mathcal{O}_Z) \tag{2.4}$$

is a smooth ruled variety. Here  $v_Z$  denotes the normal bundle of  $Z$  in  $X$ , and  $\mathcal{O}_Z$  denotes the trivial line bundle over  $Z$ . Of course,  $v_Z \oplus \mathcal{O}_Z$  is the normal bundle of  $Z \times \{[0 : 1]\}$  in  $X \times \mathbb{P}^1$ . Note that  $X_0$  is the blow-up of  $X$  at  $Z$ . Thus, if  $Z$  is a divisor in  $X$ , then  $X_0$  is simply a copy of  $X$ .

Denote by  $\pi_0$  the morphism  $p_X \circ \pi_Z|_{X_0} : X_0 \rightarrow X$ , which is just the blow-down map of  $Z$  in  $X$ . In fact,  $E_Z$  intersect  $X_0$  exactly at the exceptional locus of  $\pi_0$  (here we slightly abuse language, since when  $Z$  is a divisor this locus is not exceptional, but is just a copy of  $Z$ , the proper transform of  $Z$ ).

Let  $\beta = (\beta_1, \dots, \beta_m) \in (0, 1]^m$  be a vector, and let  $L_\beta$  be an ample  $\mathbb{R}$ -divisor on  $X$  that may depend on the vector  $\beta$ . Put

$$\mathcal{L}_{\beta,c} := (p_X \circ \pi_Z)^* L_\beta - cE_Z \tag{2.5}$$

for some  $c > 0$ . Recall the definition of the Seshadri constant of  $(X, Z)$  with respect to  $L_\beta$ ,

$$\epsilon(X, Z, L_\beta) = \sup \{c > 0 : \pi_0^*(L_\beta) - cE_Z|_{X_0} \text{ is ample}\}. \tag{2.6}$$

Thus, if  $c \geq \epsilon(X, Z, L_\beta)$ , then  $\mathcal{L}_\beta$  is not  $p$ -ample. The following is a special case of [27, Lemma 4.1]. We give a simple direct proof for the reader's convenience. We make use of the following simple fact more than once in this article, so we record it here:

$$\text{if } C \text{ is a curve contained in the central fiber then } C.X_0 = -C.E_Z. \quad (2.7)$$

Indeed, since  $C$  is contained in a fiber of  $p : \mathcal{X} \rightarrow \mathbb{P}^1$  and  $X_0 \cup E_Z$  is such a fiber (the central fiber) then  $C.(X_0 + E_Z) = 0$ .

**Lemma 2.2.** *Suppose that  $c \in (0, \epsilon(X, Z, L_\beta))$ . Then  $\mathcal{L}_{\beta,c}$  is  $p$ -ample.*

*Proof.* Since  $L_\beta$  is ample, by Kleiman's criterion there is a positive constant  $\gamma_0$  depending only on  $L_\beta$  such that

$$L_\beta.C \geq \gamma_0$$

for every curve  $C$  in  $X$ . Similarly, there is a positive constant  $\gamma_1$  depending on  $L_\beta$  and  $c$  alone such that

$$(\pi_0^*(L_\beta) - cE_Z|_{X_0}).C \geq \gamma_1$$

for every curve  $C \subset X_0$ , because  $c < \epsilon(X, Z, L_\beta)$ .

Put

$$\gamma := \min\{c, \gamma_0, \gamma_1\}.$$

We claim that  $\mathcal{L}_{\beta,c}.C \geq \gamma$  for every curve  $C \subset \mathcal{X}$  such that  $p(C)$  is a point. The latter implies  $p$ -ampleness of the divisor  $\mathcal{L}_{\beta,c}$ .

Let  $C$  be a curve in  $\mathcal{X}$  such that  $p(C)$  is a point (so that  $C$  lies in some fiber). If  $C$  is not in the central fiber  $E_Z \cup X_0$ , then

$$\mathcal{L}_{\beta,c}.C = L_\beta.p_X \circ \pi_Z(C) \geq \gamma_0 \geq \gamma.$$

If  $C$  is in the central fiber and is contracted by  $\pi_Z$  to a point, *i.e.*,  $C$  is contained in a fiber of  $E_Z \mapsto Z$  (a  $\mathbb{P}^{n-1}$  bundle), then  $\mathcal{L}_{\beta,c}.C = -cE_Z.C \geq c \geq \gamma$ , since  $-E_Z.C \geq 1$  in this case, as  $-E_Z$  restricts to the hyperplane bundle on each fiber. If  $C \subset E_Z$  and  $C \not\subset X_0$  and  $C$  is not contracted by  $\pi_Z$  to a point, then using (2.7),

$$\begin{aligned} \mathcal{L}_{\beta,c}.C &= L_\beta.p_X \circ \pi_Z(C) - cE_Z.C = L_\beta.p_X \circ \pi_Z(C) + cX_0.C \\ &\geq L_\beta.p_X \circ \pi_Z(C) \geq \gamma_0 \geq \gamma. \end{aligned}$$

If  $C \subset X_0$ , then

$$\mathcal{L}_{\beta,c}.C = ((p_X \circ \pi_Z)^*(L_\beta) - cE_Z)|_{X_0}.C = (\pi_0^*(L_\beta) - cE_Z|_{X_0}).\pi_Z(C) \geq \gamma_1 \geq \gamma,$$

concluding the proof. □



Let  $D = \sum_{i=1}^m D_i$  be a simple normal crossing divisor on  $X$ , where the  $D_i$  are distinct smooth prime divisors on  $X$ . We assume that  $\mathcal{D}_i$  is the proper transform of  $D_i \times \mathbb{P}^1$  in  $\mathcal{X}$  (recall Subsection 2.1).

For every  $c \in (0, \epsilon(X, Z, L_\beta))$ ,  $(\mathcal{X}, \mathcal{L}_{\beta,c}, \mathcal{D}_\beta)$  is a test configuration for  $(X, L_\beta, D_\beta)$ . Recall the following definition due to Ross-Thomas and Li-Sun.

**Definition 2.3.** The triple  $(X, L_\beta, D_\beta)$  is *slope unstable* with respect to  $Z$  if  $F(\mathcal{X}, \mathcal{L}_{\beta,c}, \mathcal{D}, \beta) < 0$  for some  $c \in (0, \epsilon(X, Z, L_\beta))$ .

Note that according to (2.6) and Lemma 2.2, the assumption on  $c$  in Definition 2.3 guarantees that Theorem 2.1 is applicable.

**Corollary 2.4.** *If  $(X, L_\beta, D_\beta)$  is slope unstable with respect to  $Z$ , then  $(X, D, \beta)$  does not admit a KEE metric.*

The importance of this corollary is that the number  $F(\mathcal{X}, \mathcal{L}_{\beta,c}, \mathcal{D}, \beta)$  is readily computable for the test configuration described in this subsection (compared to a general test configuration).

**Remark 2.5.** In all cases we considered so far, if  $F(\mathcal{X}, \mathcal{L}_{\beta,c}, \mathcal{D}, \beta) < 0$  for some  $c \in (0, \epsilon(X, Z, L_\beta))$ , then  $F(\mathcal{X}, \mathcal{L}_{\beta,c}, \mathcal{D}, \beta) < 0$  for  $c = \epsilon(X, Z, L_\beta)$ .

In the next section, we compute  $F(\mathcal{X}, \mathcal{L}_{\beta,c}, \mathcal{D}, \beta)$  in a particular situation.

### 2.3. Slope stability for logarithmic surfaces

Let us use the notation and assumptions of Subsection 2.2. Suppose, in addition, that  $D$  is a smooth curve in a smooth surface  $X$ , i.e.,  $m = 1, n = 2$ , and  $D = D_1, \mathcal{D} = \mathcal{D}_1$ , and  $Z$  is a smooth curve in  $X$ . For transparency, we put

$$S = X, S_0 = X_0, p_X = p_S, C = D = D_1, \beta = \beta_1, D_\beta = (1 - \beta)C.$$

Then  $\mathcal{X}$  is a threefold, and the fiber over  $[0 : 1] \in \mathbb{P}^1$  is the union of two surfaces  $E_Z \cup S_0$ , where  $S_0$  is the proper transform of the fiber of  $p_{\mathbb{P}^1}$  over  $[0 : 1]$ . Since  $C$  is a curve, we have  $S_0 \cong S$ . Note that the exceptional divisor  $E_Z \cong \mathbb{P}(v_Z \oplus \mathcal{O}_Z)$  is a smooth ruled surface, where  $v_Z$  denotes the normal bundle of  $Z$  in  $S$ , and  $\mathcal{O}_Z$  denotes the trivial line bundle over  $Z$ .

In the case when  $L_\beta \sim_{\mathbb{R}} -K_S - (1 - \beta)C$ , there is an explicit formula for  $F(\mathcal{X}, \mathcal{D}, \mathcal{L}_\beta, \beta)$ . First, recall some intersection formulas.

**Lemma 2.6.** *It holds,*

$$E_Z^3 = -\text{deg}(N_{Z/\mathcal{X}}) = -Z^2, \tag{2.8}$$

and

$$(p_S^* L_\beta)^3 = 0 \quad \text{and} \quad ((p_S \circ \pi_Z)^* L_\beta).E_Z^2 = -(p_S^* L_\beta).Z = -L_\beta.Z. \tag{2.9}$$

*Proof.* The first equality in (2.8) follows from [13, page 608] while the second equality follows from the fact that  $N_{Z/\mathcal{X}}$  decomposes as  $\nu_Z \oplus \mathcal{O}_Z$ . Since  $E_Z$  is the projectivization of  $N_{Z/\mathcal{X}}$ , the previous decomposition also implies (2.9) as  $\pi_Z(E_Z) = Z$ .  $\square$

**Proposition 2.7.** *Suppose that (2.2) holds. Then,*

$$F(\mathcal{X}, \mathcal{L}_{\beta,c}, \mathcal{D}, \beta) = \begin{cases} (6\beta c - 3c^2)L_{\beta,Z} + (2c^3 - 3c^2\beta)Z^2 & \text{if } Z = C \\ (6c - 3c^2)L_{\beta,Z} + (2c^3 - 3c^2)Z^2 & \text{if } Z \neq C. \end{cases}$$

*Proof.* First, using Lemma 2.6,

$$\begin{aligned} \mathcal{L}_{\beta,c}^3 &= \left( (p_S \circ \pi_Z)^* L_{\beta} - cE_Z \right)^3 \\ &= \left( (p_S \circ \pi_Z)^* L_{\beta} \right)^3 - 3c \left( (p_S \circ \pi_Z)^* L_{\beta} \right)^2 \cdot E_Z \\ &\quad + 3c^2 \left( (p_S \circ \pi_Z)^* L_{\beta} \right) \cdot E_Z^2 - c^3 E_Z^3 \\ &= (p_S^* L_{\beta})^3 + 3c^2 \left( (p_S \circ \pi_Z)^* L_{\beta} \right) \cdot E_Z^2 - c^3 E_Z^3 \\ &= 3c^2 \left( (p_S \circ \pi_Z)^* L_{\beta} \right) \cdot E_Z^2 - c^3 E_Z^3 \\ &= -3c^2 (p_S^* L_{\beta}) \cdot Z - c^3 E_Z^3 = -3c^2 L_{\beta,Z} + c^3 Z^2. \end{aligned}$$

For the second term in (2.3), suppose first that  $Z = C$  (this is only used in the second line in computing  $\mathcal{D}$ ). Using Lemma 2.6 and the formula for the canonical bundle and a general divisor under a blow-up, see [13, pages 187, 476],

$$\begin{aligned} &\left( K_{\mathcal{X}} - p^* K_{\mathbb{P}^1} + (1 - \beta)\mathcal{D} \right) \cdot \mathcal{L}_{\beta,c}^2 \\ &= \left( (p_S \circ \pi_Z)^* K_S + E_Z + (1 - \beta)\pi_Z^*(C \times \mathbb{P}^1) - (1 - \beta)E_Z \right) \\ &\quad \cdot \left( (p_S \circ \pi_Z)^* L_{\beta} - cE_Z \right)^2 \\ &= \left( (p_S \circ \pi_Z)^* K_S + (1 - \beta)(p_S \circ \pi_Z)^* C + \beta E_Z \right) \\ &\quad \cdot \left( \left( (p_S \circ \pi_Z)^* L_{\beta} \right)^2 - 2cE_Z \cdot (p_S \circ \pi_Z)^* L_{\beta} + c^2 E_Z^2 \right) \\ &= c^2 (p_S \circ \pi_Z)^* K_S \cdot E_Z^2 + (1 - \beta)c^2 (p_S \circ \pi_Z)^* C \cdot E_Z^2 \\ &\quad - 2\beta c (p_S \circ \pi_Z)^* L_{\beta} \cdot E_Z^2 + \beta c^2 E_Z^3 \\ &= -c^2 p_S^* K_S \cdot Z - (1 - \beta)c^2 \pi_Z^* C \cdot Z + 2\beta c L_{\beta,Z} - \beta c^2 Z^2 \\ &= -c^2 K_S \cdot Z - (1 - \beta)c^2 C \cdot Z + 2\beta c L_{\beta,Z} - \beta c^2 Z^2. \end{aligned}$$

If  $Z \neq C$ , then  $\mathcal{D} = (1 - \beta)\pi_Z^*(C \times \mathbb{P}^1)$ , so the previous calculation gives

$$(K_{\mathcal{X}} - p^*K_{\mathbb{P}^1} + (1 - \beta)\mathcal{D}) \cdot \mathcal{L}_{\beta,c}^2 = -c^2K_S \cdot Z - (1 - \beta)c^2C \cdot Z + 2cL_{\beta} \cdot Z - c^2Z^2.$$

Thus, if  $Z = C$ , we have

$$F(\mathcal{X}, \mathcal{L}_{\beta,c}, \mathcal{D}, \beta) = 2[-3c^2L_{\beta} \cdot Z + c^3Z^2] + 3[-c^2K_S \cdot Z + 2\beta cL_{\beta} \cdot Z - c^2Z^2],$$

while if  $Z \neq C$ , we have

$$F(\mathcal{X}, \mathcal{L}_{\beta,c}, \mathcal{D}, \beta) = 2[-3c^2L_{\beta} \cdot Z + c^3Z^2] + 3[-c^2K_S \cdot Z - (1 - \beta)c^2C \cdot Z + 2cL_{\beta} \cdot Z - c^2Z^2].$$

Plugging in (2.2) now yields the desired formulas. □

In the next section, we will show how to apply Proposition 2.7 to compute  $F(\mathcal{X}, \mathcal{L}_{\beta}, \mathcal{D}, \beta)$  in some cases (cf. Li-Sun [21, Proposition 3.15, Example 3.16]). Before doing so, we illustrate a simple example.

**Example 2.8.** Suppose that  $S = \mathbb{F}_1$  and  $C$  is a smooth rational curve in  $|E + F|$ , where  $F$  is a fiber of the natural projection  $S \rightarrow \mathbb{P}^1$ , and  $E$  is the unique  $-1$ -curve in  $S$ . Then  $L_{\beta}$  is ample for every  $\beta \in (0, 1]$ . The automorphism group of the pair  $(S, C)$  is reductive [6, Proposition 7.1] so the edge version of Matsushima’s obstruction [6, Theorem 1.12] is not applicable. If  $Z = C$  or  $Z = E$ , then  $\epsilon(S, L_{\beta}, Z) = 1 + \beta$ . In addition, if  $Z = C$ , Proposition 2.7 gives

$$F(\mathcal{X}, \mathcal{L}_{\beta,c}, \mathcal{D}, \beta) = 2(1 + \beta)(\beta^2 + 2\beta - 2)$$

for  $c = 1 + \beta$ , so  $F(\mathcal{X}, \mathcal{L}_{\beta,c}, \mathcal{D}, \beta) < 0$  for  $\beta < \sqrt{3} - 1$ . Similarly, if  $Z = E$ , Proposition 2.7 gives

$$F(\mathcal{X}, \mathcal{L}_{\beta,c}, \mathcal{D}, \beta) = (1 + \beta)(2 - \beta^2 - 2\beta),$$

for  $c = 1 + \beta$ , so  $F(\mathcal{X}, \mathcal{L}_{\beta,c}, \mathcal{D}, \beta) < 0$  for all  $\beta > \sqrt{3} - 1$ .

### 3. Maeda’s class

Let  $C$  be a smooth curve on a smooth surface  $S$ . Suppose that  $(S, C)$  is asymptotically log Del Pezzo. Put

$$L_{\beta} \sim_{\mathbb{R}} -K_S - (1 - \beta)C,$$

where  $\beta \in (0, 1]$  is such that  $L_{\beta}$  is ample.

We say  $(S, C)$  belongs to the Maeda class when  $-K_S - C$  is ample. The following result proves in a unified manner than whenever  $(S, C)$  belongs to the Maeda class, Conjecture 1.5 holds.

Alternatively, this result also follows by combining [6, Proposition 7.1] with [21] and Example 2.8.

**Proposition 3.1.** *Suppose that  $-K_S - C$  is ample. Then  $(S, C, \beta)$  does not admit a KEE metric for all sufficiently small  $\beta$ .*

**Remark 3.2.** By [6, Corollary 2.3]  $C$  is rational.

*Proof.* Pick any positive  $\gamma < \epsilon(S, Z, -K_S - C)$ . By definition,

$$-K_S - C \sim_{\mathbb{R}} \gamma Z + H$$

for some ample  $\mathbb{R}$ -divisor  $H$ . Letting  $Z := C$  then

$$L_{\beta} \sim_{\mathbb{R}} -K_S - C + \beta C \sim_{\mathbb{R}} (\gamma + \beta)C + H,$$

which implies that  $\epsilon(S, L_{\beta}, Z) \geq \gamma + \beta > \gamma$ .

Pick some  $c \in (0, \gamma]$ . Let us use notation and assumptions of Subsection 2.3. Then  $\mathcal{L}_{\beta,c}$  is  $p$ -ample by Lemma 2.2. By Remark 3.2,  $L_{\beta}.C = -(K_S + (1 - \beta)C).C = 2 + \beta C^2$ . Therefore, using Proposition 2.7, with  $c = \gamma$ ,

$$\begin{aligned} F(\mathcal{X}, \mathcal{L}_{\beta,c}, \mathcal{D}, \beta) &= -3\gamma^2 L_{\beta}.C + 2\gamma^3 C^2 + \beta(6\gamma L_{\beta}.C - 3\gamma^2 C^2) \\ &= -\gamma^2 L_{\beta}.C - 2\gamma^2(L_{\beta} - \gamma C).C + \beta(6\gamma L_{\beta}.C - 3\gamma^2 C^2) \\ &= -\gamma^2(2 + \beta C^2) - 2\gamma^2(L_{\beta} - \gamma C).C + \beta(6\gamma L_{\beta}.C - 3\gamma^2 C^2) \\ &< -2\gamma^2 + \beta(6\gamma L_{\beta}.C - 4\gamma^2 C^2), \end{aligned}$$

so  $\lim_{\beta \rightarrow 0^+} F(\mathcal{X}, \mathcal{L}_{\beta,c}, \mathcal{D}, \beta) \leq -2\gamma^2 < 0$ . Thus, Theorem 2.1 implies the desired result. □

**Remark 3.3.** One cannot drop the ampleness condition in Proposition 3.1. Indeed, if  $-K_S - C$  is not ample, then it follows from the classification in [6] and Lemma 4.3 (i) below that  $\epsilon(S, L_{\beta}, C) \leq \beta$  so the arguments used in the proof of Proposition 3.1 are no longer valid.

### 4. Flop-slope stability

We follow the notation and assumptions of Subsection 2.3. In addition, denote by  $O_1, \dots, O_r$ , distinct points on the curve  $Z$ , and let

$$\pi_O: S' \rightarrow S$$

be the blow-up of the union of these points, whose exceptional curves are

$$C'_1, \dots, C'_r \subset S',$$

with  $\pi_O(C'_i) = O_i$ . Denote by

$$C', Z' \subset S'$$

the  $\pi_O$ -proper transforms of the curves  $C, Z \subset S$ , respectively. Let

$$p_{S'}: S' \times \mathbb{P}^1 \rightarrow S', \quad p'_{\mathbb{P}^1}: S' \times \mathbb{P}^1 \rightarrow \mathbb{P}^1,$$


be the natural projections. Put

$$L_i := \{O_i\} \times \mathbb{P}^1 \subset S \times \mathbb{P}^1, \tag{4.1}$$

and let

$$\pi_L: S' \times \mathbb{P}^1 \rightarrow S \times \mathbb{P}^1$$

be the blow-up of the union of the smooth disjoint curves  $L_1, \dots, L_r$ . From now on, by abuse of notation, we identify  $S$  and  $S'$  with the fibers of  $p_{\mathbb{P}^1}$  and  $p'_{\mathbb{P}^1}$  over the point  $[0 : 1] \in \mathbb{P}^1$ , respectively. The blow-up  $\pi_O: S' \rightarrow S$  is induced by the blow-up  $\pi_L$ . In sum, there exists a commutative diagram



Let

$$\pi_{Z'}: \mathcal{X}' \rightarrow S' \times \mathbb{P}^1$$

be the blow-up of the curve  $Z' \subset S' \subset S' \times \mathbb{P}^1$ , and let

$$E_{Z'} \subset \mathcal{X}'$$

be the  $\pi_{Z'}$ -exceptional divisor. If  $Z$  is rational, then  $E_{Z'} \cong \mathbb{F}_k$  where  $k = |Z|^2$  (to see this recall (2.4)). Denote by

$$S'_0 \subset \mathcal{X}'$$

the  $\pi_{Z'}$ -proper transform of the surface  $S' \subset S' \times \mathbb{P}^1$ . Put

$$p' := p'_{\mathbb{P}^1} \circ \pi_{Z'}: \mathcal{X}' \rightarrow \mathbb{P}^1. \tag{4.2}$$

Then, it holds

$$S'_0 \cong S' \tag{4.3}$$

and

$$S'_0 \cup E_{Z'}$$

is the fiber of  $p'$  over the point  $[0 : 1]$  (the ‘‘central fiber’’). Denote by

$$C_1, \dots, C_r \subset \mathcal{X}'$$

the  $\pi_{Z'}$ -proper transform on of the curves  $C'_1, \dots, C'_r \subset S' \subset S' \times \mathbb{P}^1$ , respectively. Then  $C_i \cong \mathbb{P}^1$ .

**Lemma 4.1.** *The normal bundle of  $C_i$  in  $\mathcal{X}'$  is isomorphic to  $\mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1)$ .*

*Proof.* Since  $C_i$  is rational, by Grothendieck’s lemma [13, page 516] it holds  $N_{C_i|\mathcal{X}'} = \mathcal{O}(a) \oplus \mathcal{O}(b)$ . Thus,

$$0 \rightarrow T_{C_i} \rightarrow T_{\mathcal{X}'|C_i} \rightarrow \mathcal{O}(a) \oplus \mathcal{O}(b) \rightarrow 0,$$

implies (considering the first Chern classes) that

$$a + b + 2 - 2g(C_i) = c_1(\mathcal{X}') \cdot C_i = -K_{\mathcal{X}'} \cdot C_i. \tag{4.4}$$

Note that  $C_i \cdot E_{Z'} = 1$  since  $C'_i$  and  $Z'$  intersect transversally at one point downstairs (in  $S' \subset S' \times \mathbb{P}^1$ ). In addition,  $K_{\mathcal{X}'} = \pi_{Z'}^* K_{S' \times \mathbb{P}^1} + E_{Z'}$ . Thus,

$$\begin{aligned} K_{\mathcal{X}'} \cdot C_i &= \pi_{Z'}^* K_{S' \times \mathbb{P}^1} \cdot C_i + 1 = K_{S' \times \mathbb{P}^1} \cdot C'_i + 1 = K_{S' \times \mathbb{P}^1} \cdot C'_i + 1 \\ &= 2g(C'_i) - 2 - (C'_i)^2 + 1 = 0. \end{aligned}$$

Thus, from (4.4) we conclude that  $a + b = -2$ . Next,

$$0 \rightarrow N_{C_i|S'_0} \rightarrow N_{C_i|\mathcal{X}'} \rightarrow N_{S'_0|\mathcal{X}'|C_i} \rightarrow 0. \tag{4.5}$$

Observe that  $N_{C_i|S'_0} = \mathcal{O}_{\mathbb{P}^1}(-1)$  since  $C_i$  is a  $-1$ -curve in  $S'_0$ . Thus, taking first Chern classes and using the previous paragraph, we must have  $N_{S'_0|\mathcal{X}'|C_i} = \mathcal{O}_{\mathbb{P}^1}(-1)$ . The long exact sequence associated to (4.5) gives


$$0 = H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(-1)) \rightarrow H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(a) \oplus \mathcal{O}_{\mathbb{P}^1}(b)) \rightarrow H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(-1)) = 0,$$

implying that  $a, b < 0$ ; thus,  $a = b = -1$ . □

Thus, as described in Appendix A, we can simultaneously flop the curves  $C_1, \dots, C_r \subset \mathcal{X}'$ . Denote this composition of simple flops by  $f: \mathcal{X}' \rightarrow \hat{\mathcal{X}}'$ . Moreover, there exists a surjective morphism

$$\hat{p}': \hat{\mathcal{X}}' \rightarrow \mathbb{P}^1$$

that makes the diagram



commute. Note that  $\hat{p}'$  is flat [15, Proposition 9.7]. Let us show how to obtain  $\hat{\mathcal{X}}'$  even more explicitly by blowing up the threefold  $\mathcal{X}$ . This will also show that  $\hat{\mathcal{X}}'$  is projective.

**Remark 4.2.** Recall from Subsection 2.3 that we have a blow up  $\pi_Z: \mathcal{X} \rightarrow S \times \mathbb{P}^1$  of the curve  $Z \subset S \subset S \times \mathbb{P}^1$ , and we denoted the  $\pi_Z$ -exceptional divisor by  $E_Z$ . If  $Z$  is rational, then  $E_Z \cong \mathbb{F}_{|Z^2|}$ .

Denote by

$$\tilde{L}_1, \dots, \tilde{L}_r \subset \mathcal{X}$$

the  $\pi_Z$ -proper transforms of the curves  $L_1, \dots, L_r$  (defined in (4.1)). Then, each  $\tilde{L}_i$  intersects  $E_Z$  in a unique point, because each curve  $L_i$  intersects the curve  $Z$  transversally by the point  $O_i$ . Then there exists a birational morphism


$$\pi_{\tilde{L}}: \hat{\mathcal{X}}' \rightarrow \mathcal{X}$$

that is in fact the blow-up of the union of disjoint smooth curves  $\tilde{L}_1 \cup \dots \cup \tilde{L}_r$ . In particular, the threefold  $\hat{\mathcal{X}}'$  is projective.

Denote by

$$\hat{C}_1, \dots, \hat{C}_r \subset \hat{\mathcal{X}}'$$

the  $\pi_{\tilde{L}}$ -proper transform of the fibers of the morphism  $\pi_Z|_{E_Z}: E_Z \rightarrow Z$  over the points  $O_1, \dots, O_r$  in  $Z$ , respectively. Then there exists a commutative diagram



such that  $q$  is the blow-up of the (singular curve)  $Z + L_1 + \dots + L_r$ ,  $c_C$  is the contraction of the curves  $C_1, \dots, C_r$  to the  $r$  singular points (ordinary double points) of the threefold  $\tilde{\mathcal{X}}$ ,  $c_{\hat{C}}$  contracts the curves  $\hat{C}_1, \dots, \hat{C}_r$  on the threefold  $\hat{\mathcal{X}}'$  to the same points.

Recall from Subsection 2.3 that  $S$  is equipped with an ample divisor  $L_\beta$ . Let  $L'_\beta$  be an ample  $\mathbb{R}$ -divisor on the surface  $S'$  such that

$$L'_\beta \sim_{\mathbb{R}} \pi_O^*(L_\beta) - \sum_{i=1}^r \delta_i C'_i \tag{4.7}$$

for some real numbers  $\delta_1, \dots, \delta_r$ . Then all numbers  $\delta_1, \dots, \delta_r$  must be positive. Denote by  $\epsilon(S', Z', L'_\beta)$  the Seshadri constant of  $(S', Z')$  with respect to  $L'_\beta$ . Denote by  $\tau(S', Z', L'_\beta)$  the pseudoeffective threshold of  $(S', Z')$  with respect to  $L'_\beta$ , *i.e.*, the number

$$\sup \{c > 0 : L'_\beta - cZ' \text{ is big}\}.$$

Let  $c$  be a positive real number.

**Lemma 4.3.**

- (i) One has  $\epsilon(S, Z, L_\beta) \geq \epsilon(S', Z', L'_\beta)$  and  $\epsilon(S', Z', L'_\beta) \leq \delta_i$  for every  $i$ ;
- (ii) If  $c < \epsilon(S, Z, L_\beta)$  and  $c \geq \delta_i$  for every  $i$ , then the divisor  $L'_\beta - cZ'$  is big and, in particular,  $\tau(S', Z', L'_\beta) > \epsilon(S', Z', L'_\beta)$ .

*Proof.* The inequality  $\epsilon(S, Z, L_\beta) \geq \epsilon(S', Z', L'_\beta)$  is obvious. The inequality  $\epsilon(S', Z', L'_\beta) \leq \delta_i$  follows from  $L'_\beta \cdot C'_i = \delta_i$  and  $Z' \cdot C'_i = 1$ . Suppose that  $c < \epsilon(S, Z, L_\beta)$ . Then  $L_\beta - cZ$  is ample. Since

$$L'_\beta - cZ' \sim_{\mathbb{R}} \pi_O^*(L_\beta - cZ) + \sum_{i=1}^r (c - \delta_i)C'_i,$$

we see that the divisor  $L'_\beta - cZ'$  is big provided that  $c \geq \delta_i$  for every  $i$ . □

Let  $\mathcal{D}'$  be the proper transform of the divisor  $\mathcal{D}$  on  $\mathcal{X}'$ . Put

$$\mathcal{L}'_{\beta,c} := (p_{S'} \circ \pi_{Z'})^*(L'_\beta) - cE_{Z'}. \tag{4.8}$$

If  $c < \epsilon(S', Z', L'_\beta)$ ,  $\mathcal{L}'_{\beta,c}$  is  $p'$ -ample by Lemma 2.2.

**Remark 4.4.** If  $\mathcal{L}'_{\beta,c}$  is  $p'$ -ample, then the triple  $(\mathcal{X}', \mathcal{L}'_{\beta,c}, \mathcal{D}')$  is the test configuration obtained via deformation to the normal cone of  $Z'$  in  $S'$ .

**Definition 4.5.** Denote by  $R' \subset S' \times \mathbb{P}^1$ ,  $R_{\mathcal{X}'} \subset \mathcal{X}'$ , and  $R_{\hat{\mathcal{X}'}} \subset \hat{\mathcal{X}'}$  the proper transforms of the surface  $Z \times \mathbb{P}^1 \subset S \times \mathbb{P}^1$  with respect to the maps  $\pi_L$ , and  $\pi_L \circ \pi_{Z'}$  and  $\pi_{\hat{L}} \circ \pi_Z$ , respectively.

**Lemma 4.6.** Suppose that  $\epsilon(S', Z', L'_\beta) < c < \epsilon(S, Z, L_\beta)$  and  $c \geq \delta_i$  for every  $i$ . Then  $\mathcal{L}'_{\beta,c}$  is  $p'$ -big. Moreover, the curves  $C_1, \dots, C_r$  are the only curves in  $\mathcal{X}'$  that are mapped by  $p'$  to points and have negative intersections with  $\mathcal{L}'_{\beta,c}$ .



*Proof.* One has

$$\begin{aligned} \mathcal{L}'_{\beta,c} &\sim_{\mathbb{R}} (p_{S'} \circ \pi_{Z'})^*(L'_\beta) - cE_{Z'} \\ &\sim_{\mathbb{R}} (p_{S'} \circ \pi_{Z'})^*(L'_\beta - cZ') + c(p_{S'} \circ \pi_{Z'})^*(Z') - cE_{Z'} \\ &\sim_{\mathbb{R}} (p_{S'} \circ \pi_{Z'})^*(L'_\beta - cZ) + c\pi_{Z'}^*(R') - cE_{Z'} \\ &\sim_{\mathbb{R}} (p_{S'} \circ \pi_{Z'})^*(L'_\beta - cZ) + cR_{\mathcal{X}'}. \end{aligned}$$

Since  $L'_\beta - cZ'$  is big by Lemma 4.3, we see that  $(p_{S'} \circ \pi_{Z'})^*L'_\beta - cE_{Z'}$  is  $p_{\mathbb{P}^1} \circ \pi$ -big.

Let  $\Gamma$  be an irreducible curve in  $\mathcal{X}'$  such that  $p(\Gamma)$  is the point  $[0 : 1]$  and  $\mathcal{L}'_{\beta,c}.\Gamma < 0$ . Let us show that  $\Gamma$  is one of the curves  $C_1, \dots, C_r$ . If  $\pi_{Z'}(\Gamma)$  is a point, then

$$\mathcal{L}'_{\beta,c}.\Gamma = \left( (p_{S'} \circ \pi_{Z'})^*(L'_\beta) - cE_{Z'} \right).\Gamma = -cE_{Z'}.\Gamma > 0.$$

So,  $\pi_{Z'}(\Gamma)$  is not a point. Thus, if  $\Gamma \subset E_{Z'}$ , then

$$\begin{aligned} 0 > \mathcal{L}'_{\beta,c}.\Gamma &= \left( (p_{S'} \circ \pi_{Z'})^*(L'_\beta) - cE_{Z'} \right).\Gamma \geq L'_\beta.Z' - cE_{Z'}.\Gamma \\ &= L'_\beta.Z' + cS'_0.\Gamma > cS'_0.\Gamma, \end{aligned}$$

which implies that  $S'_0.\Gamma < 0$ . Thus,  $\Gamma \subset S'_0$ . Then,

$$\mathcal{L}'_{\beta,c}.\Gamma = \left( (p_{S'} \circ \pi_{Z'})^*(L'_\beta) - cE_{Z'} \right).\Gamma = (L'_\beta - cZ').\Gamma,$$

where we used that  $S'_0 \cong S'$ . On the other hand, (4.7) gives

$$L'_\beta - cZ' \sim_{\mathbb{R}} \pi_O^*(L_\beta - cZ) + \sum_{i=1}^r (c - \delta_i)C'_i,$$

where  $L_\beta - cZ$  is ample on  $S$ . Since  $c \geq \delta_i$  for every  $i$  by assumption, we see that the curve  $\Gamma$  must be one of the curves  $C'_1, \dots, C'_r$ .  $\square$

Let  $\hat{\mathcal{L}}'_\beta$  be the proper transform of (the class in  $\text{Pic}(\hat{\mathcal{X}}') \otimes \mathbb{R}$  of) the divisor  $\mathcal{L}'_\beta$  on the threefold  $\hat{\mathcal{X}}'$  (note that  $\hat{\mathcal{L}}'_\beta$  is well-defined, since  $f$  is an isomorphism in codimension one).

A sufficient condition for the  $\hat{p}'$ -ampleness of the divisor  $\hat{\mathcal{L}}'_\beta$  is given by the next result.

**Lemma 4.7.** *Suppose that  $\epsilon(S', Z', L'_\beta) < c < \epsilon(S, Z, L_\beta)$  and  $c \geq \delta_i$  for every  $i$ . Then  $\hat{\mathcal{L}}'_\beta$  is  $\hat{p}'$ -ample.*

*Proof.* Since  $L'_\beta$  is ample, there is a constant  $\gamma_0 > 0$  (that depend only on  $L'_\beta$ ) such that

$$L'_\beta \cdot \Omega' \geq \gamma_0$$

for every curve  $\Omega'$  in  $S'$ . Similarly, there is a constant  $\gamma_1 > 0$  (that depend on  $L'_\beta$  and  $c$  alone) such that

$$(L_\beta - cZ) \cdot \Omega \geq \gamma_1$$

for every curve  $\Omega \subset S$ , because  $c < \epsilon(X, Z, L_\beta)$ . Put

$$\gamma = \min \{c, \gamma_0, \gamma_1, \delta_1, \dots, \delta_r, c - \delta_1, \dots, c - \delta_r\}.$$

Let  $\Gamma$  be an irreducible curve in  $\hat{\mathcal{X}}'$  that is contracted by  $\hat{p}'$  to a point. To show that  $\hat{\mathcal{L}}'_\beta$  is  $\hat{p}'$ -ample, it is enough to prove that  $\hat{\mathcal{L}}'_\beta \cdot \Gamma \geq \gamma$ .

Denote by  $\hat{S}_0$  and  $\hat{E}_Z$  the proper transforms of the surfaces  $S_0$  and  $E_Z$  on the threefold  $\hat{\mathcal{X}}'$ , respectively. If  $\Gamma \not\subset \hat{E}_Z \cup \hat{S}_0$ , then

$$\hat{\mathcal{L}}'_\beta \cdot \Gamma \geq \gamma_0 \geq \gamma.$$

Thus, we may assume that  $\Gamma \subset \hat{E}_Z \cup \hat{S}_0$ . One the other hand, it follows from (4.6) that

$$\hat{S}_0 \cong S$$

and  $\hat{\mathcal{L}}'_\beta|_{\hat{S}_0} \sim_{\mathbb{R}} L_\beta - cZ$ . Thus, if  $\Gamma \subset \hat{S}_0$ , then

$$\hat{\mathcal{L}}'_\beta \cdot \Gamma \geq \gamma_1 \geq \gamma.$$

Hence, we may assume that  $\Gamma \subset \hat{E}_Z$ .

Denote by  $\hat{F}_1, \dots, \hat{F}_r$  the exceptional divisors of  $\pi_{\tilde{L}}$ . We may assume that  $\pi_{\tilde{L}}(\hat{F}_i) = \tilde{L}_i$  for every  $i$ . Using (4.8) and (4.7) gives

$$\hat{\mathcal{L}}'_\beta \sim_{\mathbb{R}} (p_S \circ \pi_Z \circ \pi_{\tilde{L}})^*(L_\beta) - \sum_{i=1}^r \delta_i \hat{F}_i - c\hat{E}_Z. \quad (4.9)$$

If  $\pi_{\tilde{L}}(\Gamma)$  is a point  $\tilde{L}_i \cap E_Z$ , then

$$\hat{\mathcal{L}}'_\beta \cdot \Gamma = \delta_i \geq \gamma.$$

If  $\Gamma = \hat{C}_i$ , then

$$\hat{\mathcal{L}}'_\beta \cdot \Gamma = c - \delta_i \geq \gamma.$$

If  $\Gamma$  is contracted by  $\pi_Z \circ \pi_{\tilde{L}}$  to a point in  $Z$  that is different from  $O_1, \dots, O_r$ , then

$$\hat{\mathcal{L}}'_\beta \cdot \Gamma = c \geq \gamma.$$

Thus, we may assume that  $\pi_{\tilde{L}} \circ \pi_Z(\Gamma) = Z$ . In particular, we see that

$$\Gamma \text{ is not contained in any divisor } \hat{F}_i. \tag{4.10}$$

Rewriting (4.9) and using the fact that  $\hat{E}_Z \cup_i \hat{F}_i$  is the exceptional divisor of  $\pi_Z \circ \pi_{\tilde{L}}$  gives (recall Definition 4.5),

$$\begin{aligned} \hat{\mathcal{L}}'_\beta &\sim_{\mathbb{R}} (p_S \circ \pi_Z \circ \pi_{\tilde{L}})^*(L_\beta - cZ) + c(p_S \circ \pi_Z \circ \pi_{\tilde{L}})^*Z - \sum_{i=1}^r \delta_i \hat{F}_i - c\hat{E}_Z \\ &\sim_{\mathbb{R}} (p_S \circ \pi_Z \circ \pi_{\tilde{L}})^*(L_\beta - cZ) + c(\pi_Z \circ \pi_{\tilde{L}})^*(Z \times \mathbb{P}^1) - \sum_{i=1}^r \delta_i \hat{F}_i - c\hat{E}_Z \\ &\sim_{\mathbb{R}} (p_S \circ \pi_Z \circ \pi_{\tilde{L}})^*(L_\beta - cZ) + cR_{\hat{\mathcal{X}}'} + \sum_{i=1}^r (c - \delta_i) \hat{F}_i. \end{aligned}$$

Thus, if  $\Gamma \not\subset R_{\hat{\mathcal{X}}'}$ , then since  $\Gamma$  is a finite cover of  $Z$ , degree consideration give

$$\begin{aligned} \hat{\mathcal{L}}'_\beta \cdot \Gamma &= (p_S \circ \pi_Z \circ \pi_{\tilde{L}})^*(L_\beta - cZ) \cdot \Gamma + cR_{\hat{\mathcal{X}}'} \cdot \Gamma + \sum_{i=1}^r (c - \delta_i) \hat{F}_i \cdot \Gamma \\ &\geq (L_\beta - cZ) \cdot Z + cR_{\hat{\mathcal{X}}'} \cdot \Gamma + \sum_{i=1}^r (c - \delta_i) \hat{F}_i \cdot \Gamma \\ &\geq (L_\beta - cZ) \cdot Z \geq \gamma_1 \geq \gamma, \end{aligned}$$

where we also used (4.10). Thus, we may assume that  $\Gamma \subset R_{\hat{\mathcal{X}}'}$ . Then  $\Gamma$  is the proper transform of the curve  $E_Z \cap R_{\mathcal{X}}$ . Since the surfaces  $S_0$  and  $R_{\mathcal{X}}$  are disjoint, we have  $S_0 \cdot \pi_{\tilde{L}}(\Gamma) = 0$ . Then

$$\begin{aligned} \hat{\mathcal{L}}'_\beta \cdot \Gamma &= (p_S \circ \pi_{\tilde{L}} \circ \pi_Z)^*(L_\beta) \cdot \Gamma - \sum_{i=1}^r \delta_i \hat{F}_i \cdot \Gamma - c\hat{E}_Z \cdot \Gamma \\ &= L_\beta \cdot Z - \sum_{i=1}^r \delta_i \hat{F}_i \cdot \Gamma - c\hat{E}_Z \cdot \Gamma \\ &= L_\beta \cdot Z - \sum_{i=1}^r \delta_i - c\hat{E}_Z \cdot \Gamma \\ &= L_\beta \cdot Z - \sum_{i=1}^r \delta_i - cE_Z \cdot \pi_{\tilde{L}}(\Gamma) \\ &= L_\beta \cdot Z - \sum_{i=1}^r \delta_i + cS_0 \cdot \pi_{\tilde{L}}(\Gamma) \\ &= L_\beta \cdot Z - \sum_{i=1}^r \delta_i = L'_\beta \cdot Z' \geq \gamma_0 \geq \gamma. \end{aligned}$$

This completes the proof of the lemma. □

Let  $\hat{\mathcal{D}}'$  be the proper transform of the divisor  $\mathcal{D}$  on the threefold  $\hat{\mathcal{X}}'$ , and let  $\hat{\mathcal{L}}'_\beta$  be as defined before Lemma 4.7.

**Corollary 4.8.** *Suppose that  $\epsilon(S', Z', L'_\beta) < c < \epsilon(S, Z, L_\beta)$  and  $c \geq \delta_i$  for every  $i$ . Then the quadruple  $(\hat{\mathcal{X}}', \hat{\mathcal{L}}'_\beta, \hat{\mathcal{D}}', \beta)$  is a test configuration.*

Next, we compute the generalized Futaki invariant of the flopped test configuration.

**Proposition 4.9.** *Suppose that*

$$L'_\beta \sim_{\mathbb{R}} -K_{S'} - (1 - \beta)C'.$$

Then (recall (2.3)),

$$\begin{aligned} F(\hat{\mathcal{X}}', \hat{\mathcal{L}}'_\beta, \hat{\mathcal{D}}', \beta) &= F(\mathcal{X}', \mathcal{L}'_{\beta,c}, \mathcal{D}', \beta) - 2 \sum_{i=1}^r (\mathcal{L}'_{\beta,c} \cdot C_i)^3 \\ &\quad - 3(1 - \beta) \sum_{i=1}^r (\mathcal{L}'_{\beta,c} \cdot C_i)^2 (\mathcal{D}' \cdot C_i) \\ &= 2(\mathcal{L}'_{\beta,c})^3 + 3(K_{\mathcal{X}'} - (p')^*(K_{\mathbb{P}^1}) + (1 - \beta)\mathcal{D}') \cdot (\mathcal{L}'_{\beta,c})^2 \\ &\quad - 2 \sum_{i=1}^r (\mathcal{L}'_{\beta,c} \cdot C_i)^3 - 3(1 - \beta) \sum_{i=1}^r (\mathcal{L}'_{\beta,c} \cdot C_i)^2 (\mathcal{D}' \cdot C_i). \end{aligned}$$

*Proof.* Recall from Subsection 2.1 that

$$F(\hat{\mathcal{X}}', \hat{\mathcal{L}}'_\beta, \hat{\mathcal{D}}', \beta) = 2(\hat{\mathcal{L}}'_\beta)^3 + 3(K_{\hat{\mathcal{X}}'} - (\hat{p}')^*(K_{\mathbb{P}^1}) + (1 - \beta)\hat{\mathcal{D}}') \cdot (\hat{\mathcal{L}}'_\beta)^2.$$

The assertion now follows from (2.3) and Lemma A.3, together with the fact that, as in (4.4),  $K_{\mathcal{X}'} \cdot C'_i = 0$ , while  $p^*K_{\mathbb{P}^1} \cdot C'_i = 0$  since the  $C_i$  are contained in the central fiber of  $p$ .  $\square$

## 5. Proof of Theorem 1.6

According to [6, Theorem 1.4], all asymptotically log Fano surfaces  $(S, C)$  such that  $-K_S - C$  is big satisfy either  $-K_S - C$  is ample, or  $S$  is obtained from an asymptotically log Fano surface  $(s, c)$  such that  $-K_s - c$  is ample by blowing-up  $s$  at  $r > 0$  distinct points on  $c$  and letting  $C$  denote the proper transform of  $c$ . Proposition 3.1 already established Theorem 1.6 in the first case. To complete the proof of Theorem 1.6 it remains to handle the latter case.

To that end, we switch back to the notation and assumptions of Section 4. We suppose that  $(S, C)$  is such that  $-K_S - C$  is ample (hence asymptotically log Fano), and that  $(S', C')$  is still asymptotically log Fano, *i.e.*,

$$L'_\beta := -K_{S'} - (1 - \beta)C'$$

is ample for all sufficiently small  $\beta$ . Note that  $-K_{S'} - C' = \pi_O^*(-K_S - C)$  is big being the pull-back under a birational map of an ample class. Thus,  $(S', C')$  satisfies the assumptions of Theorem 1.6. However, it is not possible to slope destabilize this latter pair in the same way as was done for  $(S, C)$  in Section 3. Indeed,

$$L'_\beta \sim_{\mathbb{R}} \pi_O^*(-K_S - (1 - \beta)C) - \beta \sum_{i=1}^r C'_i, \tag{5.1}$$

so that by Lemma 4.3 (i) (putting  $\delta_i = \beta$  and  $Z = C$ ),  $\epsilon(S', Z', L'_\beta) \leq \beta$ , and in particular using Proposition 2.7 one checks that the generalized Futaki invariant  $F(\mathcal{X}', \mathcal{L}'_{\beta,c}, \mathcal{D}, \beta)$  of the degeneration to the normal cone is positive for  $c \in (0, \beta)$ , and so  $(S', C')$  is not slope destabilized in this way. In what follows, we apply the results of Section 4 to destabilize our pair nevertheless.

Before proving Theorem 1.6, let us consider a model example.

**Example 5.1.** Suppose that  $S = \mathbb{P}^2$  and  $C$  is a smooth conic. Then  $\epsilon(S', L'_\beta, Z') = \beta$ . Thus, if  $c < \beta$ , then  $\mathcal{L}'_{\beta,c}$  is  $p'$ -ample. By Proposition 2.7, we have

$$F(\mathcal{X}', \mathcal{L}'_{\beta,c}, \mathcal{D}', \beta) = (6\beta c - 3c^2)(2 + \beta(4 - r)) + (2c^3 - 3c^2\beta)(4 - r).$$

In particular, this invariant is always positive for  $\beta$  sufficiently small (depending on  $r$ ). On the other hand,

$$\tau(S', L'_\beta, Z') = \epsilon(S, L_\beta, Z) = \frac{1}{2} + \beta.$$

Thus, if  $\beta < c < \frac{1}{2} + \beta$ , then  $\hat{\mathcal{L}}'_\beta$  is  $\hat{p}'$ -ample by Lemma 4.7. By Proposition 4.9, one has

$$\begin{aligned} F(\hat{\mathcal{X}}', \hat{\mathcal{L}}'_{\beta,c}, \hat{\mathcal{D}}', \beta) &= F(\mathcal{X}', \mathcal{L}'_{\beta,c}, \mathcal{D}', \beta) + 2r(c - \beta)^3 \\ &= (6\beta c - 3c^2)(2 + \beta(4 - r)) + (2c^3 - 3c^2\beta)(4 - r) + 2r(c - \beta)^3 \end{aligned}$$

(see Appendix Appendix A). If we put  $c = \frac{1}{2} + \beta$ , then

$$\lim_{\beta \rightarrow 0^+} F(\hat{\mathcal{X}}', \hat{\mathcal{L}}'_\beta, \hat{\mathcal{D}}', \beta) = -\frac{1}{2}.$$

Recalling the discussion at the beginning of this section, Theorem 1.6 follows from the following result.

**Proposition 5.2.** *The triple  $(S', C', \beta)$  is flop-slope unstable for all sufficiently small  $\beta$ .*

*Proof.* Let  $\epsilon(S, Z, -K_S - C)$  be the Seshadri constant of  $Z \subset S$  with respect to  $-K_S - C$ . Pick any positive  $\gamma < \epsilon(S, Z, -K_S - C)$ . Then

$$-K_S - C \sim_{\mathbb{R}} \gamma Z + H$$

for some ample  $\mathbb{R}$ -divisor  $H$ . Then

$$L_{\beta} \sim_{\mathbb{R}} -K_S - C + \beta C \sim_{\mathbb{R}} (\gamma + \beta)C + H,$$

hence  $\epsilon(S, L_{\beta}, Z) \geq \gamma + \beta$ , so  $\epsilon(S, L_{\beta}, Z) > \beta \geq \epsilon(S', L'_{\beta}, Z')$  by Lemma 4.3 (i). By taking  $\beta$  small, we may suppose that  $\gamma > \beta$ . Letting  $c$  be a real number such that

$$\epsilon(S', Z', L'_{\beta}) \leq \beta < c \leq \gamma < \epsilon(S, L_{\beta}, Z), \tag{5.2}$$

Lemma 4.7 implies that  $\hat{\mathcal{L}}'_{\beta,c}$  is  $\hat{\rho}'$ -ample. By Proposition 4.9, we have

$$\begin{aligned} F(\hat{\mathcal{X}}', \hat{\mathcal{L}}'_{\beta,c}, \hat{\mathcal{D}}', \beta) &= F(\mathcal{X}', \mathcal{L}'_{\beta,c}, \mathcal{D}', \beta) - 2 \sum_{i=1}^r (\mathcal{L}'_{\beta,c} \cdot C_i)^3 \\ &\quad - 3 \sum_{i=1}^r (\mathcal{L}'_{\beta,c} \cdot C_i)^2 (\mathcal{D}' \cdot C_i). \end{aligned}$$

Moreover, by Proposition 2.7

$$F(\mathcal{X}', \mathcal{L}'_{\beta,c}, \mathcal{D}', \beta) = (6\beta c - 3c^2)L'_{\beta} \cdot C' + (2c^3 - 3c^2\beta)(C')^2.$$

Note that using (4.8) and (5.1),

$$\mathcal{L}'_{\beta,c} \cdot C_i = ((p_{S'} \circ \pi_{Z'})^* L'_{\beta} - cE_{Z'}) \cdot C_i = L'_{\beta} \cdot C'_i - cZ' \cdot C'_i = \beta - c.$$

In addition, before the blow-up  $\pi_{Z'}$ , the intersection of  $\mathcal{D} = C' \times \mathbb{P}^1$  and  $S' \subset S' \times \mathbb{P}^1$  is precisely  $Z' \subset S' \subset S' \times \mathbb{P}^1$  (this is precisely where we use that  $Z' = C'$ ). Thus, after blowing-up  $Z'$ , the surfaces  $\mathcal{D}'$  and  $S'_0 \cong S'$  (recall (4.3)) no longer intersect. Since  $C_i$  is contained in  $S'_0$ ,

$$\mathcal{D}' \cdot C_i = 0.$$

Combining these facts,

$$F(\hat{\mathcal{X}}', \hat{\mathcal{L}}'_{\beta,c}, \hat{\mathcal{D}}', \beta) = (6\beta c - 3c^2)L'_{\beta} \cdot C' + (2c^3 - 3c^2\beta)(C')^2 + 2r(c - \beta)^3.$$

By Remark 3.2,  $C$  and hence also  $C'$  are rational, so  $L'_{\beta} \cdot C' = -(K_{S'} - (1 - \beta)C') \cdot C' = 2 + \beta C'^2$ . Thus, putting  $c = \gamma$  and grouping most terms of order  $\beta$

together yields,

$$\begin{aligned}
 F(\hat{\mathcal{X}}', \hat{\mathcal{L}}'_{\beta,\gamma}, \hat{\mathcal{D}}', \beta) &= -\gamma^2 L'_\beta \cdot C' - 2\gamma^2 (L'_\beta - \gamma C') \cdot C' + 2r\gamma^3 \\
 &\quad + \beta(6\gamma L'_\beta \cdot C' - 3\gamma^2 C'^2 - 6r\gamma^2 + 6r\beta\gamma - 2r\beta^2) \\
 &= -\gamma^2(2 + \beta C'^2) - 2\gamma^2 \left( \pi_{O'}^*(L_\beta - \gamma C) + (\gamma - \beta) \sum_{i=1}^r C'_i \right) \cdot C' + 2r\gamma^3 \\
 &\quad + \beta(6\gamma L'_\beta \cdot C' - 3\gamma^2 C'^2 - 6r\gamma^2 + 6r\beta\gamma - 2r\beta^2) \tag{5.3} \\
 &= -\gamma^2(2 + \beta C'^2) - 2\gamma^2 (L_\beta - \gamma C) \cdot C - 2\gamma^2(\gamma - \beta)r + 2r\gamma^3 \\
 &\quad + \beta(6\gamma L'_\beta \cdot C' - 3\gamma^2 C'^2 - 6r\gamma^2 + 6r\beta\gamma - 2r\beta^2) \\
 &= -2\gamma^2 - 2\gamma^2 (L_\beta - \gamma C) \cdot C \\
 &\quad + \beta(6\gamma L'_\beta \cdot C' - 4\gamma^2 C'^2 - 4r\gamma^2 + 6r\beta\gamma - 2r\beta^2),
 \end{aligned}$$

so by (5.2),

$$\begin{aligned}
 F(\hat{\mathcal{X}}', \hat{\mathcal{L}}'_{\beta,c}, \hat{\mathcal{D}}', \beta) &< -2\gamma^2 \\
 &\quad + \beta(6\gamma L'_\beta \cdot C' - 4\gamma^2 C'^2 - 4r\gamma^2 + 6r\beta\gamma - 2r\beta^2); \tag{5.4}
 \end{aligned}$$

implying that  $\lim_{\beta \rightarrow 0^+} F(\hat{\mathcal{X}}', \hat{\mathcal{L}}'_{\beta,c}, \hat{\mathcal{D}}', \beta) \leq -2\gamma^2 < 0$ . □

### 6. Further examples

We conclude by illustrating the advantage of using flop-slope stability over slope stability with two simple examples.

**Example 6.1** ( $\mathbb{F}_1$ ). According to Ross-Thomas [27, Examples 5.27, 5.35] (cf. Panov-Ross [25, Example 3.8]),  $\mathbb{F}_1$  is (for  $\beta = 1$ ) slope destabilized by the  $-1$ -curve. More generally, by Li-Sun [21], the Futaki invariant of the slope test configuration of the triple  $(\mathbb{F}_1, C, \beta)$  with  $C$  smooth in  $|-K_{\mathbb{F}_1}|$  and with respect to the  $-1$ -curve equals  $-3c^2\beta - 2c^3 + 3c^2 + 6c\beta$ , which for  $c = 2\beta$  (the Seshadri constant in this case), gives  $4\beta^2(6 - 7\beta)$ , showing that there exists no KEE metric when  $\beta \in (6/7, 1]$ . However,  $\mathbb{F}_1$  is *not* destabilized by any fiber of its natural projection to  $\mathbb{P}^1$  [25, Theorem 1.3]. We now show that  $\mathbb{F}_1$  is destabilized by a fiber after one flop, and this even holds for  $\beta \in (12/13, 1]$ .

To show this, it is most convenient to carry over the notation and assumptions of Section 4. Thus, we let  $S$  be  $\mathbb{P}^2$ ,  $C$  be a smooth cubic, and  $Z$  be a line. Then  $S' = \mathbb{F}_1$  is the blow-up of  $S$  at a point  $O_1 \in Z \cap C$ ,  $C'$  is an elliptic (anticanonical) curve, and  $Z'$  is a fiber of the natural projection  $\mathbb{F}_1 \rightarrow \mathbb{P}^1$ . In addition  $\mathcal{D}$  is  $C' \times \mathbb{P}^1$  and  $\mathcal{D}'$  is its proper transform with respect to the blow-up of  $Z' \subset S' \times \mathbb{P}^1$ . Let

$L'_\beta := -K_{S'} - (1 - \beta)C' = \beta C'$ , so  $\epsilon(S', L'_\beta, Z') = \beta$ . As  $L'_\beta \cdot Z' = 2\beta$  and  $Z'^2 = 0$ , Proposition 2.7 gives

$$F(\mathcal{X}', \mathcal{L}'_{\beta,c}, \mathcal{D}', \beta) = 6c\beta(2 - c). \tag{6.1}$$

Thus, if  $c < \beta$ , then  $F(\mathcal{X}', \mathcal{L}'_{\beta,c}, \mathcal{D}', \beta) > 0$ . On the other hand, we have

$$\tau(S', L'_\beta, Z') = \epsilon(S, L_\beta, Z) = 3\beta.$$

Thus, it follows from Lemma 4.7 that  $\hat{\mathcal{L}}_\beta$  is ample for every  $c \in (\beta, 3\beta)$ . By Proposition 4.9 and (6.1),

$$\begin{aligned} F(\hat{\mathcal{X}}', \hat{\mathcal{L}}'_{\beta,c}, \hat{\mathcal{D}}', \beta) &= F(\mathcal{X}', \mathcal{L}'_{\beta,c}, \mathcal{D}', \beta) - 2(\beta - c)^3 - 3(1 - \beta)(\beta - c)^2 \\ &= 6c\beta(2 - c) - 2(\beta - c)^3 - 3(1 - \beta)(\beta - c)^2. \end{aligned} \tag{6.2}$$

If  $c = 3\beta$ , then  $F(\hat{\mathcal{X}}', \hat{\mathcal{L}}'_{\beta,c}, \hat{\mathcal{D}}', \beta) = 24\beta^2 - 26\beta^3$ , which implies that  $F(\hat{\mathcal{X}}', \hat{\mathcal{L}}'_{\beta,c}, \hat{\mathcal{D}}', \beta) < 0$  (for some  $c \in (\beta, 3\beta)$ ) provided that  $\beta > \frac{12}{13}$ .

In fact, one can show that  $(\mathbb{F}_1, C', \beta)$  does not admit a KEE metric for  $\beta \in (\frac{4}{5}, 1]$  [30]. On the other hand,  $(\mathbb{F}_1, C', \beta)$  admits a KEE metric for  $\beta \in (0, \frac{3}{10})$ , and, moreover, if  $C'$  is a *general* curve in  $|-K_{\mathbb{F}_1}|$ , then  $(\mathbb{F}_1, C', \beta)$  admits a KEE metric for  $\beta \in (0, \frac{3}{7})$  [5, Corollary 1.16].

**Example 6.2** ( $\text{Bl}_{O_1, O_2} \mathbb{P}^2$ ). We take, as in the previous subsection,  $S = \mathbb{P}^2$ ,  $C$  a smooth cubic, and  $Z$  a line, but now blow-up two points  $O_1, O_2 \in C \cap Z$  to obtain  $S'$ , and let  $C', Z'$  be the proper transforms of  $C, Z$ , respectively. According to Panov-Ross [25, Example 7.6], the surface  $S'$  (with  $\beta = 1$ ) is slope stable. We will show that it is *not* flop-slope stable, and moreover this holds also for  $(S', C', \beta)$  with  $\beta \in (21/25, 1]$ . By comparison, Székelyhidi [30] constructed a destabilizing toric degeneration for  $\beta \in (\frac{7}{9}, 1]$  in the case when  $C'$  does not contain either of the points  $Z' \cap C'_1$  or  $Z' \cap C'_2$ , where  $C'_i$  are the exceptional curves of the blow-down map to  $\mathbb{P}^2$ . It is interesting to note that the value 21/25 also arises in the related smooth continuity method [29, Proposition 10] and [20, Example 2].

By Proposition 2.7, we have

$$F(\mathcal{X}', \mathcal{L}'_{\beta,c}, \mathcal{D}', \beta) = 3\beta c(2 - c) - c^2(2c - 3).$$

Here  $c < \epsilon(S', L'_\beta, Z') = \beta$ . Thus,  $F(\mathcal{X}', \mathcal{L}'_{\beta,c}, \mathcal{D}', \beta) > 0$  for every  $c \in (0, \beta)$  (*i.e.*, slope stable). On the other hand, we have

$$\tau(S', L'_\beta, Z') = \epsilon(S, L_\beta, Z) = 3\beta.$$

By Lemma 4.7, the divisor  $\hat{\mathcal{L}}'_\beta$  is ample for  $c \in (\beta, 3\beta)$ . Note that  $C'_i \cdot Z' = 1$  and as in (4.4) (see also (A.1))  $K_{\mathcal{X}'} \cdot C'_i = 0$ . Therefore,

$$\mathcal{L}'_{\beta,c} \cdot C_i = -\beta K_{S'} \cdot C'_i - cZ' \cdot C'_i = \beta - cZ' \cdot C'_i = \beta - c,$$



and by Proposition 4.9, one has

$$F(\hat{\mathcal{X}}', \hat{\mathcal{L}}'_\beta, \hat{\mathcal{D}}', \beta) = F(\mathcal{X}', \mathcal{L}'_{\beta,c}, \mathcal{D}', \beta) - 4(\beta - c)^3 - 6(\beta - c)^2(1 - \beta).$$

Plugging-in  $c = 3\beta$  yields

$$\begin{aligned} F(\hat{\mathcal{X}}', \hat{\mathcal{L}}'_\beta, \hat{\mathcal{D}}', \beta) &= 9\beta^2(2 - 3\beta) - 9\beta^2(6\beta - 3) + 32\beta^3 - 24\beta^2(1 - \beta) \\ &= \beta^2(21 - 25\beta) < 0, \end{aligned}$$

when  $\beta > \frac{21}{25}$ .


Note that  $(S', C', \beta)$  admits a KEE metric for  $\beta \in (0, \frac{3}{7})$ , and, moreover, if  $C'$  does not contain neither of the points  $Z \cap C'_1$  and  $Z \cap C'_2$ , then a KEE metric exists for  $\beta \in (0, \frac{1}{2})$  [5, Corollary 1.16].

### Appendix A. Simple flops

Let  $V$  be a smooth projective variety, and let

$$C \subset V$$

be a smooth rational curve such that its normal bundle in  $V$  is isomorphic to  $\mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1)$ . Then there exists a commutative diagram



such that the threefold  $\hat{V}$  is smooth, the threefold  $X$  has an isolated ordinary double point,  $\pi_C$  is a blow-up of the curve  $C$ ,  $\pi_{\hat{C}}$  is the contraction of the  $\pi_C$ -exceptional surface, let us call it  $E \cong \mathbb{P}^1 \times \mathbb{P}^1$ , to a smooth rational curve, let us call it  $\hat{C}$ . We define the map  $f$  by declaring the diagram to be commutative. This defines  $f$  as a birational map away from  $C$ . It is important in this construction that  $\pi_C \neq \pi_{\hat{C}}$ , so that the map  $f$  is not an isomorphism. Finally,  $c_C$  and  $c_{\hat{C}}$  are (small) contractions of the curves  $C$  and  $\hat{C}$ , respectively, to the isolated ordinary double point of  $X$ .

**Remark A.1.** The birational map  $f : V \dashrightarrow \hat{V}$  is called the *simple flop* of the curve  $C$ . Sometimes it is called an Atiyah flop [2]. Later it was explicitly introduced by Kulikov in [19, Section 4.2] as *perestroika I*.

Note that the normal bundle of  $\hat{C}$  in  $U$  is isomorphic to  $\mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1)$ . As in (4.4),

$$K_V.C = 0. \tag{A.1}$$

In fact, another way to see this equality is by noting that since  $c_C$  is an isomorphism away from codimension 2, then  $K_V \sim_{\mathbb{Q}} c_C^* K_V$ , and of course  $c_C^*(K_X).C = 0$  since  $c_C$  contracts  $C$ . Similarly,  $K_{\hat{V}}.\hat{C} = 0$  by construction. Moreover, we have  $E|_E$  is a divisor on  $E \cong \mathbb{P}^1 \times \mathbb{P}^1$  of bi-degree  $(-1, -1)$ . Furthermore, the morphism  $c_C \circ \pi_C = c_{\hat{C}} \circ \pi_{\hat{C}}$  is just the contraction of the surface  $E$  to the the isolated ordinary double point of  $\hat{X}$ , *i.e.*, its inverse map is the blow-up of this point.

**Remark A.2.** Note that in general  $\hat{V}$  is not necessarily projective. However, it is not hard to see that  $\hat{V}$  is projective in many cases, either by explicit construction or by using log MMP. In all our applications,  $\hat{V}$  is projective by construction, see Section 4.

Given an irreducible reduced Weyl divisor  $D$  on  $V$ , we denote by  $\hat{D}$  the unique divisor on  $\hat{V}$  such that

$$\hat{D} := \overline{f(D \setminus C)}.$$

By linearity, we extend the same notation to all  $\mathbb{R}$ -divisors on  $V$ . The following formula may be known, but we provide a proof since we were not able to find a reference for it.

**Lemma A.3.** *Let  $H_i, i = 1, 2, 3$ , be  $\mathbb{R}$ -divisors on  $V$ . Then,*

$$\hat{H}_1.\hat{H}_2.\hat{H}_3 = H_1.H_2.H_3 - (H_1.C)(H_2.C)(H_3.C).$$

*Proof.* Let  $\tilde{H}_1, \tilde{H}_2$  and  $\tilde{H}_3$  be the proper transforms of the divisors  $H_1, H_2$  and  $H_3$  on  $W$ , respectively. Recall that  $E = \mathbb{P}^1 \times \mathbb{P}^1$  denotes the exceptional divisor of  $\pi_C$  (and of  $\pi_{\hat{C}}$ ). Then,

$$\begin{cases} \tilde{H}_1 \sim_{\mathbb{R}} c_C^* H_1 - m_1 E \sim_{\mathbb{R}} c_{\hat{C}}^* \hat{H}_1 - \hat{m}_1 E \\ \tilde{H}_2 \sim_{\mathbb{R}} c_C^* H_2 - m_2 E \sim_{\mathbb{R}} c_{\hat{C}}^* \hat{H}_2 - \hat{m}_2 E \\ \tilde{H}_3 \sim_{\mathbb{R}} c_C^* H_3 - m_3 E \sim_{\mathbb{R}} c_{\hat{C}}^* \hat{H}_3 - \hat{m}_3 E, \end{cases}$$

for some real numbers  $m_i, \hat{m}_i$ . Put

$$r_i := H_i.C \quad \text{and} \quad \hat{r}_i := \hat{H}_i.\hat{C}.$$

Then each  $\tilde{H}_i|_E$  is a divisor (in  $\mathbb{P}^1 \times \mathbb{P}^1$ ) of bi-degree

$$(r_i + m_i, m_i) = (\hat{m}_i, \hat{r}_i + \hat{m}_i);$$

this is because  $E|_E = N_{E|V}$  is a line bundle of bi-degree  $(-1, -1)$ , while since  $c_C(E) = C$  and  $c_{\hat{C}}(E) = \hat{C}$ ,

$$c_C^* H_i|_E = H_i.c_C(E) \times (\text{fiber of projection of } \pi_C) = r_i \times (\text{bi-degree } (1,0) \text{ curve}), \tag{A.2}$$

and

$$c_{\hat{C}}^* \hat{H}_i|_E = \hat{H}_i \cdot c_{\hat{C}}(E) \times (\text{fiber of projection of } \pi_{\hat{C}}) = \hat{r}_i \times (\text{bi-degree } (0,1) \text{ curve}).$$

Thus,

$$\hat{m}_i = r_i + m_i, \quad \text{and } \hat{r}_i = -r_i. \quad (\text{A.3})$$

Now,  $E^3 = E|_E \cdot E|_E = c_1(\mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1))^2 = 2$ , and by (A.2),

$$c_C^* H_i \cdot E^2 = c_C^* H_i|_E \cdot E = -H_i|_E \cdot C = -r_i.$$

On the other hand,

$$c_C^* H_i \cdot c_C^* H_j \cdot E = c_C^* H_i|_E \cdot c_C^* H_j|_E = 0$$

since by (A.2),  $c_C^* H_i|_E$  and  $c_C^* H_j|_E$  are fibers of the same projection in  $\mathbb{P}^1 \times \mathbb{P}^1 = E$ . Altogether,

$$\begin{aligned} \tilde{H}_1 \cdot \tilde{H}_2 \cdot \tilde{H}_3 &= (c_C^* H_1 - m_1 E) \cdot (c_C^* H_2 - m_2 E) \cdot (c_C^* H_3 - m_3 E) \\ &= H_1 \cdot H_2 \cdot H_3 + (m_1 m_2 c_C^* H_3 + m_1 m_3 c_C^* H_2 + m_2 m_3 c_C^* H_1) \cdot E^2 - m_1 m_2 m_3 E^3 \\ &= H_1 \cdot H_2 \cdot H_3 - (m_1 m_2 r_3 + m_1 m_3 r_2 + m_2 m_3 r_1) - 2m_1 m_2 m_3. \end{aligned}$$

Similarly,

$$\begin{aligned} \tilde{\hat{H}}_1 \cdot \tilde{\hat{H}}_2 \cdot \tilde{\hat{H}}_3 &= (c_{\hat{C}}^* \hat{H}_1 - \hat{m}_1 E) \cdot (c_{\hat{C}}^* \hat{H}_2 - \hat{m}_2 E) \cdot (c_{\hat{C}}^* \hat{H}_3 - \hat{m}_3 E) \\ &= \hat{H}_1 \cdot \hat{H}_2 \cdot \hat{H}_3 + (\hat{m}_1 \hat{m}_2 c_{\hat{C}}^* \hat{H}_3 + \hat{m}_1 \hat{m}_3 c_{\hat{C}}^* \hat{H}_2 + \hat{m}_2 \hat{m}_3 c_{\hat{C}}^* \hat{H}_1) \cdot E^2 - \hat{m}_1 \hat{m}_2 \hat{m}_3 E^3 \\ &= \hat{H}_1 \cdot \hat{H}_2 \cdot \hat{H}_3 - (\hat{m}_1 \hat{m}_2 \hat{r}_3 + \hat{m}_1 \hat{m}_3 \hat{r}_2 + \hat{m}_2 \hat{m}_3 \hat{r}_1) - 2\hat{m}_1 \hat{m}_2 \hat{m}_3. \end{aligned}$$

Thus,

$$\begin{aligned} H_1 \cdot H_2 \cdot H_3 - m_1 m_2 r_1 - m_1 m_3 r_2 - m_2 m_3 r_1 - 2m_1 m_2 m_3 \\ = \hat{H}_1 \cdot \hat{H}_2 \cdot \hat{H}_3 - \hat{m}_1 \hat{m}_2 \hat{r}_3 - \hat{m}_1 \hat{m}_3 \hat{r}_2 - \hat{m}_2 \hat{m}_3 \hat{r}_1 - 2\hat{m}_1 \hat{m}_2 \hat{m}_3. \end{aligned}$$

By (A.3), this yields  $\hat{H}_1 \cdot \hat{H}_2 \cdot \hat{H}_3 = H_1 \cdot H_2 \cdot H_3 - r_1 r_2 r_3$ .  $\square$

## References

- [1] C. AREZZO, A. DELLA VEDOVA and G. LA NAVE, *Singularities and K-semistability*, Int. Math. Res. Not. IMRN **4** (2012), 849–869.
- [2] M. ATIYAH, *On analytic surfaces with double points*, Proc. Roy. Soc. Lond. Ser. A **247** (1958), 237–244.
- [3] R. BERMAN, *K-polystability of  $\mathbb{Q}$ -Fano varieties admitting Kähler–Einstein metrics*, Invent. Math. **203** (2016), 973–1025.

- [4] R. J. BERMAN, S. BOUCKSOM, P. EYSSIDIEUX, V. GUEDJ and A. ZERIAHI, *Kähler–Ricci flow and Ricci iteration on log Fano varieties*, J. Reine Angew. Math., doi: <https://doi.org/10.1515/crelle-2016-0033>
- [5] I. CHELTSOV and J. MARTINEZ-GARCIA, *Dynamic  $\alpha$ -invariants of Del Pezzo surfaces*, Int. Math. Res. Not. IMRN **2016** (2016), 2994–3028.
- [6] I. A. CHELTSOV and Y. A. RUBINSTEIN, *Asymptotically log Fano varieties*, Adv. Math. **285** (2015), 1241–1300.
- [7] T. DARVAS and Y. A. RUBINSTEIN, *Tian’s properness conjectures and Finsler geometry of the space of Kähler metrics*, J. Amer. Math. Soc. **30** (2017), 347–387.
- [8] W.-Y. DING and G. TIAN, *Kähler–Einstein metrics and the generalized Futaki invariant*, Invent. Math. **110** (1992), 315–335.
- [9] S. K. DONALDSON, *Scalar curvature and stability of toric varieties*, J. Differential Geom. **62** (2002), 289–349.
- [10] S. K. DONALDSON, *Discussion of the Kähler–Einstein problem*, preprint, <http://www2.imperial.ac.uk/~skdona/KENOTES.PDF>
- [11] K. FUJITA, *On log  $K$ -stability for asymptotically log Fano varieties*, preprint, arxiv:1509:02808.
- [12] A. FUTAKI, *An obstruction to the existence of Einstein–Kähler metrics*, Invent. Math. **73** (1983), 437–443.
- [13] P. GRIFFITHS and J. HARRIS, “Principles of Algebraic Geometry”, Wiley Interscience, 1978.
- [14] H. GUENANCIA and M. PĂUN, *Conic singularities metrics with prescribed Ricci curvature: The case of general cone angles along normal crossing divisors*, J. Differential Geom. **103** (2016), 15–57.
- [15] R. HARTSHORNE, “Algebraic Geometry”, Springer, 1977.
- [16] J.-M. HWANG, H. KIM, Y. LEE and J. PARK, *Slopes of smooth curves on Fano manifolds*, Bull. Lond. Math. Soc. **43** (2011), 827–839.
- [17] V. ISKOVSKIKH and YU. PROKHOROV, “Fano Varieties”, Springer, 1999.
- [18] T. JEFFRES, R. MAZZEO and Y. A. RUBINSTEIN, *Kähler–Einstein metrics with edge singularities* (with an appendix by C. Li and Y.A. Rubinstein), Ann. of Math. **183** (2016), 95–176.
- [19] V. KULIKOV, *Degenerations of  $K3$  surfaces and Enriques surfaces*, Izv. Akad. Nauk SSSR Ser. Mat. **41** (1977), 1008–1042, 1199.
- [20] C. LI, *Greatest lower bounds on Ricci curvature for toric Fano manifolds*, Adv. Math. **226** (2011), 4921–4932.
- [21] C. LI and S. SUN, *Conical Kähler–Einstein metric revisited*, preprint, arxiv:1207.5011v2.
- [22] C. LI and C.-Y. XU, *Special test configuration and  $K$ -stability of Fano varieties*, Ann. of Math. **180** (2014), 197–232.
- [23] Y. MATSUSHIMA, *Sur la structure du groupe d’homéomorphismes analytiques d’une certaine variété kählérienne*, Nagoya Math. J. **11** (1957), 145–150.
- [24] Y. ODAKA, *A generalization of the Ross–Thomas slope theory*, Osaka J. Math. **50** (2013), 171–185.
- [25] D. PANOV and J. ROSS, *Slope stability and exceptional divisors of high genus*, Math. Ann. **343** (2009), 79–101.
- [26] E. PICARD, *De l’équation  $\Delta u = ke^u$  sur une surface de Riemann fermée*, Journ. de Math. (4) **9** (1893), 273–292.
- [27] J. ROSS and R. THOMAS, *An obstruction to the existence of constant scalar curvature Kähler metrics*, J. Differential Geom. **72** (2006), 429–466.
- [28] Y. A. RUBINSTEIN, *Smooth and singular Kähler–Einstein metrics*, In: “Geometric and Spectral Analysis”, P. Albin *et al.* (eds.), Contemp. Math. 630, Amer. Math. Soc. and Centre Recherches Mathématiques, 2014, 45–138.
- [29] G. SZÉKELYHIDI, *Greatest lower bounds on the Ricci curvature of Fano manifolds*, Compos. Math. **147** (2011), 319–331.

- [30] G. SZÉKELYHIDI, *A remark on conical Kähler-Einstein metrics*, Math. Res. Lett. **20** (2013), 581–590.
- [31] V. V. SHOKUROV, *A nonvanishing theorem*, Izv. Akad. Nauk SSSR Ser. Mat. **49** (1985), 635–651; Math. USSR-izv. **26** (1986), 591–604.
- [32] J. STOPPA, *K-stability of constant scalar curvature Kähler manifolds*, Adv. Math. **221** (2009), 1397–1408.
- [33] G. TIAN, *Kähler-Einstein metrics on algebraic manifolds*, In: “Transcendental Methods in Algebraic Geometry” (Cetraro 1994), Lecture Notes in Math., Vol. 1646, 1996, 143–185.
- [34] G. TIAN, *Kähler-Einstein metrics with positive scalar curvature*, Invent. Math. **130** (1997), 1–37.
- [35] G. TIAN, *Futaki invariant and CM polarization*, preprint, <http://bicmr.pku.edu.cn/tian/?ddownload=204>
- [36] M. TROYANOV, *Prescribing curvature on compact surfaces with conic singularities*, Trans. Amer. Math. Soc. **324** (1991), 793–821.
- [37] X.-W. WANG, *Height and GIT weight*, Math. Res. Lett. **19** (2012), 909–926.

James Clerk Maxwell Building  
School of Mathematics  
University of Edinburgh  
Peter Guthrie Tait Road  
King’s Building Campus  
Edinburgh EH9 3FD  
I.Cheltsov@ed.ac.uk

University of Maryland  
Department of Mathematics  
College Park  
MD 20742-4015, USA  
yanir@umd.edu