

On the structure of affine flat group schemes over discrete valuation rings, I

NGUYEN DAI DUONG, PHUNG HO HAI AND JOÃO PEDRO P. DOS SANTOS

Abstract. We study affine group schemes over a discrete valuation ring R using two techniques: Neron blowups and Tannakian categories. We employ the theory developed to define and study differential Galois groups of \mathcal{D} -modules on a scheme over a R . This throws light on how differential Galois groups of families degenerate.

Mathematics Subject Classification (2010): 14F10 (primary); 14L15 (secondary).

1. Introduction

The thoughts the reader is about to see in this text come from our will to understand “models” of group schemes defined over a DVR and their Tannakian interpretation. In fact, we set out to identify if a more general theory would be able to accommodate the following two examples. (For unexplained notation, see the end of this introduction.)

Example 1.1 ([10, Exp. VI_B, 13.3]). Let R be a discrete valuation ring with uniformiser π and quotient field K . Consider the pro-system of affine group schemes

$$\dots \longrightarrow \mathbf{G}_{a,R} \xrightarrow{\times\pi} \mathbf{G}_{a,R} \xrightarrow{\times\pi} \mathbf{G}_{a,R}, \quad (1.1)$$

which corresponds, on the level of rings, to the inductive system

$$R[x_0] \longrightarrow R[x_1] \longrightarrow \dots, \quad x_i \longmapsto \pi x_{i+1}. \quad (1.2)$$

The limit G of diagram (1.1) is a flat affine group scheme over R whose associated ring $R[G]$ is the colimit of (1.2): $\{P \in K[T] \mid P(0) \in R\}$. Another meaningful way

The research of the first and the second named authors is supported in part by Vietnam National Foundation for Science and Technology Development (NAFOSTED), grant number 101.04-2016.19.

Received September 4, 2015; accepted in revised form March 14, 2017.

Published online May 2018.

to express G is by writing it as a join of R -schemes:

$$G = \mathbf{G}_{a,K} \coprod_{e, \text{Spec } K} \text{Spec } R.$$

Note that the R -module $R[G]$, being isomorphic to $R \oplus K \oplus K \oplus \dots$ is *not* projective and that $G \otimes R_n$ is always the trivial group scheme over $R_n = R/(\pi^{n+1})$.

Example 1.2 (cf. [2, 3.2.1.5]). Let R be a discrete valuation ring with uniformiser π and quotient field K . Let $\mathbf{G}_{m,R} = \text{Spec } R[z, 1/z]$. If we set $x_0 = z - 1$ and $y_0 = 1/z - 1$, so that $R[z, 1/z] = R[x_0, y_0]/(x_0 + y_0 + x_0 y_0)$, co-multiplication is given by

$$\begin{aligned} x_0 &\longmapsto x_0 \otimes 1 + 1 \otimes x_0 + x_0 \otimes x_0 \\ y_0 &\longmapsto y_0 \otimes 1 + 1 \otimes y_0 + y_0 \otimes y_0. \end{aligned} \tag{1.3}$$

Now we write

$$G_n = \text{Spec } R[x_n, y_n]/(x_n + y_n + \pi^n x_n y_n),$$

and define co-multiplication by

$$\begin{aligned} x_n &\longmapsto x_n \otimes 1 + 1 \otimes x_n + \pi^n x_n \otimes x_n, \\ y_n &\longmapsto y_n \otimes 1 + 1 \otimes y_n + \pi^n y_n \otimes y_n. \end{aligned} \tag{1.4}$$

It is immediately verified that

$$\begin{aligned} x_{n-1} &\longmapsto \pi x_n \\ y_{n-1} &\longmapsto \pi y_n \end{aligned}$$

defines a morphism of R -algebras $R[G_{n-1}] \rightarrow R[G_n]$ which induces an isomorphism $K[G_{n-1}] \rightarrow K[G_n]$. This means that $R[G_n]$ is none other than the subring of $K[z, 1/z]$ generated by $\pi^{-n} x_0$ and $\pi^{-n} y_0$. It is also easily verified that the co-multiplication (1.4) on $R[G_n]$ is obtained from that in (1.3). We then arrive at the projective system of affine group schemes

$$\dots \longrightarrow G_{n+1} \longrightarrow G_n \longrightarrow G_{n-1} \longrightarrow \dots$$

whose limit G is considered in [2, 3.2.1.5]. Note that, as in Example 1.1,

$$R[G] = \{P \in K[z, 1/z] : \varepsilon P \in R\},$$

where ε is the co-identity $z \mapsto 1$. Again $G \otimes R_n = \text{Spec } R_n$ for all n .

Furthermore, in [2], the author shows that the group scheme G is a ‘‘differential Galois group’’; a fact which kindred our interest in building a heftier theory.

Our method is to put together the theory of Tannakian categories over a DVR [9] and the theory of Neron blowups [32]. Once this is done, two basic principles appear:

- P1.** Tannakian theory over a DVR meets much more frequently group schemes of infinite type. More technically, ‘‘Galois groups’’ or ‘‘images’’ can be of infinite type;

P2. The theory of Neron blowups serves to render these inconvenients more tractable.

We then focus on two main tasks:

T1. Understand to what extent **P2** is responsible for the difficulty in **P1**;

T2. Detect in concrete cases when the difficulty in **P1** is avoided.

Our findings concerning **T1**, respectively **T2**, is in Section 6, respectively Section 8. We now summarise the contents of the text in more detail.

In Section 2 we introduce and study some basic properties of the Neron blowup of a group scheme over a DVR along a closed subscheme of the special fibre. This is a central technique in studying group schemes over discrete valuation rings. The text relies heavily on the work of Waterhouse and Weisfeiler [32] and for most of the time (Subsections 2.1-2.4) we simply present their findings in our perspective so that further explanations become more effective. In doing so, we explain how one result from [32] fits comfortably in general mathematical culture (Theorem 2.4), we elaborate on a hint appearing in [32, Proposition 2.7], we give a more important role to the notion of standard sequence (Definition 2.12) by including it as part of the structure theorem on generic isomorphisms (Theorem 2.11) and we derive an expression for an arbitrary affine flat group scheme as a limit of flat *algebraic* ones (Theorem 2.17). On the other hand, Subsection 2.5 already starts the program mentioned in this introduction: we define automatic blowups of group schemes (Definition 2.18). This is an abstraction of Example 1.1 and Example 1.2, and fits into the theory of standard sequences. We then go on to nuance some basic properties of the automatic blowup.

In Section 3, we begin to put emphasis on the Tannakian approach and study how to produce faithful representations of Neron blowups. This is motivated by our will to study differential Galois groups. Indeed, differential Galois groups are the reflex in the Tannakian mirror of certain tensor categories *generated by one object*. Performing blowups modifies the group on one side and our task is to understand how to modify the generating object on the other. We begin by finding faithful representations of Neron blowups of the identity (Proposition 3.1) and then proceed to treat the general case (Proposition 3.5). To repeat, the constructions involved in these last two propositions are tailored to serve a situation where the objects under the lenses is the category of representations, so that manipulations should take place in there. Once this principle is abandoned, we can offer the reader some simpler results (Proposition 3.8 and Proposition 3.9).

Section 4 studies possible candidates for images of group schemes. For a morphism $\rho : G \rightarrow H$ between affine group schemes over a field k , it is well-known that the schematic image $\text{Im}(\rho)$ is a closed subgroup scheme of H and that the natural morphism $G \rightarrow \text{Im}(\rho)$ is faithfully flat [31, Chapter 14]. (This is a very pleasant feature of this theory.) Said differently, ρ can be written as the composition $\tau \circ \sigma$, with σ a closed embedding and τ faithfully flat. But, in the setting of group schemes over a DVR, the situation gets a bit more complicated since the aforementioned factorization ceases to exist in general. One is then led

to consider factorizations into three morphisms (Definition 4.1), which gives two “images”. This is not as simple as the case of affine group schemes over fields, but is not as complicated as that of affine schemes. In possession of these definitions, we then move on to study the behavior of these images under reduction modulo π . The structure then becomes more sophisticated and a third “image” appears, see diagram (4.1). While these “images” are of course related to each other, they need not be identical (Example 4.8). We put in evidence some of these relations in Corollary 4.6 and Proposition 4.7. The section ends with a direct translation of some of the results found in terms of the accompanying tensor categories, which are properly defined in Definition 4.9 and recognized in Proposition 4.10.

In Section 5 we introduce the Neron blowup along a formal closed flat subgroup scheme (Definition 5.6). This is a generalisation of the notion of automatic blowup introduced in Subsection 2.5 and is the beginning of the march towards understanding how Neron blowups help in controlling “Galois groups” of infinite type. This section works out some fundamental properties of the new concept and special attention should be paid to Theorem 5.14, which says that the standard sequence of the blowup along a formal closed flat subgroup is “constant.” This property is then isolated and studied from an abstract point of view in Section 6. There, the goal is simple: when is a standard sequence with “constant” centres the standard sequence of a blowup along a formal flat closed subgroup scheme? (The motivation for this question should be understood in the light of task **T1** mentioned earlier since standard sequences give rise to affine group schemes.) We give a partial answer, cf. Corollary 6.10.

In Section 7 we apply some of our previous results to differential Galois theory. (As we understand, the object of this theory is the study of \mathcal{O} -coherent \mathcal{D} -modules, where \mathcal{D} is the ring of all differential operators.) Here we make fundamental use of Tannakian categories over R . The upshot is that for a \mathcal{D} -module \mathcal{M} on a smooth scheme X over R , one has *two differential Galois groups* in sight: the full $\text{Gal}'(\mathcal{M})$ and the restricted $\text{Gal}(\mathcal{M})$ (see Definition 7.6 and Definition 7.7). This, of course, is a manifestation of the theory developed in Section 4 so that, accordingly, $\text{Gal}(\mathcal{M})$ is always of finite type over R , and $\text{Gal}'(\mathcal{M})$ can be quite big (see Example 7.11, where we reinterpret Example 1.2). Over the residue field k , we arrive at *three group schemes*: the reductions $\text{Gal}(\mathcal{M}) \otimes k$ and $\text{Gal}'(\mathcal{M}) \otimes k$, and the differential Galois group of the \mathcal{D} -module $\mathcal{M}|_{X_k}$, $\text{Gal}(\mathcal{M} \otimes k)$. A simple example where all these groups differ is given (Example 7.9).

In Section 8, we concentrate on the “geometric case” and suppose that R contains a copy of its residue field. We then present a result, Theorem 8.2, which moves in the direction of Bolibrukh’s theorem on complete integrability and isomonodromic deformations [25, Theorem 1.2]. It says that for inflated (also called completely integrable) \mathcal{D} -modules over proper and smooth ambient spaces, the aforementioned differential Galois groups coincide. This is of course an accomplishment of task **T2** introduced above. In this same section, and then again on Section 9, we also explain a number of technical results concerning the approximation of R by smooth subalgebras (stemming mainly from [16, IV₃, Section 8] and [3]) which

come to be necessary in order to circumvent the fact that R is not of finite type over the ground field.

The proof of Theorem 8.2 makes use of two other elaborate webs of ideas which were developed in [12, 27] and [33], and the objective of Section 9 is to review these works. In fact, the examination we offer of [27] and [33] (see Theorem 9.1) only touches upon the main output of these, the exactness of the homotopy sequence. On the other hand the reading we make of [12] is more incisive and as a result, we extract deeper consequences from the methods: compare Theorem 9.2 with its inspiration, [12, Theorem 5.10].

Notation, conventions and standard terminology

- (1) Throughout the text, R stands for a discrete valuation ring with quotient field K , residue field k , and uniformizer π . All the Hom-sets and tensor products, when not explicitly indicated, are understood to be taken over R . The quotient ring $R/(\pi^{n+1})$ is denoted by R_n ;
- (2) Given an object X over R (a scheme, a module, etc.), we sometimes find useful to write X_k instead of $X \otimes_R k$, X_K instead of $X \otimes_R K$, etc.;
- (3) To avoid repetitions, by a **group scheme** over some ring A , we understand an **affine group scheme** over A ;
- (4) The category of group schemes over a ring A will be denoted by (\mathbf{GSch}/A) ; the full subcategory whose objects are A -flat will be denoted by (\mathbf{FGSch}/A) ;
- (5) If V is a free R -module of finite rank, we write $\mathbf{GL}(V)$ for the general linear group scheme representing $A \mapsto \text{Aut}_A(V \otimes A)$. If $V = R^n$, then $\mathbf{GL}(V) = \mathbf{GL}_n$;
- (6) If G is a group scheme over R , we let $\text{Rep}_R(G)$ stand for the category of representations of G which are, as modules, of *finite type* over R . (We adopt Jantzen's and Waterhouse's definition of representation. See [19, Part I, 2.7 and 2.8, 29ff] and [31, 3.1-2, 21ff].);
- (7) The full subcategory $\text{Rep}_R(G)^o$ of $\text{Rep}_R(G)$ has for objects those V whose underlying R -module is free. If $V \in \text{Rep}_R(G)^o$, we abuse notation and make no graphical distinction between the homomorphism $G \rightarrow \mathbf{GL}(V)$ and the coaction $V \rightarrow V \otimes R[G]$;
- (8) An object V of $\text{Rep}_R(G)^o$ is said to be a faithful representation if the resulting morphism $G \rightarrow \mathbf{GL}(V)$ is a closed immersion. Similar conventions are in force for group schemes over k , R_n , etc. We admonish the reader that *this is not* the terminology of the authoritative [10], where a faithful action is decreed to be one having no kernel (see Definition 2.3.6.2 of Exposé I). Since the literature on the subject is scant and the definition of [10] permits some bizzareries (see Example 1.2), we allow ourselves to make such a modification;
- (9) For an affine scheme X over R , we let $R[X]$ stand for the ring $\mathcal{O}(X)$. More generally, if A is any R -algebra, we write $A[X]$ to denote $\mathcal{O}(X \otimes_R A)$;
- (10) If G is a group scheme over R , the coidentity or counit is the morphism $\varepsilon : R[G] \rightarrow R$ associated to the identity $\text{Spec } R \rightarrow G$. The augmentation ideal of $R[G]$ is the kernel of ε ;

- (11) A vector bundle on an locally noetherian scheme X is a locally free \mathcal{O}_X -module of finite rank;
- (12) If $M \subset N$ is an inclusion of R -modules, we write M^{sat} to stand for the saturation of M in N , i.e. $\cup_m (M : \pi^m)$.

ACKNOWLEDGEMENTS. We wish to thank the referee of this journal for an intelligent reading of the work. The third named author thanks Mark Spivakovsky for help with the theory of smoothing of morphisms.

2. Introducing Neron blowups

2.1. Definition and basic properties

Let $G \in (\mathbf{FGSch}/R)$. In what follows we regard $R[G]$ as a subring of $K[G]$.

Let H_0 be a closed subgroup of G_k defined by an ideal J . Note that, in this case, $\pi \in J$. The Neron blowup G' of G at H_0 (cf. [32, page 551] and [1, 2.1.2]) is the spectrum of the subring $R[G']$ of $K[G]$ generated by $R[G]$ and the elements of $\pi^{-1}J$. (It is obviously sufficient to adjoin elements $\pi^{-1}f$, where f runs over a system of generators of J .) The natural morphism $R[G] \rightarrow R[G']$ is a morphism of Hopf algebras over R as is readily verified. (See also [32, Proposition 1.1].) By construction, $G'_K \rightarrow G_K$ is an isomorphism and $G'_k \rightarrow G_k$ factors through H_0 . In fact we have:

Lemma 2.1 (Universal property, see [32, page 551]). *Let $\mathcal{G} \rightarrow G$ be a morphism in \mathbf{FGSch}/R such that $\mathcal{G}_k \rightarrow G_k$ factors through $H_0 \rightarrow G_k$. Then there exists a unique $\mathcal{G} \rightarrow G'$ rendering the diagram*

$$\begin{array}{ccc}
 \mathcal{G} & \longrightarrow & G' \\
 & \searrow & \downarrow \\
 & & G
 \end{array}$$

commutative.

Remark 2.2. It should be noted that this very elementary definition of blowup in fact relates to the usual notion as follows. Let $(\pi, a_1, \dots, a_r) \subset R[G]$ be an ideal cutting out on G_k a closed subgroup scheme H_0 . Write G' for its Neron blow-up. Then

$$\frac{R[G][\xi_1, \dots, \xi_r]}{(\pi \xi_1 - a_1, \dots, \pi \xi_r - a_r)^{\text{sat}}} \longrightarrow K[G], \quad \xi_i \longmapsto \frac{a_i}{\pi},$$

defines an isomorphism onto $R[G']$. As is well-known, in case $R[G]$ is noetherian, the ring on the left above is the ring of an affine piece of the blowup [20, Lemma 1.2(e), page 318].

2.2. A result stemming directly from the theory of blowups

We use this section to state a generalization of a result appearing in [32] and which is more transparently treated by using (nowadays) commonplace algebraic geometry. We begin by recalling the following important theorem, whose proof the reader can find in [7, Chapter X, Section 9, number 7].

Theorem 2.3. *Let $\mathcal{I} \subset \mathcal{O}_X$ be an ideal of the locally noetherian scheme X . Let $\psi : \tilde{X} \rightarrow X$ stand for the blowup of \mathcal{I} . Let $U = \text{Spec } A$ be an affine open of X where \mathcal{I} is generated by a regular A -sequence x_1, \dots, x_c . Then the U -scheme $\psi^{-1}(U)$ is isomorphic to*

$$\text{Proj} \frac{A[\xi_1, \dots, \xi_c]}{(\xi_i x_j - \xi_j x_i)}.$$

From this and [32, Theorem 1.5], we arrive at the following generalization of [32, Theorem 1.7]. (Recall from [16, IV₄, 17.12.1, page 85] that a closed immersion of smooth schemes is always regular.)

Theorem 2.4. *Let $G \in (\mathbf{FGSch}/R)$ be of finite type. Let $H_0 \subset G_k$ be a regularly immersed closed sub-group scheme. Write c for the codimension of H_0 in G_k . If $G' \rightarrow G$ stands for the Neron blowup of G at H_0 , then $G' \otimes k \rightarrow H_0$ is a smooth surjective morphism whose kernel is isomorphic to $\mathbf{G}_{a,k}^c$.*

Example 2.5. Let k be of positive characteristic p . Let $G = \mathbf{G}_{a,R}$ and denote the Neron blowup of $\alpha_p \subset G_k$ by $G' \rightarrow G$. Then $G'_k \rightarrow \alpha_p$ is a smooth morphism with kernel $\mathbf{G}_{a,k}$.

2.3. The action of the centre of the Neron blowup on the kernel

Let $G \in (\mathbf{FGSch}/R)$ be of finite type. Let $H_0 \subset G_k$ be a regularly immersed closed subgroup scheme. From Theorem 2.4 we know that $G'_k \rightarrow H_0$ is a faithfully flat morphism so that H_0 acts by group scheme automorphisms on the *right* of the kernel F . This is explained in [8, Chapter III, Section 6, number 1, 431ff], but is really a triviality: “lift and conjugate.” Since $F \simeq \mathbf{G}_{a,k}^c$, then one might ask if this action is linear and if it can be obtained from something simpler. The answer is Proposition 2.7 below, which is the theme of this section.

To grasp the meaning of Proposition 2.7, we need the notion of the *conormal* representation. Let \mathcal{G} be an affine group scheme of finite type over k and let $\mathcal{H} \subset \mathcal{G}$ be a closed subgroup cut out by the ideal $I \subset k[\mathcal{G}]$. We let \mathcal{H} act on \mathcal{G} on the *right* by conjugation: $g \cdot h = h^{-1}gh$. In this way, we obtain an \mathcal{H} -module structure on $k[\mathcal{G}]$. If $\mathfrak{a} \subset k[\mathcal{G}]$ stands for the augmentation ideal, it is easily seen that \mathfrak{a} , I and $\mathfrak{a}I$ are all sub- \mathcal{H} -modules of $k[\mathcal{G}]$. Note that $I/\mathfrak{a}I$ is a finite dimensional k -space.

Definition 2.6. The \mathcal{H} -module $I/\mathfrak{a}I$ is called the *co-normal representation*. It will be denoted in this text by $\nu : \mathcal{H} \rightarrow \mathbf{GL}(I/\mathfrak{a}I)$.

Obviously, $I/\mathfrak{a}I$ is simply the fibre at e of the conormal sheaf of the immersion $\mathcal{H} \subset \mathcal{G}$ [16, IV₄, page 5]. We invite the reader to explicate the relation between ν and the pertaining adjoint representations.

Proposition 2.7. *We maintain the setting of the first paragraph of this section. Then there exists an isomorphism of group schemes*

$$F \simeq \mathbf{G}_{a,k}^c \simeq \text{Spec } k[\xi_1, \dots, \xi_c]$$

such that the action of H_0 on F is determined by

$$\xi_j \mapsto \sum_{i=1}^c \xi_i \otimes v_{ij},$$

where $[v_{ij}] \in \text{GL}_c(k[H_0])$ defines the conormal representation of H_0 .

Proof. In the argument, we employ some useful notations:

- We write “ \times ” for the fibre product over R and note that if X and Y are k -schemes, then $X \times Y$ is canonically isomorphic to $X \times_{\text{Spec } k} Y$ [16, I, 3.2.4, page 105];
- If A and B are R -algebras and \mathfrak{A} is an ideal of A , we write $\langle \mathfrak{A} \otimes 1 \rangle$ for the ideal generated by the image of $\mathfrak{A} \otimes_R B \rightarrow A \otimes_R B$;
- If \mathfrak{p} is a prime ideal of a ring A and $I \subset \mathfrak{p}$ is a sub-ideal, we write $I_{\mathfrak{p}}$ for $I A_{\mathfrak{p}}$.

Let $\gamma : G \times G \rightarrow G$ stand for the morphism defined by $(x, g) \mapsto g^{-1}xg$; it produces a right action of G on itself. Letting $\gamma' : G' \times G' \rightarrow G'$ stand for the analogous action of G' on itself, we arrive at a commutative diagram

$$\begin{array}{ccc} G' \times G' & \xrightarrow{\gamma'} & G' \\ \downarrow & & \downarrow \\ G \times G & \xrightarrow{\gamma} & G. \end{array}$$

Let $V \subset G$ be any affine open subset whose intersection with H_0 is the source of a scheme-theoretic section to $G'_k \rightarrow H_0$:

$$\sigma : H_0 \cap V \longrightarrow G'_k.$$

(The existence of V is a consequence of the description of the blowup offered in Theorem 2.3.) By definition of the right action

$$\alpha : F \times H_0 \longrightarrow F$$

we have a commutative diagram

$$\begin{array}{ccc} G'_k \times G'_k & \xrightarrow{\gamma'_k} & G'_k \\ \uparrow & & \uparrow \\ F \times G'_k & \xrightarrow{\quad} & F \\ \uparrow \text{id} \times \sigma & \nearrow \alpha & \\ F \times (H_0 \cap V) & & \end{array} \tag{2.1}$$

Let n be a closed point of $H_0 \cap V$ and complete diagram (2.1) as follows:

$$\begin{array}{ccc}
 G'_k \times G'_k & \xrightarrow{\gamma'_k} & G'_k \\
 \uparrow & & \uparrow \\
 F \times G'_k & \xrightarrow{\quad} & F \\
 \text{id} \times \sigma \uparrow & \nearrow \alpha & \\
 F \times \text{Spec } k[H_0]_n & \longrightarrow & F \times (H_0 \cap V).
 \end{array} \tag{2.2}$$

We will show that the obvious composition

$$F \times \text{Spec } k[H_0]_n \longrightarrow F \times (H_0 \cap V) \xrightarrow{\alpha} F$$

coincides with the composition

$$F \times \text{Spec } k[H_0]_n \longrightarrow F \times (H_0 \cap V) \xrightarrow{\nu} F,$$

where ν is the conormal representation.

As usual, let $\varepsilon : R[G] \rightarrow R$ be the co-identity and write \mathfrak{a} for its kernel. Write \mathfrak{m} for the ideal (π, \mathfrak{a}) – it corresponds to the identity on the closed fibre G_k – and note that ε extends to $R[G]_{\mathfrak{m}}$. It goes without saying that the kernel of $\varepsilon : R[G]_{\mathfrak{m}} \rightarrow R$ is just $\mathfrak{a}R[G]_{\mathfrak{m}}$.

Lemma 2.8. *There exist functions $x_{1,\mathfrak{m}}, \dots, x_{c,\mathfrak{m}} \in R[G]_{\mathfrak{m}}$ such that*

- (1) For each j , $\varepsilon(x_{j,\mathfrak{m}}) = 0$;
- (2) The sequence $\pi, x_{1,\mathfrak{m}}, \dots, x_{c,\mathfrak{m}}$ is regular and generates the ideal of H_0 at $R[G]_{\mathfrak{m}}$.

Proof. Let $\pi, x_{1,\mathfrak{m}}^*, \dots, x_{c,\mathfrak{m}}^*$ be a regular sequence generating the ideal of H_0 at \mathfrak{m} . We can always write $x_{j,\mathfrak{m}}^* = c_j + x_{j,\mathfrak{m}}$, with $c_j \in R$ and $x_{j,\mathfrak{m}} \in \mathfrak{a}R[G]_{\mathfrak{m}}$. Moreover, we know that $\varepsilon(x_{j,\mathfrak{m}}^*) \equiv 0 \pmod{\pi}$, since $x_{j,\mathfrak{m}}^* + (\pi) \in k[G]$ belongs to the ideal of H_0 . It then follows that $c_j \equiv 0 \pmod{\pi}$. Write $c_j^* := \pi^{-1}c_j$. Then $x_{j,\mathfrak{m}} = x_{j,\mathfrak{m}}^* - \pi c_j^*$, which means that we have an equality of ideals

$$(\pi, x_{1,\mathfrak{m}}, \dots, x_{c,\mathfrak{m}}) = (\pi, x_{1,\mathfrak{m}}^*, \dots, x_{c,\mathfrak{m}}^*).$$

The proof is finished by [16, IV₄, 16.9.5, page 47]. □

The morphism γ induces an arrow between local rings

$$\gamma^\# : R[G]_{\mathfrak{m}} \longrightarrow R[G]_{\mathfrak{m}} \otimes_R R[G]_n.$$

Since H_0 is a subgroup of G_k , if $J \subset R[G]$ stands for its ideal, it follows that

$$\gamma^\#(J_m) \subset \langle J_m \otimes 1 \rangle + \langle 1 \otimes J_n \rangle.$$

If $y_{1,n}, \dots, y_{c,n}$ denote elements of $R[G]_n$ such that $\pi, y_{1,n}, \dots, y_{c,n}$ forms a regular sequence generating J_n , we conclude that

$$\gamma^\#(x_{j,m}) = \sum_i a_{ij} x_{i,m} \otimes a'_{ij} + \sum_i b'_{ij} \otimes b_{ij} y_{i,n} + \pi s_j \tag{2.3}$$

for a_{ij}, b'_{ij} in $R[G]_m, a'_{ij}, b_{ij}$ in $R[G]_n$ and s_j in $R[G]_m \otimes_R R[G]_n$.

Since $\varepsilon(x_{j,m}) = 0$ (see Lemma 2.8), equation (2.3) gives

$$0 = \sum_i \varepsilon(b'_{ij}) \cdot b_{ij} y_{i,n} + \pi \cdot (\varepsilon \otimes_R \text{id}_{R[G]_n})(s_j),$$

which proves that

$$\pi \cdot (\varepsilon \otimes_R \text{id}_{R[G]_n})(s_j) \in (y_n).$$

Since the ideal (y_n) has no π -torsion, because the sequence $\{y_{1,n}, \dots, y_{c,n}, \pi\}$ is regular, it follows that

$$(\varepsilon \otimes_R \text{id}_{R[G]_n})(s_j) \in (y_n).$$

Hence, abusing notation and confusing \mathfrak{a} with $\mathfrak{a}R[G]_m$,

$$s_j \in \langle \mathfrak{a} \otimes_R 1 \rangle + \langle 1 \otimes_R (y_n) \rangle. \tag{2.4}$$

Since the sequence $\{\pi, x_{1,m}, \dots, x_{c,m}\}$ is regular, it follows that we have an isomorphism of $R[G]_m$ -algebras:

$$R[G']_m = \frac{R[G]_m[\xi]}{(\pi\xi - x_m)}. \tag{2.5}$$

(Here $R[G']_m$ stands for the ring obtained by inverting the elements of $R[G']$ in the image of $R[G] \setminus m$.) Now, using [16, IV₁, 15.2.4, page 21], there exists some affine neighbourhood V of \mathfrak{n} in G and functions $y \in R[V]$ inducing y_n such that $\{\pi, y_1, \dots, y_c\}$ is $R[V]$ -regular. Consequently, if $G'|_V$ stands for the inverse image of V in G' , we have

$$R[G'|_V] \simeq \frac{R[V][\eta]}{(\pi\eta - y)} \tag{2.6}$$

as $R[V]$ -algebras. In particular, $V \cap H$ is the source of a section $\sigma : H_0 \cap V \rightarrow G'_k$ defined on the level of rings by $\eta \mapsto \mathbf{0}$.

Now, $\gamma^\#$ fits into a diagram

$$\begin{array}{ccc} R[G']_m & \xrightarrow{\gamma^\#} & R[G']_m \otimes_R R[G']_n \\ \uparrow & & \uparrow \\ R[G]_m & \xrightarrow{\gamma^\#} & R[G]_m \otimes_R R[G]_n \end{array}$$

Then, from equations (2.3) and (2.5), we have

$$\pi \gamma^{\#}(\xi_j) = \pi \sum_i a_{ij} \xi_i \otimes a'_{ij} + \pi \sum_i b'_{ij} \otimes b_{ij} \eta_i + \pi s_j,$$


so that

$$\gamma^{\#}(\xi_j) = \sum_i a_{ij} \xi_i \otimes a'_{ij} + \sum_i b'_{ij} \otimes b_{ij} \eta_i + s_j. \tag{2.7}$$

Consequently, equations (2.4) and (2.6) show that

$$\gamma^{\#}(\xi_j) \equiv \sum_i \varepsilon(a_{ij}) \xi_i \otimes a'_{ij} \pmod{(\pi) + \langle \mathfrak{a} \otimes_R 1 \rangle + \langle 1 \otimes_R \eta \rangle}. \tag{2.8}$$

Now we write diagram (2.2) on the level of rings and complete:



It then follows from equation (2.8) that the image of ξ_j via

$$k[F] \xrightarrow{\alpha^{\#}} k[F] \otimes k[H_0 \cap V] \longrightarrow k[F] \otimes k[H_0]_n$$

is $\sum_i \varepsilon(a_{ij}) \xi_i \otimes a'_{ij}$. By definition, this is the co-normal representation v . □

Question 2.9. Is a regularly immersed closed affine algebraic subgroup scheme cut out by a regular sequence?

Remark 2.10. Proposition 2.7 is mentioned on [32, page 554].

2.4. The standard blowup sequence

As remarked in [32], Neron blowups allow us to decompose any morphism of group schemes which is an isomorphism on generic fibres.

Theorem 2.11 (cf. [32, Theorem 1.4]). *Let $\rho : \mathcal{G} \rightarrow G$ be an arrow of \mathbf{FGSch}/R which is an isomorphism on generic fibres. Write $\rho_0 = \rho$ and $G_0 = G$. Let $\rho_n : \mathcal{G} \rightarrow G_n$ be constructed and define G_{n+1} , respectively ρ_{n+1} , as the blowup of $\text{Im}(\rho_n \otimes k)$, respectively the natural morphism $\mathcal{G} \rightarrow G_{n+1}$. Then $\varprojlim_n \rho_n : \mathcal{G} \rightarrow \varprojlim_n G_n$ is an isomorphism.*

Proof. For the convenience of the reader and to avoid the hypothesis of finite generation present in [32], we summarize the proof.

We prove by induction that $\pi^{-n}R[G_0] \cap R[\mathcal{G}] \subset R[G_n]$. As the ideal of $R[G_0]$ cutting out $\text{Im}(\rho_0 \otimes k)$ is $R[G_0] \cap \pi R[\mathcal{G}]$, the case $n = 1$ is readily proved. We now assume that $\pi^{-n}R[G_0] \cap R[\mathcal{G}] \subset R[G_n]$. Let $f_0 \in R[G_0]$ be such that $\pi^{-n-1}f_0$ lies in $R[\mathcal{G}]$. Then $\pi^{-n}f_0$ belongs to $\pi R[\mathcal{G}]$. On the other hand, the inclusion $\pi^{-n}R[G_0] \cap R[\mathcal{G}] \subset R[G_n]$ forces $\pi^{-n}f_0$ to be in $R[G_n]$, so that $\pi^{-n}f_0$ belongs to $R[G_n] \cap \pi R[\mathcal{G}]$. The latter ideal cuts out $\text{Im}(\rho_n \otimes k)$, and therefore $\pi^{-1}\pi^{-n}f_0$ lies in $R[G_{n+1}]$.

Using the above inclusion, we arrive at $R[\mathcal{G}] \subset \cup R[G_n]$. Since each $R[G_n]$ is contained in $R[\mathcal{G}]$, the proof is finished. \square

Definition 2.12. The sequence of morphism described in the statement of Theorem 2.11 is called the *standard blowup sequence* associated to $\mathcal{G} \rightarrow G$. If context prevents any misunderstanding, we simply speak of the standard sequence of $\mathcal{G} \rightarrow G$.

The concept of standard sequence also allows the following point of view [32, page 552, Remarks].

Definition 2.13. A diagram

$$\cdots \longrightarrow G_{n+1} \xrightarrow{\rho_n} \cdots \xrightarrow{\rho_0} G_0$$

in \mathbf{FGSch}/R is a *standard sequence* if

- (1) Each arrow ρ_n is a blowup of some closed subgroup $B_n \subset G_n \otimes k$;
- (2) The induced morphism $B_{n+1} \rightarrow B_n$ is faithfully flat.

The overburdening of the term ‘‘standard sequence’’ caused by Definitions 2.12 and Definition 2.13 is ephemeral in view of:

Lemma 2.14. *Let*

$$\cdots \longrightarrow G_{n+1} \xrightarrow{\rho_n} \cdots \xrightarrow{\rho_0} G_0$$

be a standard sequence as in Definition 2.13. Then $\varprojlim_n G_n \rightarrow G_0$ is an isomorphism on generic fibres and its standard blowup sequence is the above diagram.

We shall apply Theorem 2.11 to give a description of affine group schemes over R , see Theorem 2.17 below. First we give a definition.

Definition 2.15. Let H' be a flat Hopf algebra over R . A *Hopf subalgebra* H of H' is an R -submodule equipped with a Hopf algebra structure such that the inclusion $H \rightarrow H'$ is a homomorphism of Hopf algebras. We say that H is *saturated* in H' if H'/H is flat as an R -module.

Remarks 2.16.

- (1) The image of a Hopf algebra homomorphism is a Hopf subalgebra (of the target);

- (2) One can always saturate a Hopf subalgebra. Adopting the notations explained at the end of Section 1 and Definition 2.15, we put

$$\Delta_{H^{\text{sat}}}(h) := \pi^{-m} \Delta_H(\pi^m h), \quad \text{if } \pi^m h \in H.$$

It can be extracted from the proof of [9, Lemma 3.1.2] that this is a coproduct for H^{sat} ;

- (3) Some authors define Hopf subalgebras as our saturated Hopf subalgebras. But Neron blowups justify our definition, see also Theorem 2.17.

Theorem 2.17. *Let G be a flat group scheme over R . Then we can present it as the limit of a pro-system of flat group schemes*

$$G := \varprojlim_i G_i,$$

in which all morphisms are faithfully flat and the generic fiber of each G_i is of finite type over K . Further, each G_i can be obtained from a flat group scheme of finite type by a standard sequence.

Proof. Consider $R[G]$ as a (right) regular representation of G (i.e., as comodule on itself by means of the coproduct). According to Serre [30], $R[G]$ is the union of its sub-representations, which are finite as R -modules. Let V be such a sub-representation (note that V is free over R). Then V induces a Hopf algebra homomorphism $R[\mathbf{GL}(V)] \rightarrow R[G]$, the image of which is a Hopf subalgebra of $R[G]$, denoted by $R[G(V)]$. Notice that V is a subset of $R[G(V)]$. Thus $R[G]$ is a union of its Hopf subalgebras $R[G(V)]$. One can take the G_i in Theorem to be the saturation of the $R[G(V)]$ as V runs in the set of finite sub-representations of $R[G]$.

Now by means of Theorem 2.11, the saturation $R[G(V)]^{\text{sat}}$ is obtained from $R[G(V)]$ by iterated Neron blowups. □

2.5. Study of a particular case

Examples 1.1 and 1.2 are a particular case of the following process. Let G be a flat group scheme over R .

Definition 2.18. The automatic blowup

$$N \longrightarrow G,$$

of the identity is the limit of the diagram

$$\cdots \longrightarrow G_n \longrightarrow \cdots \longrightarrow G_0,$$

where $G_0 = G$ and $G_{n+1} \rightarrow G_n$ is the Neron blowup of $\{e\} \subset G_n \otimes k$.

Remark 2.19. The construction put forth in the above definition will be generalized in Section 5 below.

Proposition 2.20. *Write \mathfrak{a} for the augmentation ideal of $R[G]$. Then, the algebra $R[N]$ of the automatic blowup of the identity is the $R[G]$ -subalgebra of $K[G]$ generated by $\cup_n \pi^{-n} \mathfrak{a}$. The latter union also generates the augmentation ideal of $R[N]$.*

Proof. We let G_n be as in Definition 2.18 and write \mathfrak{a}_n for the augmentation ideal of $R[G_n]$. It is easily proved that $(\pi^{-1} \mathfrak{a}_n) = \mathfrak{a}_{n+1}$. Let us now show by induction that $R[G_0][\pi^{-n} \mathfrak{a}_0] = R[G_n]$ and $\mathfrak{a}_n = (\pi^{-n} \mathfrak{a}_0)$. As there is nothing to be checked for $n = 0$, we assume that $R[G_0][\pi^{-n} \mathfrak{a}_0] = R[G_n]$ and $\mathfrak{a}_n = (\pi^{-n} \mathfrak{a}_0)$. Now, if $A \subset K[G]$ is any R -algebra and S is any subset of A generating an ideal \mathfrak{S} , it is clear that $A[\pi^{-1} \mathfrak{S}] = A[\pi^{-1} S]$. Hence, $R[G_{n+1}] = R[G_n][\pi^{-1} \mathfrak{a}_n]$ is just $R[G_n][\pi^{-n-1} \mathfrak{a}_0]$, which equals $R[G_0][\pi^{-n-1} \mathfrak{a}_0]$. This proves the first claim. As $\mathfrak{a}_{n+1} = (\pi^{-1} \mathfrak{a}_n)$ and $\mathfrak{a}_n = (\pi^{-n} \mathfrak{a}_0)$, we conclude that $\mathfrak{a}_{n+1} = (\pi^{-n-1} \mathfrak{a}_0)$, and the second claim follows as well. \square

Some noteworthy properties present in the examples remain valid. Others easily come forward. The next statement employs the notion of fibre product of rings, cf. [14, Section 1].

Corollary 2.21. *If $\varepsilon_K : K[G] \rightarrow K$ stands for the co-identity, then the first projection*

$$K[G] \times_{\varepsilon_K, K} R \longrightarrow K[G]$$


induces an isomorphism onto $R[N]$. Geometrically, we have an isomorphism of R -schemes

$$N \simeq G_K \coprod_{e, \text{Spec } K} \text{Spec } R.$$

Proof. It is clear that the first projection is an isomorphism onto the subalgebra $A := \{f \in K[G] : \varepsilon_K(f) \in R\}$. Equally clear is the inclusion $R[N] \subset A$, so that we are left with the verification of $A \subset R[N]$. If $f \in A$, then $f = c + f'$ with $c \in R$ and $\varepsilon_K(f') = 0$; this means that $f' = \pi^{-m} f''$ with f'' in the augmentation ideal of $R[G]$. By Proposition 2.20, $f' \in R[N]$ and $A \subset R[N]$ is verified.

The proof of the final statement follows directly from the first and from the fact that the functor Spec sends the cartesian diagram of rings in sight into a co-cartesian diagrams of schemes [14, Theorem 5.1, page 568]. \square

Corollary 2.22. *Let $\mathcal{G} \rightarrow G$ be a morphism of flat group schemes over R which induces an isomorphism on generic fibres. Then there exists a unique arrow $N \rightarrow G$ rendering commutative the diagram*



Said differently, $N \rightarrow G$ is an initial object in the category of flat group schemes $\mathcal{G} \rightarrow G$ over G which induce an isomorphism on generic fibres.

Proof. We need to prove that $R[\mathcal{G}] \subset K[G]$ is contained in $R[N]$. For that, let \mathfrak{a} and $\mathfrak{a}_{\mathcal{G}}$ stand respectively for the augmentation ideals of $R[G]$ and $R[\mathcal{G}]$. Writing an $f \in \mathfrak{a}_{\mathcal{G}}$ as $\pi^{-m} f'$ with $f' \in \mathfrak{a}$, we conclude that $f \in \cup_n \pi^{-n} \mathfrak{a}$. But then $f \in R[N]$ (Proposition 2.20) and we are done since $R \cdot 1 \oplus \mathfrak{a}_{\mathcal{G}} = R[\mathcal{G}]$. \square

Corollary 2.23. *Let $N \rightarrow G$ be the automatic blowup of the identity. Then, for each $n \geq 0$, the group scheme $N \otimes R_n$ over R_n is trivial, i.e. isomorphic to $\text{Spec } R_n$.*

Conversely, let $\rho : \mathcal{G} \rightarrow G$ be an arrow of \mathbf{FGSch}/R which is an isomorphism on generic fibres. If $\mathcal{G} \otimes R_n$ is trivial for all n , then ρ is the automatic blowup of the identity.

Proof. In this argument, we let \mathfrak{a} , \mathfrak{a}_N and $\mathfrak{a}_{\mathcal{G}}$ stand respectively for the augmentation ideals of $R[G]$, $R[N]$ and $R[\mathcal{G}]$. We know that $R[N] = \cup_n R[G][\pi^{-n} \mathfrak{a}]$ and that $\mathfrak{a}_N = (\cup_n \pi^{-n} \mathfrak{a})$. Hence, $\mathfrak{a}_N \subset (\pi^{n+1})$ for any n ; the first claim follows. On the other hand if $R_n[\mathcal{G}]$ is always trivial, then $\mathfrak{a}_{\mathcal{G}}$ is contained in (π^{n+1}) . Hence, $\mathfrak{a} \cdot R[\mathcal{G}] \subset (\pi^{n+1})$, which shows that $R[\mathcal{G}] \supset \cup_n R[G][\pi^{-n} \mathfrak{a}] = R[N]$. In view of Corollary 2.22, the inclusion $R[N] \supset R[\mathcal{G}]$ always holds and the proof is finished. \square

In fact, the second statement in the above corollary allows the following relevant amplification in case G is a closed subgroup scheme of $\mathbf{GL}_{n,R}$.

Corollary 2.24. *Let $V \in \text{Rep}_R(G)^o$ afford a faithful representation of G . Let $\rho : \mathcal{G} \rightarrow G$ be an arrow of \mathbf{FGSch}/R which is an isomorphism on generic fibres. If $V \otimes R_n$ is the trivial representation of $\mathcal{G} \otimes R_n$ for all n , then ρ is the automatic blowup of the identity.*

Proof. Let $[a_{ij}] \in \mathbf{GL}_r(R[G])$ be the matrix associated to V on some unspecified basis. Write $b_{ij} = a_{ij} - \delta_{ij}$. Since $V \otimes R_n$ is the trivial representation of $\mathcal{G} \otimes R_n$, we conclude that $\pi^{n+1} | b_{ij}$ in $R[\mathcal{G}]$ for all n . Now, the augmentation ideal \mathfrak{a} of $R[G]$ is generated by the functions b_{ij} , so that $\mathfrak{a} \cdot R[\mathcal{G}] \subset (\pi^{n+1})$. Hence, $R[\mathcal{G}] \supset \cup_n R[G][\pi^{-n} \mathfrak{a}] = R[N]$. Using Corollary 2.22, we have $R[N] = R[\mathcal{G}]$. \square

3. Faithful representations of Neron blowups

Let G be a flat group scheme over R . We assume furthermore that G is of finite type over R , so that the Neron blowup (see Section 2) of any closed subgroup of G_k is again of finite type. As such a group scheme, it admits a closed embedding into some $\mathbf{GL}_{r,R}$ (adapt the *proofs* in [31, 3.3]), or, according to the terminology at the end of section 1, it possesses a faithful representation. (The reader is again warned that this terminology *differs from* [10], where in Exposé I, Definition 2.3.6.2, a faithful action is one having a trivial kernel.) In this section we describe a means

to find faithful representations of Neron blowups. Of course, put this way, our task might seem pointless since it is sufficient to run a general argument. But our point of view is to produce faithful representations by performing linear algebraic constructions in Rep_R . This is most desirable when the category Rep_R is in fact one side of a Tannakian correspondence, see Example 7.9.

The principle is quite simple and we begin with a particular case:

$$G' \longrightarrow G$$

is the Neron blowup of the identity in the closed fibre. Let $V \in \text{Rep}_R(G)^o$ be a faithful representation of rank r ; since $G'_k \rightarrow G_k$ is the *trivial* morphism, we know that $\mathbf{1}^r \otimes k \simeq V \otimes k$ as representations of G' . We then obtain a diagram

$$\begin{array}{ccc} & & V \\ & & \downarrow \\ \mathbf{1}^r & \xrightarrow{\varphi} & V \otimes k \end{array}$$

in the category $\text{Rep}_R(G')$. (Here φ is the obvious morphism.) Let V' stand for its pull-back. In concrete terms,

$$V' = \{(\mathbf{v}, \mathbf{e}) \in V \oplus \mathbf{1}^r : \mathbf{v} \otimes 1 = \varphi(\mathbf{e})\}. \tag{3.1}$$

Proposition 3.1. *In the above setting, the representation V' of G' is faithful.*

The proof hinges on the following lemma, whose verification is omitted.

Lemma 3.2. *Let M and E be free R -modules. Let $\{\mathbf{m}_1, \dots, \mathbf{m}_r\}$ be a basis for M and $\{\mathbf{e}_1, \dots, \mathbf{e}_s\}$ be one for E . Let $q \leq \min\{r, s\}$ and define $\varphi : E \rightarrow M \otimes k$ by*

$$\varphi(\mathbf{e}_j) = \begin{cases} \mathbf{m}_j \otimes 1, & 1 \leq j \leq q \\ \mathbf{0}, & \text{otherwise.} \end{cases}$$

Then the R -module

$$\{(\mathbf{m}, \mathbf{e}) \in M \oplus E : \mathbf{m} \otimes 1 = \varphi(\mathbf{e})\}$$

is free on the basis

$$\begin{aligned} &(\pi \mathbf{m}_1, \mathbf{0}), \dots, (\pi \mathbf{m}_r, \mathbf{0}) \\ &(\mathbf{m}_1, \mathbf{e}_1), \dots, (\mathbf{m}_q, \mathbf{e}_q) \\ &(\mathbf{0}, \mathbf{e}_{q+1}), \dots, (\mathbf{0}, \mathbf{e}_s). \end{aligned} \quad \square$$

Proof of Proposition 3.1. This is plain linear algebra. Let $\{\mathbf{v}_1, \dots, \mathbf{v}_r\}$ be a basis of V and let $a_{ij} \in R[G]$ stand for the matrix coefficients inducing the representation of G . Let $\varepsilon : R[G] \rightarrow R$ stand for the counit; by definition of a representation we have $\varepsilon(a_{ij}) = \delta_{ij}$, so that

$$a_{ij} = \delta_{ij} \cdot 1 + b_{ij}, \tag{3.2}$$

with $b_{ij} \in \text{Ker } \varepsilon$. By definition of G' , there exist $b'_{ij} \in R[G']$ such that

$$\pi b'_{ij} = b_{ij}. \tag{3.3}$$

Let $\{\mathbf{e}_1, \dots, \mathbf{e}_r\}$ be an ordered basis of $\mathbf{1}^r$ which is sent by φ to $\{\mathbf{v}_1 \otimes 1, \dots, \mathbf{v}_r \otimes 1\}$. According to Lemma 3.2, V' is free on

$$\{\pi \mathbf{v}_1, \dots, \pi \mathbf{v}_r, \mathbf{v}_1 + \mathbf{e}_1, \dots, \mathbf{v}_r + \mathbf{e}_r\}.$$

Employing this basis and the equations (3.2) and (3.3), we conclude that the matrix defining the representation of G' on V' is

$$\left[\begin{array}{c|c} a_{ij} & b'_{ij} \\ \hline 0 & \delta_{ij} \end{array} \right].$$

Since the functions a_{ij} together with $1/\det[a_{ij}]$ generate the R -algebra $R[G]$, the functions b_{ij} generate the ideal $\text{Ker } \varepsilon$; by definition the functions a_{ij} , $1/\det[a_{ij}]$ and $b'_{ij} = \pi^{-1}b_{ij}$ generate $R[G']$ and the proof is finished. \square

Before moving to the search for a faithful representation of a general Neron blowup $G' \rightarrow G$, we record some valuable properties of V' and give an example.

Corollary 3.3. *We maintain the above notation.*

- (1) *The representation V' is an extension of $\mathbf{1}^r$ by V . In particular, V is a sub-representation of V' ;*
- (2) *Let ξ be the evident extension class in $\text{Ext}_{G'}(V \otimes k, V)$. Then the class of V' in $\text{Ext}_{G'}(\mathbf{1}^r, V)$ is simply the image of ξ under the morphism induced by $\mathbf{1}^r \rightarrow V \otimes k$;*
- (3) *The cokernel of the injection $V' \rightarrow V \oplus \mathbf{1}^r$ is annihilated by π ;*
- (4) *The natural arrow $V' \otimes K \rightarrow (V \oplus \mathbf{1}^r) \otimes K$ is an isomorphism of representations of $G' \otimes K$;*
- (5) *Let $\{\mathbf{v}_1, \dots, \mathbf{v}_r\}$ and $\{\mathbf{e}_1, \dots, \mathbf{e}_r\}$ be respectively ordered bases of V and $\mathbf{1}^r$ as constructed above. Write $\rho_K : G_K \rightarrow \mathbf{GL}_{2r, K}$ for the homomorphism associated to the representation $(V \oplus \mathbf{1}^r) \otimes K$ by means of these bases. Then, if*

$$\beta = \left[\begin{array}{c|c} \pi & \\ \hline & \text{Id}_r \\ \hline & \pi \\ \hline & \text{Id}_r \end{array} \right],$$

it follows that G' is the closure of the image of $\beta^{-1} \cdot \rho_K \cdot \beta$ in $\mathbf{GL}_{2r, R}$;

- (6) *As representations of G' , V' is a sub-representation of $V \oplus \mathbf{1}^r$, and V is a sub-representation of V' .*

Proof. The only statements which need comment are (2) and (5). Concerning (2), we only need to inform the reader that we follow [18, page 87] in constructing $\text{Ext}(V \otimes k, V) \rightarrow \text{Ext}(\mathbf{1}', V)$. Concerning (5), we produce the following justifications. Let $[a_{ij}] \in \text{GL}_r(R[G])$ stand for the matrix associated to the representation V and the basis $\{v_1, \dots, v_r\}$. We observe that

$$\beta^{-1} \cdot \rho_K \cdot \beta = \left[\begin{array}{c|c} a_{ij} & (a_{ij} - \delta_{ij})/\pi \\ \hline O & \delta_{ij} \end{array} \right].$$

Now the above matrix is also the matrix of a faithful representation $G' \rightarrow \mathbf{GL}_{2r,R}$, see the proof of Proposition 3.1. Since $G' \otimes K \xrightarrow{\sim} G \otimes K$, the closure of the image of $\beta^{-1} \cdot \rho_K \cdot \beta$ is the same as the closure of the image of $G' \otimes K$ by $\beta^{-1} \cdot \rho_K \cdot \beta$. This is G' , since two flat and closed subschemes having the same generic fibre must be equal. □

Example 3.4. Let G' be the Neron blowup of $\mathbf{G}_{m,R} = \text{Spec } R[u, 1/u]$ at the origin in $\mathbf{G}_{m,k}$. If V is the obvious representation of $\mathbf{G}_{m,R}$ (corresponding to id), then V' is the representation of G' corresponding to the matrix

$$\begin{bmatrix} \pi & 1 \\ 0 & 1 \end{bmatrix}^{-1} \cdot \begin{bmatrix} u & \\ & 1 \end{bmatrix} \cdot \begin{bmatrix} \pi & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} u & (u-1)/\pi \\ 0 & 1 \end{bmatrix}.$$

It follows that G' is

$$\left\{ \begin{bmatrix} u & v \\ 0 & 1 \end{bmatrix} : \pi v + 1 = u \right\}.$$

Let us now assume that

$$G' \longrightarrow G,$$

is the Neron blowup of a closed subgroup scheme $H_0 \subset G \otimes k$. The idea to construct a faithful representation of G' by means of a faithful representation of G is to express H_0 as the stabilizer of some line. This means that the problem is divided into two:

Step 1. Express H_0 as a stabilizer.

Step 2. Construct a faithful representation of the Neron blow of a stabilizer in the special fibre.

We start by discussing Step 2 and presenting our findings as Proposition 3.5. Step 1 follows from standard material in the theory of affine group schemes and is explained in Lemma 3.7 below.

Let $V \in \text{Rep}_R(G)^\circ$ be faithful and let $v \otimes 1 \in V \otimes k$ be a nonzero vector. Let $H_0 \subset G_k$ stand for the stabilizer of the line

$$\ell := v \otimes k.$$

Clearly, it is possible to find an ordered basis $\{\mathbf{v}_1, \dots, \mathbf{v}_r\}$ of V such that $\mathbf{v}_1 = \mathbf{v}$. Let $[a_{ij}] \in \text{GL}_r(R[G])$ be the matrix of coefficients associated to $\{\mathbf{v}_1, \dots, \mathbf{v}_r\}$. It then follows that the ideal cutting out H_0 in $R[G]$ is simply

$$(\pi, a_{21}, \dots, a_{r1}).$$

For future use, we write

$$a_{21} = \pi a'_{21}, \dots, a_{r1} = \pi a'_{r1} \tag{3.4}$$

for functions $a'_{21}, \dots, a'_{r1} \in R[G']$.

The line ℓ is now fixed by G'_k and we obtain a character $G'_k \rightarrow \mathbf{G}_{m,k}$ (defined by the group-like element $a_{11} + (\pi)$ in $k[G']$). In order to follow the method developed to treat the particular case, we should find a representation $L \in \text{Rep}_R(G')^o$ lifting ℓ . So here we need to modify the argument since there is no reason for L to exist. Let $E \in \text{Rep}_R(G')^o$ be the source of a surjection

$$\varphi : E \longrightarrow \ell$$

in $\text{Rep}_R(G')$. (That E exists is proved in [30, Proposition 3, page 41]). We fix an element $\mathbf{e}_1 \in E$ above $\mathbf{v} \otimes 1 \in \ell$. A bit of common sense shows that there exists elements $\mathbf{e}_2, \dots, \mathbf{e}_s \in \text{Ker } \varphi$ which together with \mathbf{e}_1 form a basis of E . Let $[b_{ij}] \in \text{GL}_s(R[G'])$ be the matrix associated to the representation E and the ordered basis $\{\mathbf{e}_1, \dots, \mathbf{e}_s\}$. Note that in this case we have

$$\begin{aligned} b_{11} &= a_{11} + \pi c_{11} \\ b_{12} &= \pi c_{12} \\ \dots &= \dots \\ b_{1s} &= \pi c_{1s}, \end{aligned} \tag{3.5}$$

since $\varphi(\mathbf{e}_1) = \mathbf{v}_1 \otimes 1$ and the subspace of E_k generated by $\mathbf{e}_2 \otimes 1, \dots, \mathbf{e}_s \otimes 1$ is stable under G'_k . We then consider the pull-back diagram

$$\begin{array}{ccc} V' & \longrightarrow & V \\ \downarrow & \square & \downarrow \\ E & \xrightarrow{\varphi} & V_k, \end{array}$$

so that V' is naturally a sub- G' -module of $V \oplus E$. From Lemma 3.2 we know that

$$\begin{aligned} &(\pi \mathbf{v}_1, \mathbf{0}), \dots, (\pi \mathbf{v}_r, \mathbf{0}) \\ &(\mathbf{v}_1, \mathbf{e}_1), \\ &(\mathbf{0}, \mathbf{e}_2), \dots, (\mathbf{0}, \mathbf{e}_s) \end{aligned}$$

is a basis of V' . A simple computation using equations (3.4) and (3.5) shows that the matrix of the representation of G' on V' associated to this basis is

$$\left[\begin{array}{ccc|ccc} a_{11} & \cdots & a_{1r} & -c_{11} & -c_{12} & \cdots & -c_{1s} \\ \vdots & \ddots & \vdots & a'_{21} & 0 & \cdots & 0 \\ \vdots & & \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{r1} & \cdots & a_{rr} & a'_{r1} & 0 & \cdots & 0 \\ \hline 0 & \cdots & 0 & b_{11} & b_{12} & \cdots & b_{1s} \\ \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & b_{s1} & b_{s2} & \cdots & b_{ss} \end{array} \right].$$

Since the a_{ij} together with $1/\det [a_{ij}]$ generate $R[G]$, and a'_{21}, \dots, a'_{r1} generate $R[G']$ over $R[G]$, we conclude that V' is a faithful representation of G' . We have then proved the following.

Proposition 3.5. *Let $V \in \text{Rep}_R(G)$ be faithful and let $\ell \subset V_k$ be a line whose stabilizer is denoted by $H_0 \subset G_k$. Let G' be the Neron blowup of G at H_0 and let $\varphi : E \rightarrow \ell$ be a surjective morphism in $\text{Rep}_R(G')$ such that the R -module E is free. Then $V' := V \times_{V_k} E$ is a faithful representation of G' . \square*

Corollary 3.6. *Let $V \in \text{Rep}_R(G)^\circ$ be faithful and let $\mathbf{v} \otimes 1 \in V_k \setminus 0$ be a vector whose stabilizer is denoted by $H_0 \subset G_k$. (Note that H_0 is smaller than the stabilizer of the line!) Let G' be the Neron blowup of G at H_0 and note that there is an obvious G' -equivariant morphism $\mathbf{1} \rightarrow V_k$ whose image is the line $\mathbf{v} \otimes k$. Then $V' := V \times_{V_k} \mathbf{1}$ is a faithful representation of G' . Moreover, if $[a_{ij}] \in \text{GL}_r(R[G])$ is the matrix associated to some basis $\{\mathbf{v}_1, \dots, \mathbf{v}_r\}$ with $\mathbf{v}_1 = \mathbf{v}$, then the matrix of the representation V' of G' is*

$$\left[\begin{array}{ccc|c} a_{11} & \cdots & a_{1r} & \alpha_{11} \\ \vdots & \ddots & \vdots & a'_{21} \\ \vdots & & \vdots & \vdots \\ a_{r1} & \cdots & a_{rr} & a'_{r1} \\ \hline 0 & \cdots & 0 & 1 \end{array} \right],$$

where $\pi\alpha_{11} = a_{11} - 1$ and $\pi a'_{i1} = a_{i1}$ for each $i \geq 2$. \square

To address Step 1 we only need the following:

Lemma 3.7. *Let $H_0 \subset G_k$ be a closed subgroup scheme of G . Then there exists a faithful representation V of G and a line $\ell \subset V_k$ whose stabilizer is exactly H_0 .*

Proof. By [31, 16.1, Corollary] there exists $\rho : G_k \rightarrow \mathbf{GL}_{r,k}$ and a line $\ell \subset k^r$ whose stabilizer is H_0 . Using [30, Proposition 3, page 41] we can find $W \in \text{Rep}_R(G)^\circ$ together with a G_k -equivariant injection $k^r \rightarrow W_k$. Obviously, H_0 is still the stabilizer of the line $\ell \subset W_k$. Replacing W by $V = W \oplus F$, with F faithful, we are done. \square

Let us now criticize Proposition 3.5. Finding explicitly the “covering” representation $\varphi : E \rightarrow \ell$ is by no means a simple task. (But in many important cases, the character of $G'_k \rightarrow \mathbf{G}_{m,k}$ associated to ℓ will be a reduction of a character $G' \rightarrow \mathbf{G}_{m,R}$.) Also, if we abandon the need to construct the faithful representation V' by means of linear algebra in the abelian category $\text{Rep}_R(G')$, a more efficient path is:

Proposition 3.8. *Let V be a free R -module on the basis $\mathbf{v}_1, \dots, \mathbf{v}_r$ affording a faithful representation of G . Let $H_0 \subset G_k$ be the stabilizer of the line spanned by $\mathbf{v}_1 \otimes 1$ in V_k and denote by $G' \rightarrow G$ the Neron blowup of H_0 . Write $[a_{ij}] \in \text{GL}_r(R[G])$ for the matrix associated to the representation of G on V , and let a'_{21}, \dots, a'_{r1} be functions of $R[G']$ which, when multiplied by π , become respectively a_{21}, \dots, a_{r1} . Then the following claims are true.*

- (1) *The R -submodule of V_K freely generated by $\mathbf{v}_1 \otimes \pi^{-1}, \mathbf{v}_2 \otimes 1, \dots, \mathbf{v}_r \otimes 1$ has the structure of an $R[G']$ -comodule. Its associated matrix is*

$$\left[\begin{array}{c|ccc} a_{11} & \pi a_{12} & \cdots & \pi a_{1r} \\ \hline a'_{21} & a_{22} & \cdots & a_{2r} \\ \vdots & \vdots & \ddots & \vdots \\ a'_{r1} & a_{r2} & \cdots & a_{rr} \end{array} \right];$$

- (2) *If V' stands for the representation of G' considered in the previous item, then $V \oplus V'$ is a faithful representation of G' .*

Proof. Once one knows what to look for, the proof is a triviality. □

Provided the centre of the Neron blowup has certain particular features, the point of view of [28] provides another means to construct faithful representations.

Proposition 3.9. *Let $H \subset G$ be an R -flat, normal subgroup scheme and write A for the quotient group scheme. Denote by G' and A' the Neron blowups of G at H_k and of A at $\{e\} \subset A_k$, respectively. Let $G' \rightarrow A'$ be the morphism obtained by the “universal property” (Lemma 2.1). If $\rho : G \rightarrow \mathbf{GL}_r$ and $\sigma : A' \rightarrow \mathbf{GL}_s$ are faithful representations, then $\rho \oplus \sigma$ is a faithful representation of G' .*

Proof. Here are the properties of the quotient which we shall need: $R[A]$ is a subring of $R[G]$ and if \mathfrak{a}_A stands for the augmentation ideal of $R[A]$, then $\mathfrak{a}_A \cdot R[G]$ cuts out H . Hence, if a_1, \dots, a_m are generators of \mathfrak{a}_A , it follows that

$$R[G'] = R[G] \left[\pi^{-1}a_1, \dots, \pi^{-1}a_m \right].$$

We now verify that

$$\rho^* \otimes \sigma^* : R[\mathbf{GL}_r] \otimes R[\mathbf{GL}_s] \longrightarrow R[G']$$

is a surjection. By definition of ρ , $R[G] \subset \text{Im}(\rho^* \otimes \sigma^*)$. By definition of σ , the elements $\pi^{-1}a_i$ belong to $\text{Im}(\sigma^*)$. This finishes the proof. □

4. Images of morphisms between flat group schemes

Let $\rho : \Pi \rightarrow G$ be a morphism in \mathbf{FGSch}/R . There are two natural ways of defining “images” of ρ .

Definition 4.1 (The diptych). Define Ψ_ρ as the group scheme whose Hopf algebra is the image of $R[G]$ in $R[\Pi]$. Define $R[\Psi'_\rho]$ as the saturation of the latter inside $R[\Pi]$. The obvious commutative diagram

$$\begin{array}{ccc}
 \Psi'_\rho & \longrightarrow & \Psi_\rho \\
 \uparrow & & \downarrow \\
 \Pi & \xrightarrow{\rho} & G
 \end{array}$$

is called the *diptych* of ρ .

Implicit in the above definition is the fact that $R[\Psi'_\rho]$ is a Hopf algebra. This can be extracted from the proof of [9, Lemma 3.1.2] (see Remarks 2.16). Another relevant fact, whose proof the reader can find in [9, Theorem 4.1.1], is the following.

Theorem 4.2. *The morphism $\Pi \rightarrow \Psi'_\rho$ is faithfully flat.*

A basic property of Ψ_ρ is as follows.

Lemma 4.3. *The group scheme Ψ_ρ is the closure of the image of $\Pi_K \rightarrow G_K$ in G .*

Other useful facts having simple proofs are collected in the next lemma.

Lemma 4.4. *Let $\rho : \Pi \rightarrow G$ be as before and consider a factorization of ρ in the category \mathbf{FGSch}/R :*

$$\Pi \longrightarrow G' \longrightarrow G.$$

- (1) *If $\Pi \rightarrow G'$ faithfully flat, then there exists a unique dotted arrow rendering the diagram*

$$\begin{array}{ccccc}
 & & \Psi'_\rho & & \\
 & \nearrow & \uparrow & \searrow & \\
 \Pi & \longrightarrow & G' & \longrightarrow & G,
 \end{array}$$

commutative;

- (2) *If $G' \rightarrow G$ is a closed embedding, then there exists a unique dotted arrow*

$$\begin{array}{ccccc}
 & & \Psi_\rho & & \\
 & \nearrow & \uparrow & \searrow & \\
 \Pi & \longrightarrow & G' & \longrightarrow & G,
 \end{array}$$

rendering this diagram commutative;

- (3) If $G' \rightarrow G$ is a closed embedding, and the arrow $\Psi_\rho \rightarrow G'$ produced in (2) induces an isomorphism on generic fibres, then $\Psi_\rho \rightarrow G'$ is an isomorphism;
- (4) If $\Pi \rightarrow G'$ is faithfully flat and the arrow $G' \rightarrow \Psi'_\rho$ produced in (1) induces an isomorphism on generic fibres, then $G' \rightarrow \Psi'_\rho$ is an isomorphism.

Lemma 4.5. *If $\Psi'_{\rho,k} \rightarrow \Psi_{\rho,k}$ is faithfully flat, then $\Psi'_\rho \rightarrow \Psi_\rho$ is an isomorphism.*

Proof. By construction, $R[\Psi_\rho] \rightarrow R[\Psi'_\rho]$ is injective. If $k[\Psi_\rho] \rightarrow k[\Psi'_\rho]$ is also injective, which follows from the assumption, then $R[\Psi_\rho] \subset R[\Psi'_\rho]$ is saturated. The equality $K[\Psi_\rho] = K[\Psi'_\rho]$ then finishes the proof. \square

Over the residue field k , there is yet another interesting group scheme in sight: the image of ρ_k . We then have the *tritych* of ρ , which is the commutative diagram

$$\begin{array}{ccc}
 \Psi'_{\rho,k} & \xrightarrow{\quad} & \Psi_{\rho,k} \\
 \uparrow & \searrow & \nearrow \\
 & \text{Im}(\rho_k) & \\
 \uparrow & \nearrow & \searrow \\
 \Pi_k & \xrightarrow{\rho_k} & G_k
 \end{array}
 \quad
 \begin{array}{l}
 \hookrightarrow = \text{closed immersion} \\
 \twoheadrightarrow = \text{faithfully flat.}
 \end{array}
 \tag{4.1}$$

Together with Lemma 4.5, diagram (4.1) proves the following:

Corollary 4.6. *The following claims are true.*

- i) *If $\text{Im}(\rho_k) \rightarrow \Psi_{\rho,k}$ is an isomorphism, then $\Psi'_\rho \rightarrow \Psi_\rho$ is an isomorphism;*
- ii) *If $\Psi'_\rho \rightarrow \Psi_\rho$ is an isomorphism, then $\text{Im}(\rho_k) \rightarrow \Psi_{\rho,k}$ is an isomorphism;*
- iii) *The image of $\Psi'_{\rho,k}$ in $\Psi_{\rho,k}$ is none other than $\text{Im}(\rho_k)$.*

Proposition 4.7. *The kernel of $\Psi'_{\rho,k} \rightarrow \text{Im}(\rho_k)$ is unipotent.*

Proof. Obviously, the kernel in question is also $\text{Ker } \Psi'_{\rho,k} \rightarrow \Psi_{\rho,k}$ as we learn from diagram (4.1). Using Theorem 2.11, we are able to write Ψ'_ρ as $\varprojlim_i \Psi'_i$, where each $\Psi'_{i+1} \rightarrow \Psi'_i$ is a Neron blowup and $\Psi'_0 = \Psi_\rho$. The reader can immediately verify that

$$\text{Ker} \left(\Psi'_{\rho,k} \rightarrow \Psi_{\rho,k} \right) = \varprojlim_i \text{Ker} \left(\Psi'_{i,k} \rightarrow \Psi_{\rho,k} \right).$$

Consequently, using [8, Proposition 2.3 on page 485], we only need to show that each kernel in the above limit is unipotent. This, in turn follows directly from [32, Theorem 1.5] and two fundamental properties of unipotent group schemes: a closed subgroup scheme of a unipotent group scheme is unipotent, and the extension of a unipotent group scheme by a unipotent group scheme is still unipotent [8, Proposition 2.3 on page 485]. \square

We show in the following example that it is possible that neither $\Psi'_{\rho,k} \rightarrow \text{Im}(\rho_k)$ nor $\text{Im}(\rho_k) \rightarrow \Psi_{\rho,k}$ be an isomorphism:

Example 4.8. Let $\rho : G' \rightarrow G$ be the Neron blowup of the origin in the closed fibre of $G = \mathbf{G}_{m,R}$. The diptych of ρ is

$$\begin{array}{ccc} G' & \xrightarrow{\rho} & G \\ \text{id} \uparrow & & \uparrow \text{id} \\ G' & \xrightarrow{\rho} & G. \end{array}$$

Then $\Psi'_{\rho,k} = \mathbf{G}_{a,k}$, $\Psi_{\rho,k} = \mathbf{G}_{m,k}$ and $\text{Im}(\rho_k) = \{e\}$.

Further ahead, in Example 7.9, we will show how this example fits in a more empiric situation.

Let V be a free R -module of finite rank and assume that our G above equals $\mathbf{GL}(V)$. We now interpret Ψ_ρ and Ψ'_ρ in terms of their representation categories. This amounts simply to finding proper references in the literature.

Definition 4.9. Let $\Pi \in \mathbf{FGSch}/R$, as before.

- (1) Let V be an object of $\text{Rep}_R(\Pi)^o$. Write $\mathbf{T}^{a,b}V$ for the representation $V^{\otimes a} \otimes V^{\vee \otimes b}$ and denote by $\langle V \rangle_\otimes$ the full subcategory of $\text{Rep}_R(\Pi)$ having as objects sub-quotients of direct sums $\mathbf{T}^{a_1,b_1}V \oplus \dots \oplus \mathbf{T}^{a_m,b_m}V$ for varying a_i, b_i ;
- (2) Let $\alpha : V' \rightarrow V$ be a monomorphism in $\text{Rep}_R(\Pi)$ with both V and V' free as R -modules. If $\text{Coker}(\alpha)$ is also free as an R -module, we say, following [26, Definitions 10 and 23], that α is a *special monomorphism*. Call an object $V'' \in \text{Rep}_R(\Pi)^o$ a *special sub-quotient* of V if there exists a special monomorphism $V' \rightarrow V$ and an epimorphism $V' \rightarrow V''$. The category of all special sub-quotients of various $\mathbf{T}^{a_1,b_1}V \oplus \dots \oplus \mathbf{T}^{a_m,b_m}V$ is denoted by $\langle V \rangle_\otimes^s$.

Proposition 4.10. *Let V be an object of $\text{Rep}_R(\Pi)^o$ and ρ be the natural homomorphism $\Pi \rightarrow G := \mathbf{GL}(V)$.*

- (1) *The obvious functor $\text{Rep}_R(\Psi_\rho) \rightarrow \text{Rep}_R(\Pi)$ defines an equivalence of categories between $\text{Rep}_R(\Psi_\rho)^o$ and $\langle V \rangle_\otimes^s$;*
- (2) *The obvious functor $\text{Rep}_R(\Psi'_\rho) \rightarrow \text{Rep}_R(\Pi)$ defines an equivalence between $\text{Rep}_R(\Psi'_\rho)$ and $\langle V \rangle_\otimes$.*

Proof. (1) We know that $\text{Rep}_R(\Psi_\rho)^o \rightarrow \text{Rep}_R(\Pi)$ takes values in $\langle V \rangle_\otimes^s$ [26, Proposition 12], that it is fully faithful [9, Proposition 3.2.1(i)], and that its image is stable under special subobjects (loc. cit.). With little effort one then shows that its image is also stable under taking special subquotients.

(2) We know that $\text{Rep}_R(\Psi'_\rho) \rightarrow \text{Rep}_R(\Pi)$ is fully faithful and that its image is stable under taking subobjects [9, Proposition 3.2.1(ii)]. With little effort one then shows that its image is also stable under taking subquotients. Consequently, $\langle V \rangle_\otimes$

is a full subcategory of $\text{Rep}_R(\Psi'_\rho)$ and we set out to prove that any $W \in \text{Rep}_R(\Psi'_\rho)^o$ actually belongs to $\langle V \rangle_\otimes$; this fact, together with [30, Proposition 3, page 41] suffices to show that $\text{Rep}_R(\Psi'_\rho) = \langle V \rangle_\otimes$. Let us regard $R[\Psi'_\rho]$, respectively $R[\Psi_\rho]$, as a Ψ'_ρ -module, respectively Ψ_ρ -module, by means of the right-regular action. Restricting via $\Psi'_\rho \rightarrow \Psi_\rho$, we can also think of $R[\Psi_\rho]$ as a Ψ'_ρ -module. Since W can be embedded in $R[\Psi'_\rho]^{\oplus r}$, we may concentrate on the case where W is a Ψ'_ρ -submodule of $R[\Psi'_\rho]$. Let $m \in \mathbf{N}$ be such that $\pi^m W \subset R[\Psi_\rho]$. It follows that multiplication by π^m defines an embedding of Ψ'_ρ -modules $W \rightarrow R[\Psi_\rho]$. Using part (1) we then conclude that W belongs to $\langle V \rangle_\otimes$. \square

Note that, in general, $\text{Rep}_R(\Psi_\rho)$ is not a full subcategory of $\text{Rep}_R(\Pi)$. This means that we have the following interpretation of the diptych (Definition 4.1) in terms of representation categories:

$$\begin{array}{ccc} \langle V \rangle_\otimes & \longleftarrow & \langle V \rangle_\otimes^s \\ \downarrow & & \downarrow \\ \text{Rep}_R(\Pi) & \longleftarrow & \text{Rep}_R(\mathbf{GL}(V))^o. \end{array}$$

We now deal with the representation theoretic interpretation of the triptych (diagram (4.1)) of ρ . For that, given an R -linear category \mathcal{C} , write $\mathcal{C}_{(k)}$ to denote the full subcategory whose objects W are annihilated by π , i.e. $\pi \cdot \text{id}_W = 0$ in $\text{Hom}_{\mathcal{C}}(W, W)$. We then have a commutative diagram of *solid* arrows between k -linear abelian categories:

$$\begin{array}{ccc} \text{Rep}_R(\Psi'_\rho)_{(k)} & \longleftarrow & \text{Rep}_R(\Psi_\rho)_{(k)} \\ \downarrow & \swarrow & \dashrightarrow \\ & \langle V \otimes k \rangle_\otimes & \\ \downarrow & \swarrow & \\ \text{Rep}_R(\Pi)_{(k)}. & & \end{array} \tag{4.2}$$

From [19, Part I, 10.1, 162] the categories $\text{Rep}_R(-)_{(k)}$ are simply the corresponding representations categories of the group schemes obtained by base change $R \rightarrow k$. Since V is a faithful representation of Ψ_ρ (recall that $\Psi_\rho \rightarrow \mathbf{GL}(V)$ is a closed immersion by construction), $V \otimes k$ is a faithful representation of $\Psi_\rho \otimes k$, so that each object of $\text{Rep}_R(\Psi_\rho)_{(k)}$ is a sub-quotient of some $\bigoplus \mathbf{T}^{a_i, b_i}(V \otimes k)$. This means that the upper horizontal arrow in diagram (4.2) factors through $\langle V \otimes k \rangle_\otimes$, i.e. the dotted arrow exists and still produces a commutative diagram. We conclude that diagram (4.2) captures the essence of diagram (4.1) as the former can easily be completed by introducing the representation category of the general linear group on the lower right corner.

5. Neron blowups of formal subgroup schemes

Let G be group scheme over R which is flat and of finite type. In what follows, we fix a non-negative integer n . Let $H_n \subset G \otimes R_n$ be a closed subgroup scheme (over R_n) cut out by the ideal $I_n \subset R[G]$. We note that, in this case, $\pi^{n+1} \in I_n$.

Definition 5.1. The subring of $K[G]$ obtained by adjoining to $R[G]$ all elements of the form $\pi^{-n-1}a$ with $a \in I_n$ will be denoted by E^n .

Lemma 5.2. Let Δ , ε , and S denote respectively the co-multiplication, the co-identity and the antipode of $K[G]$. Then Δ , ε and S send E^n into $E^n \otimes E^n$, R and E^n respectively.

Proof. Let $a \in I_n$. Since $(\text{Ker } \varepsilon, \pi^{n+1}) \supset (I_n, \pi^{n+1})$, there exists some $c \in \text{Ker } \varepsilon$ and some $f \in R[G]$ such that $a = c + \pi^{n+1}f$. It then follows that $\varepsilon(a) = \pi^{n+1}\varepsilon(f)$, so that $\varepsilon(a\pi^{-n-1}) \in R$. Also, there are $a_i, a'_i \in I_n$ together with $\tau \in R[G] \otimes R[G]$ such that

$$\Delta(a) = \sum a_i \otimes x_i + x'_i \otimes a'_i + \pi^{n+1}\tau.$$

Then,

$$\Delta(\pi^{-n-1}a) = \sum \pi^{-n-1}a_i \otimes x_i + x'_i \otimes \pi^{-n-1}a'_i + \tau.$$

The verification of the statement concerning the antipode is equally trivial and is omitted. □

Definition 5.3. The group scheme

$$\text{Spec } E^n,$$

is called the *Neron blowup* of H_n .

Obviously, there is a morphism of group schemes

$$\text{Spec } E^n \longrightarrow G$$

which, when tensored with K , becomes an isomorphism. The next result, whose simple proof is left to the reader, says a bit more about the group scheme $\text{Spec } E^n$.

Lemma 5.4. *The following claims are true:*

- (1) *The morphism $\text{Spec } E^n \rightarrow G$ when tensored with R_n factors through $H_n \subset G \otimes R_n$;*
- (2) *If $f : G' \rightarrow G$ is an arrow of **FGSch**/ R such that $f \otimes R_n$ factors through $H_n \subset G \otimes R_n$, then f factors uniquely through $\text{Spec } E^n \rightarrow G$.*

We now wish to concentrate on the case “ $n = \infty$.” We assume that R is *complete*. Let \widehat{G} be the completion of G along its closed fibre (it is automatically flat over R) and let $\mathfrak{H} \subset \widehat{G}$ be a closed formal subgroup scheme which is moreover flat over R . We follow traditional notation in the theory of adic algebras and write $R\langle G \rangle$, respectively $R\langle \mathfrak{H} \rangle$, to denote the algebra associated to the formal schemes \widehat{G} , respectively \mathfrak{H} . This being so, $R\langle \mathfrak{H} \rangle$ is a quotient of $R\langle G \rangle$ by some ideal I . In order to carry on proofs, we shall find useful to let $I_n \subset R\langle G \rangle$ be the ideal of the closed subgroup H_n of $G \otimes R_n$ induced by \mathfrak{H} . (The integer n should no more be thought as fixed.) Note that $\pi^{n+1} \in I_n$ and $I_n R\langle G \rangle = (\pi^{n+1}, I)$. Using Lemma 5.4, we see that $E^n \subset E^{n+1}$.

Definition 5.5. We denote by E^∞ the algebra $\cup_n E^n$.

Of course, replacing “ n ” by “ ∞ ” in the statement of Lemma 5.2 still produces a true claim so that we have the following.

Definition 5.6. The group scheme associated to the R -algebra E^∞ of Definition 5.5 will be denoted by $\mathcal{N}_\mathfrak{H}^\infty$ or $\mathcal{N}_\mathfrak{H}^\infty(G)$ if the need presents. The group scheme $\mathcal{N}_\mathfrak{H}^\infty$ will be called the *blowup of \widehat{G} along \mathfrak{H}* .

The group scheme $\text{Spec } E^n$ associated to $H_n = \mathfrak{H} \otimes R_n$ (see Definition 5.3) will be denoted by $\mathcal{N}_\mathfrak{H}^n$ or $\mathcal{N}_\mathfrak{H}^n(G)$ if the need presents. The group scheme $\mathcal{N}_\mathfrak{H}^n$ will be called the *partial blowup of G of level n along \mathfrak{H}* .

If \mathfrak{H} is the completion of a closed subgroup H of G , we put $\mathcal{N}_H^* = \mathcal{N}_\mathfrak{H}^*$.

Example 5.7. The automatic blowup of the identity introduced in Definition 2.18 is, due to Proposition 2.20, $\mathcal{N}_{\{e\}}^\infty(G)$.

Note that the obvious arrow of groups $\mathcal{N}_\mathfrak{H}^\infty \rightarrow G$ becomes an isomorphism when tensored with K . The following is a trivial observation which will prove useful further ahead.

Lemma 5.8. *Let $\varphi : G' \rightarrow G$ be a morphism in \mathbf{FGSch}/R . Denote by $\widehat{\varphi} : \widehat{G}' \rightarrow \widehat{G}$ the morphism induced between π -adic completions. If $\mathfrak{H}' \subset \widehat{G}'$ is a closed and R -flat formal subgroup which is taken by $\widehat{\varphi}$ into \mathfrak{H} , then there exists a unique arrow of affine group schemes $\mathcal{N}_{\mathfrak{H}'}^\infty(G') \rightarrow \mathcal{N}_\mathfrak{H}^\infty(G)$ rendering the diagram*

$$\begin{array}{ccc} \mathcal{N}_{\mathfrak{H}'}^\infty(G') & \longrightarrow & G' \\ \downarrow & & \downarrow \varphi \\ \mathcal{N}_\mathfrak{H}^\infty(G) & \longrightarrow & G \end{array}$$

commutative.

In the following lines we wish to present some elementary features of the blowup along a formal subgroup. We maintain the above notations and introduce

$$K\langle G \rangle = K \otimes R\langle G \rangle \quad \text{and} \quad K\langle \mathfrak{H} \rangle = K \otimes R\langle \mathfrak{H} \rangle.$$

For what comes, the following commutative diagram can be helpful.

$$\begin{array}{ccccc}
 R[G] & \longrightarrow & R\langle G \rangle & \twoheadrightarrow & R\langle \mathfrak{H} \rangle \\
 \downarrow & & \downarrow & & \downarrow \\
 K[G] & \longrightarrow & K\langle G \rangle & \twoheadrightarrow & K\langle \mathfrak{H} \rangle.
 \end{array}$$

Lemma 5.9. *An element $b \in K[G]$ belongs to E^∞ if and only if its image in $K\langle \mathfrak{H} \rangle$ belongs to the image of $R\langle \mathfrak{H} \rangle$.*

Proof. Let $b \in R[G]$ and $m \geq 1$. We assume that the image of $\pi^{-m}b$ in $K\langle \mathfrak{H} \rangle$ coincides with the image of an element $a^* \in R\langle G \rangle$. Then $\pi^m a^*$ and b have the same image in $K\langle \mathfrak{H} \rangle$. This implies that $\pi^m a^* \equiv b \pmod{I}$. As $IR_n[G] = I_n R_n[G]$, we have $0 \equiv b \pmod{I_{m-1}R_{m-1}[G]}$. Hence $\pi^{-m}b \in E^{m-1}(G)$.

Let $a \in I_n$. Since $(I, \pi^{n+1}) = I_n R\langle G \rangle$, we can write $a = \pi^{n+1}a^* + a^{**}$, where $a^* \in R\langle G \rangle$ and $a^{**} \in I$. Then, $\pi^{-n-1}a = a^* + \pi^{-n-1}a^{**}$ in $K\langle G \rangle$ with $a^* \in R\langle G \rangle$ and $\pi^{-n-1}a^{**} \in I \cdot K\langle G \rangle$. The image of $\pi^{-n-1}a$ in $K\langle \mathfrak{H} \rangle$ is then the image of a^* . □

The next statement employs the notion of fibre product of rings, cf. [14, Section 1].

Corollary 5.10. *The first projection*

$$K[G] \times_{K\langle \mathfrak{H} \rangle} R\langle \mathfrak{H} \rangle \longrightarrow K[G]$$

induces an isomorphism between $K[G] \times_{K\langle \mathfrak{H} \rangle} R\langle \mathfrak{H} \rangle$ and E^∞ .

The next result marks a distinctive feature of the Neron blowup of a formal flat subgroup as opposed to Neron blowups centered at the special fibre (cf. the second remark after [32, Theorem 1.4]).

Corollary 5.11. *For each $n \in \mathbb{N}$, the obvious morphism of R_n -groups $\mathcal{N}^\infty \otimes R_n \rightarrow G \otimes R_n$ induces an isomorphism of $\mathcal{N}^\infty \otimes R_n$ with $H_n = \mathfrak{H} \otimes R_n$.*

Proof. In view of Corollary 5.10, it is enough to show that the obvious morphism

$$\theta : R[G] \longrightarrow K[G] \times_{K\langle \mathfrak{H} \rangle} R\langle \mathfrak{H} \rangle,$$

when tensored with R_n , gives the ensuing factorization:

$$\begin{array}{ccc}
 R_n[G] & \longrightarrow & R_n \otimes (K[G] \times_{K\langle \mathfrak{H} \rangle} R\langle \mathfrak{H} \rangle) \\
 \downarrow & \nearrow \sim & \\
 R_n \otimes R\langle \mathfrak{H} \rangle & &
 \end{array}$$

We consider the commutative diagram of R -algebras

$$\begin{array}{ccc}
 K[G] \times_{K(\mathfrak{H})} R(\mathfrak{H}) & \xrightarrow{\text{pr}_2} & R(\mathfrak{H}) \\
 \uparrow \theta & & \uparrow \\
 R[G] & \longrightarrow & R(G),
 \end{array}$$

and claim that $\text{pr}_2 \otimes R_n$ is an isomorphism. Surjectivity is easily checked and injectivity is justified as follows: If $(f, \varphi) \in K[G] \times_{K(\mathfrak{H})} R(\mathfrak{H})$ is such that $\varphi = \pi^{n+1}\varphi'$, then $(f, \varphi) = \pi^{n+1}(\pi^{-n-1}f, \varphi')$, so that (f, φ) belongs to (π^{n+1}) . The proof then follows by tensoring the above commutative square with R_n . \square

We now analyze the standard sequence (see Definition 2.12) associated to $\mathcal{N}_{\mathfrak{H}}^\infty \rightarrow G$. We will see that this sequence can be produced by a spontaneous process. For that we need the following concept.

Definition 5.12 (Strict transform). Let $\mathfrak{H} \subset \widehat{G}$ be as before. Write $I \subset R\langle G \rangle$ for the ideal of \mathfrak{H} and let $\rho : G' \rightarrow G$ be the Neron blowup of some closed subgroup of $G \otimes k$. Then the *strict transform* of \mathfrak{H} , denoted $\rho^\#(\mathfrak{H})$, is the closed formal subgroup of \widehat{G}' cut out by the saturation of the ideal $I \cdot R\langle G' \rangle$, namely $\cup_n (I \cdot R\langle G' \rangle : \pi^n)$. If $H \subset G$ is a closed R -flat subgroup, we can equally define the strict transform $\rho^\#(H)$, which now is a closed, subgroup of G' .

Remarks 5.13.

- (1) By construction, all strict transforms are flat over R ;
- (2) The strict transform of closed subgroups $H \subset G$ is used in [28, Section 1];
- (3) If $\rho : G' \rightarrow G$ and $H \subset G$ are as in Definition 5.12, then the strict transform $\rho^\#(H)$ is just the closure in G' of the closed subgroup $H \otimes K$ in $G' \otimes K$. The same reasoning can be applied in the case of a formal subgroup as $\mathfrak{H} \subset \widehat{G}$ if we use the concept of schematic closure in the setting of rigid geometry, see [5, 0.2.4(vi), page 17]. (Note that $K\langle G \rangle \rightarrow K\langle G' \rangle$ is not usually an isomorphism.);
- (4) The completion \widehat{A} of a noetherian R -algebra A is flat an an A -module [23, Theorem 8.8, page 60]. From this, we conclude that for any noetherian R -algebra A and for any ideal J of A , the equality $J^{\text{sat}}\widehat{A} = (J\widehat{A})^{\text{sat}}$ holds. This proves that, in the notation of the previous item, the completion of $\rho^\#(H)$ is $\rho^\#(\widehat{H})$.

We are now ready to state:

Theorem 5.14. *Let $\mathfrak{H} \subset \widehat{G}$ be as above.*

- (1) *Let $G_0 := G$, $\mathfrak{H}_0 := \mathfrak{H}$ and denote by $\rho_0 : G_1 \rightarrow G_0$ the Neron blowup of $\mathfrak{H}_0 \otimes k$. If \mathfrak{H}_n is defined, write $\rho_n : G_{n+1} \rightarrow G_n$ for the Neron blowup of $\mathfrak{H}_n \otimes k$ and put $\mathfrak{H}_{n+1} := \rho_n^\#(\mathfrak{H}_n)$. Then*

$$\dots \xrightarrow{\rho_1} G_1 \xrightarrow{\rho_0} G_0 = G$$

- is the standard sequence (see Definition 2.12) of $\mathcal{N}_{\mathfrak{H}}^\infty(G) \rightarrow G$;
 (2) The image of the canonical morphism $\mathcal{N}_{\mathfrak{H}}^\infty(G) \otimes k \rightarrow G_{n+1} \otimes k$ is taken isomorphically by $\rho_n \otimes k$ onto the image of $\mathcal{N}_{\mathfrak{H}}^\infty(G) \otimes k \rightarrow G_n \otimes k$.

The proof will need the following preparation. From Corollary 5.11 we know that the image of $\rho_0 \otimes k$ is just $\mathfrak{H}_0 \otimes k$. Let a_1^*, \dots, a_r^* generate the ideal I of \mathfrak{H} and let $a_i \in R[G_0]$ be such that $a_i^* \equiv a_i \pmod{\pi}$. Obviously, the ideal of $\mathfrak{H}_0 \otimes k$ is (π, a_1, \dots, a_r) . The Neron blowup $\rho_0 : G_1 \rightarrow G_0$ of $\mathfrak{H}_0 \otimes k$ is given by the inclusion

$$R[G_0] \subset E = R[G_0] \left[\frac{a_1}{\pi}, \dots, \frac{a_r}{\pi} \right].$$

In the what follows, we write $A\langle \xi_1, \dots, \xi_r \rangle$ for the π -adic completion of $A[\xi_1, \dots, \xi_r]$.

Lemma 5.15. *Let \widehat{E} stand for the π -adic completion of E . Then*

$$\widehat{E} \simeq \frac{R\langle G_0 \rangle \langle \zeta_1, \dots, \zeta_r \rangle}{(\pi \zeta_1 - a_1^*, \dots, \pi \zeta_r - a_r^*)^{\text{sat}}}.$$

Proof. We use bold letters to stand for r -tuples. It is clear that, as $R\langle G_0 \rangle$ -algebras,

$$\frac{R\langle G_0 \rangle \langle \boldsymbol{\xi} \rangle}{(\pi \boldsymbol{\xi} - \mathbf{a})} \simeq \frac{R\langle G_0 \rangle \langle \boldsymbol{\zeta} \rangle}{(\pi \boldsymbol{\zeta} - \mathbf{a}^*)},$$

by means of $\xi_i \mapsto \zeta_i - \pi^{-1}(a_i^* - a_i)$. We then recall that for a noetherian R -algebra A , its completion \widehat{A} is A -flat [23, Theorem 8.8, page 60]. From this, we conclude that for any noetherian R -algebra A and for any ideal J of A , the equality $J^{\text{sat}} \widehat{A} = (\widehat{J A})^{\text{sat}}$ holds. Now

$$\begin{aligned} \widehat{E} &= \left(\frac{R[G_0][\boldsymbol{\xi}]}{(\pi \boldsymbol{\xi} - \mathbf{a})^{\text{sat}}} \right)^\wedge && \text{(by Remark 2.2)} \\ &= \frac{R\langle G_0 \rangle \langle \boldsymbol{\xi} \rangle}{(\pi \boldsymbol{\xi} - \mathbf{a})^{\text{sat}} R\langle G_0 \rangle \langle \boldsymbol{\xi} \rangle} && \text{(by [23, Theorem 8.7, page 60])} \\ &= \frac{R\langle G_0 \rangle \langle \boldsymbol{\xi} \rangle}{(\pi \boldsymbol{\xi} - \mathbf{a})^{\text{sat}}} && \text{(by the argument above).} \quad \square \end{aligned}$$

The algebra \widehat{E} in the statement of the lemma is quite close to a fundamental object in rigid analytic Geometry. Following the terminology of [15, 4.1], the analytic space $\text{Sp } K \otimes \widehat{E}$ is just the *rational domain* $|a_i^*| \leq |\pi|$ in the rigid analytic space $\text{Sp } K \langle G_0 \rangle$.

Proposition 5.16. *The morphism of affine formal schemes $\widehat{G}_1 \rightarrow \widehat{G}_0$ induces an isomorphism between $\rho_0^\#(\mathfrak{H}_0)$ and \mathfrak{H}_0 . The same statement also holds if, instead of a formal subgroup $\mathfrak{H}_0 \subset \widehat{G}_0$, we choose a closed subgroup $H_0 \subset G_0$ which is flat over R .*

Remark 5.17. In the usual theory of blowups, this is known as the statement that the strict transform *is* the blowup of the intersection [11, Proposition IV-21] and that blowing up a “hypersurface” operates no change. In the present setting, a separate consideration has to be made due to the fact that the generic fibre of \widehat{G}_1 is not the generic fibre of \widehat{G}_0 .

Proof. We will omit the verification of the statement concerning a closed subgroup $H_0 \subset G_0$ and concentrate on the formal case.

Write $I^\#$ for the ideal of $\rho_0^\#(\mathfrak{H}_0)$. We must show that the obvious arrow $\phi : R\langle G_0 \rangle / I \rightarrow R\langle G_1 \rangle / I^\#$ is an isomorphism. By the description of $\widehat{E} = R\langle G_1 \rangle$ offered in Lemma 5.15, the equality $I = (a_1^*, \dots, a_r^*)$ and the definition of $I^\#$ as the saturation of $I \cdot R\langle G_1 \rangle$, it is evident that ϕ is surjective. Injectivity will follow from injectivity of $\phi \otimes K$, which for those accustomed to rigid analytic geometry is a triviality. Indeed, there exists a dotted arrow rendering the diagram

$$\begin{array}{ccc}
 & K\langle G_1 \rangle & \\
 & \uparrow & \searrow \text{---} \\
 K\langle G_0 \rangle & \longrightarrow & K\langle G_0 \rangle / I \cdot K\langle G_0 \rangle
 \end{array}$$

commutative, see [15, 4.1.2, page 71]. The injectivity of $\phi \otimes K$ is then obvious as $I \cdot K\langle G_1 \rangle = I^\# \cdot K\langle G_1 \rangle$. □

Proof of Theorem 5.14. We have now obtained an affine and flat group scheme G_1 together with a closed formal subgroup scheme $\rho_0^\#(\mathfrak{H}_0) = \mathfrak{H}_1 \subset \widehat{G}_1$. By functoriality (Lemma 5.8), we arrive at a commutative diagram

$$\begin{array}{ccc}
 \mathcal{N}_{\mathfrak{H}_1}^\infty(G_1) & \longrightarrow & G_1 \\
 \downarrow & & \downarrow \\
 \mathcal{N}_{\mathfrak{H}_0}^\infty(G_0) & \longrightarrow & G_0.
 \end{array}$$

Furthermore, inserting the obvious morphism $\mathcal{N}_{\mathfrak{H}_0}^\infty(G_0) \rightarrow G_1$ in the above diagram still produces a commutative one. (Recall that G_1 is the Neron blowup of $\mathfrak{H}_0 \otimes k$.) Using Corollary 5.11 and Corollary 5.16 we conclude that

$$\mathcal{N}_{\mathfrak{H}_1}^\infty(G_1) \otimes k \xrightarrow{\sim} \mathcal{N}_{\mathfrak{H}_0}^\infty(G_0) \otimes k.$$

As

$$\mathcal{N}_{\mathfrak{H}_1}^\infty(G_1) \otimes K \xrightarrow{\sim} \mathcal{N}_{\mathfrak{H}_0}^\infty(G_0) \otimes K,$$

we deduce that $\mathcal{N}_{\mathfrak{H}_1}^\infty(G_1) \rightarrow \mathcal{N}_{\mathfrak{H}_0}^\infty(G_0)$ is an isomorphism and the image of $\mathcal{N}_{\mathfrak{H}_0}^\infty(G_0) \otimes k \rightarrow G_1 \otimes k$ is the image of $\mathcal{N}_{\mathfrak{H}_1}^\infty(G_1) \otimes k \rightarrow G_1 \otimes k$. Theorem 5.14 can now be proved by induction. □

To end this section, we use Theorem 5.14 to study the standard sequence of the partial blowup of level m of $\mathfrak{X} \subset \widehat{G}$ (cf. Definition 5.3). One probable candidate for this sequence is the truncation of the standard sequence of $\mathcal{N}_{\mathfrak{X}}^{\infty}(G)$. The situation here seems to be a bit more subtle and we only propose an easy consequence of Theorem 5.14.

Corollary 5.18. *We maintain the notation of Theorem 5.14. Let $n \geq 0$ be given. Then there exists $\mu \geq 0$ depending on n and \mathfrak{X} such that the standard sequence of $\mathcal{N}_{\mathfrak{X}}^m \rightarrow G$, for all $m \geq \mu$ starts as*

$$G_{n+1} \xrightarrow{\rho_n} \cdots \xrightarrow{\rho_0} G_0 = G.$$

Proof. We write $J_\nu \subset R[G_\nu]$ for the ideal of $\mathfrak{X}_\nu \otimes k \subset \widehat{G}_\nu \otimes k = G_\nu \otimes k$. From Theorem 5.14, we know that

- (1) Each $R[G_\nu]$ is contained in $E^\infty = \cup_m E^m$;
- (2) The ideal $\pi E^\infty \cap R[G_\nu]$, which cuts out the image of $\mathcal{N}_{\mathfrak{X}}^\infty(G) \otimes k$ in $G_\nu \otimes k$, is just J_ν .

Since $G = G_0$ is assumed to be of finite type, there exists some $\mu_0 \geq 0$ such that $R[G_{n+1}] \subset E^m$ for all $m \geq \mu_0$. We now take $\mu \geq \mu_0$ to be such that the generators of each one of the ideals J_0, \dots, J_n belong to πE^μ . This μ is the one predicted in the statement as in this case, for $m \geq \mu$, the image of $\mathcal{N}_{\mathfrak{X}}^m(G) \otimes k \rightarrow G_\nu \otimes k$ is cut out by J_ν for each $\nu \leq n$. □

Question 5.19. Under which conditions does the standard sequence of the “partial blowup” $\mathcal{N}_{\mathfrak{X}}^n(G) \rightarrow G$ (see Definition 5.3) begin as $G_{n+1} \rightarrow \cdots \rightarrow G_0$, where the G_i are as in Theorem 5.14?

6. Study of a particular class of standard sequences

In this section, we assume that R is complete. Let G be a group scheme over R which is flat and of finite type. Theorem 5.14 guarantees that the centres appearing in the standard sequence of

$$\mathcal{N}_{\mathfrak{X}}^\infty(G) \longrightarrow G,$$

are all isomorphic. In this section we wish to understand a possible converse for this: Corollary 6.10.

As the next paragraph argues, standard sequences with “constant” centres may easily appear; the utility of this study is therefore not restricted to proving a converse to Theorem 5.14.

Assume that k is of characteristic zero and let

$$\cdots \longrightarrow G_n \longrightarrow G_{n-1} \longrightarrow \cdots \longrightarrow G_0,$$

be a standard sequence (cf. Definition 2.13). As $n \mapsto \dim B_n$ is nondecreasing and

$$\dim B_n \leq \dim G_{n,k} = \dim G_{0,K}$$

(the equality follows from [10, VI_B, Corollary 4.3]) there exists $n_0 \in \mathbf{N}$ such that $\dim B_{n_0} = \dim B_n$ for all $n \geq n_0$. Since the kernel of $B_{n+1} \rightarrow B_n$ is either trivial or positive dimensional (it is a subgroup scheme of some $\mathbf{G}_{a,k}^r$ due to Theorem 2.4) we conclude that for $n \geq n_0$, the arrows $B_{n+1} \rightarrow B_n$ are all *isomorphisms*.

It is quite possible that a result more general than Corollary 6.10 holds, but for the moment, our best effort needs higher control on the relation between centres (*i.e.*, the conclusion of Proposition 6.1 hold) so that more hypothesis were introduced.

Proposition 6.1. *Let*

$$\cdots \longrightarrow G_{n+1} \xrightarrow{\rho_n} \cdots \xrightarrow{\rho_0} G_0 = G$$

be a standard sequence (Definition 2.13) where the centre of ρ_n is $B_n \subset G_n \otimes k$. Assume the following particularities.

- i) *The group G_0 is smooth over R , and B_0 is smooth over k ;*
- ii) *For every $n \geq 0$, the restriction $\rho_n \otimes k : B_{n+1} \rightarrow B_n$ is an isomorphism;*
- iii) *There exists an R -flat closed subgroup $L_0 \subset G_0$ with $L_0 \otimes k = B_0$;*
- iv) *For each representation V of B_0 , the first cohomology group $H^1(B_0; V)$ vanishes (linear reductivity).*

Then, for every $n \geq 1$, there exist an inner automorphism $a_n : G_n \rightarrow G_n$ together with a closed and R -flat subgroup $L_n \subset G_n$ enjoying the following properties.

- (1) *For any $n \geq 1$, the closed subgroup L_n is $a_n(\rho_{n-1}^\# L_{n-1})$. (See Definition 5.12 for notation.);*
- (2) *The special fibre of L_n is B_n .*

Proof. We construct a_1 and L_1 . From assumption (ii) and [31, Theorem 14.1], we know that $G_1 \otimes k \rightarrow B_0$ is faithfully flat; we arrive at an exact sequence

$$1 \longrightarrow U_1 \longrightarrow G_1 \otimes k \longrightarrow B_0 \longrightarrow 1, \tag{e}$$

which, again by property (ii), ensures that $G_1 \otimes k$ is a semidirect product $U_1 \rtimes B_1$. In $G_1 \otimes k$, besides B_1 , we have another closed subgroup isomorphic to B_0 , viz. the special fibre of $\Lambda_1 := \rho_0^\#(L_0)$ (see Definition 5.12 and Proposition 5.16). As U_1 is isomorphic to $\mathbf{G}_{a,k}^r$, due to (i) and Theorem 2.4, the action of B_0 on U_1 by conjugation defines a linear action of B_0 on U_0 . (See Section 2.3 for details.) Assumption (iv) together with the exercise proposed in [29, I.5.1] shows that there exists an inner automorphism $a_1 : G_1 \otimes k \rightarrow G_1 \otimes k$ taking $\Lambda_1 \otimes k$ to B_1 .

Now G_1 is smooth over R due to the “fibre-by-fibre” smoothness criterion [16, IV₄, 17.8.2, page 79], the isomorphism $G_1 \otimes K \simeq G_0 \otimes K$ and smoothness of $G_1 \otimes k \rightarrow B_0$ (Theorem 2.4). The infinitesimal lifting criterion allows us to find an inner automorphism of G_1 , parsimoniously called a_1 , such that $a_1(\Lambda_1 \otimes k) = B_1$; define $L_1 := a_1(\Lambda_1)$. This deals with the case $n = 1$. The inductive step “ $n \Rightarrow n + 1$ ” proceeds in the exactly same fashion. □

It is perhaps worth expressing the conclusion of Proposition 6.1 in a pictorial form by means of the following *stair*:

$$\begin{array}{ccc}
 & & \text{higher level} \\
 & & \vdots \\
 & G_2 & \xrightarrow{a_2} G_2 \\
 & \downarrow \rho_1 & \\
 G_1 & \xrightarrow{a_1} & G_1 \\
 \downarrow \rho_0 & & \\
 G_0, & &
 \end{array} \tag{6.1}$$

where

$$\begin{aligned}
 \rho_0 &= \text{blowup of } L_0 \otimes k, \\
 a_1 &= \text{inner automorphism of } G_1, \\
 L_1 &= a_1 [\rho_0^\#(L_0)], \\
 \rho_1 &= \text{blowup of } L_1 \otimes k, \\
 a_2 &= \text{inner automorphism of } G_2 \\
 L_2 &= a_2 [\rho_1^\#(L_1)], \\
 \dots &= \dots .
 \end{aligned} \tag{6.2}$$

For the sake of discussion, we make some definitions.

Definition 6.2 (Spontaneous sequences). A standard sequence as depicted in the above stair is called an *almost spontaneous sequence associated to* $L_0 \subset G_0$. An almost spontaneous standard sequence is *spontaneous* if all the inner automorphisms appearing in it equal the identity.

Using Remark 5.13-(4), another way of stating Theorem 5.14(1) is then:


Theorem 6.3. *Let $H \subset G$ be a closed immersion in \mathbf{FGSch}/R . Then the standard sequence of $\mathcal{N}_H^\infty(G) \rightarrow G$ is the spontaneous sequence of $H \subset G$.*

We now wish to prove Theorem 6.8 below which sheds light on the nature of *almost spontaneous sequences*. We fix, in addition to G , a closed immersion $L_0 \subset G_0 = G$ in \mathbf{FGSch}/R and a stair as in (6.1)-(6.2), which is the almost spontaneous sequence associated to L_0 and the inner automorphisms a_i .

Proposition 6.4. *We maintain the above notation. For each $n \geq 1$, define inner automorphisms $a_{0,n}, \dots, a_{n,n}$ by decreeing that $a_{n,n} = a_n$ and that*

$$\begin{array}{ccc}
 G_m & \xrightarrow{a_{m,n}} & G_m \\
 \rho_{m-1} \downarrow & & \downarrow \rho_{m-1} \\
 G_{m-1} & \xrightarrow{a_{m-1,n}} & G_{m-1},
 \end{array}$$

commutes. In other words, we complete the stair (6.1) as suggested by:



Then


$$G_{n+1} \xrightarrow{\rho_n} \cdots \xrightarrow{\rho_0} G_0 = G,$$

is the spontaneous sequence associated to

$$a_{0,n} \cdots a_{0,2} a_{0,1}(L_0),$$


and truncated at level $n + 1$.

Proof. We proceed by induction and begin with $n = 1$. In this case, we have a commutative diagram



We remark that $a_{0,1} : G_0 \otimes k \rightarrow G_0 \otimes k$ is conjugation by an element of $L_0 \otimes k$. Hence, $L_0 \otimes k = a_{0,1}(L_0) \otimes k$ and commutativity of the above diagram guarantees that $\rho_0^\# [a_{0,1}(L_0)] = a_1 [\rho_0^\#(L_0)]$. The desired result then follows.

Let $n > 1$ and assume that the result is valid for all almost spontaneous sequences associated to $L_1 = a_1 [\rho_0^\#(L_0)] \subset G_1$ and truncated at level n . Since



is commutative, functoriality of the strict transform and the definition of L_1 give us

$$\begin{aligned} \rho_0^\# [a_{0,n} \cdots a_{0,1}(L_0)] &= a_{1,n} \cdots a_{1,1} \left(\rho_0^\#(L_0) \right) \\ &= a_{1,n} \cdots a_{1,2}(L_1). \end{aligned} \tag{6.3}$$

(Recall that by construction $a_{1,1} = a_1$.) Since $a_{0,j}$ is conjugation by an element of $G_0(R)$ belonging to the image of $G_1(R) \rightarrow G_0(R)$, we can affirm that $a_{0,j}$ leaves $L_0 \otimes k$ invariant. Consequently, L_0 and $a_{0,n} \cdots a_{0,1}(L_0)$ have the same closed fibre, from which we conclude that ρ_0 is also the blowup of

$$(a_{0,n} \cdots a_{0,1}(L_0)) \otimes k.$$

Since

$$G_{n+1} \xrightarrow{\rho_n} \cdots \xrightarrow{\rho_1} G_1,$$

is, by induction hypothesis, the spontaneous sequence of $a_{1,n} \cdots a_{1,2}(L_1) \subset G_1$ truncated at level n and, as (see equations (6.3))

$$a_{1,n} \cdots a_{1,2}(L_1) = \rho_0^\# [a_{0,n} \cdots a_{0,1}(L_0)],$$

we obtain that

$$G_{n+1} \xrightarrow{\rho_n} \cdots \xrightarrow{\rho_1} G_1 \xrightarrow{\rho_0} G_0,$$

is the spontaneous sequence associated to $a_{0,n} \cdots a_{0,1}(L_0)$ truncated at level $n + 1$. □

We maintain the above notation and write

$$C_n := a_{0,n} \cdots a_{0,1}(L_0).$$

Proposition 6.5. *For each $n \geq m$, we have $C_n \otimes R_m = C_m \otimes R_m$ as closed subschemes of $G \otimes R_m$. The limit $\mathfrak{C} := \varinjlim_n C_n \otimes R_n$ defines a closed formal subgroup of $\widehat{G}_0 = \varinjlim_n G_0 \otimes R_n$ which is furthermore flat over R .*

The proof relies on the following result.

Lemma 6.6. *Let $\mathcal{L} \subset \mathcal{G}$ be a closed immersion in \mathbf{FGSch}/R . Let*

$$\cdots \xrightarrow{\sigma_0} \mathcal{G}_0 = \mathcal{G}$$

stand for the spontaneous sequence associated to \mathcal{L} . Let $g_{n+1} \in \mathcal{G}_{n+1}(R)$. Then its image by the obvious arrow

$$\mathcal{G}_{n+1}(R) \longrightarrow \mathcal{G}(R) \longrightarrow \mathcal{G}(R_n),$$

lies in $\mathcal{L}(R_n)$.

Proof. To help with the verification, we employ the following observation:

Lemma 6.7. *Let $\mathcal{L} \subset \mathcal{G}$ be a closed immersion in \mathbf{FGSch}/R . If J stands for the ideal of \mathcal{L} and $\sigma : \mathcal{G}' \rightarrow \mathcal{G}$ for the blowup of $\mathcal{L} \otimes k$, then the ideal of the strict transform $\sigma^\#(\mathcal{L})$ of \mathcal{L} contains $\pi^{-1}J$.*

Proof. Evident since the ideal of the strict transform is the saturation $\cup_v(JR[\mathcal{G}'] : \pi^v)$.

We carry on the verification of the lemma. Write $\mathcal{L}_0 = \mathcal{L}$ and define \mathcal{L}_{n+1} as the strict transform of \mathcal{L}_n in \mathcal{G}_{n+1} . Let J_n stand for the ideal of \mathcal{L}_n in $R[\mathcal{G}_n]$. From Lemma 6.7, we know that for any $f_0 \in J_0$, there exists an element $f_{n+1} \in R[\mathcal{G}_{n+1}]$ (belonging to J_{n+1}) such that $\pi^{n+1}f_{n+1} = f_0$. Consequently, if $g_{n+1} : R[\mathcal{G}_{n+1}] \rightarrow R$ is a morphism of R -algebras, we conclude that $g_{n+1}(f_0) = \pi^{n+1}g_{n+1}(f_{n+1})$, so that the morphism $g_0 : R[\mathcal{G}_0] \rightarrow R$ associated to it satisfies $g_0(J_0) \subset (\pi^{n+1})$. Hence, $g_0 \otimes R_n : R_n[\mathcal{G}_0] \rightarrow R_n$ annihilates J_0 , i.e. the corresponding point belongs to $\mathcal{L}_0(R_n)$. □

Proof of Proposition 6.5. We wish first to verify

$$C_{n+1} \otimes_R R_n = C_n \otimes_R R_n. \tag{6.4}$$

Proposition 6.4 ensures us that for each fixed n ,

$$G_{n+1} \xrightarrow{\rho_n} \cdots \xrightarrow{\rho_0} G_0 = G,$$

is the spontaneous sequence of $C_n \subset G_0$ truncated at level $n + 1$. We know that the inner automorphism $a_{0,n+1} : G_0 \rightarrow G_0$, which sends C_n isomorphically to C_{n+1} , is conjugation by an element of the form

$$\rho_n \circ \cdots \circ \rho_0(g_{n+1}), \quad g_{n+1} \in G_{n+1}(R).$$

Hence, due to Lemma 6.6, it is true that

$$a_{0,n+1}(C_n \otimes_R R_n) = C_n \otimes_R R_n.$$

Therefore,

$$\begin{aligned} C_{n+1} \otimes_R R_n &= a_{0,n+1}(C_n) \otimes_R R_n \\ &= a_{0,n+1}(C_n \otimes R_n) \\ &= C_n \otimes R_n. \end{aligned}$$

Another way of expressing equality (6.4) is by saying that, if \mathfrak{k}_n stands for the kernel of $R_n[G_0] \rightarrow R_n[C_n]$, then $R_{n+1}[G_0] \rightarrow R_n[G_0]$ takes \mathfrak{k}_{n+1} onto \mathfrak{k}_n . Consequently, $\varprojlim_n R_n[G_0] \rightarrow \varprojlim_n R_n[C_n]$ is surjective and \mathfrak{C} is a closed formal subscheme of \widehat{G}_0 . We omit the verification that \mathfrak{C} is a subgroup of \widehat{G}_0 and refer the reader to [23, 22.3, page 174] for the statement about flatness. □

Everything is now in place for the proof of

Theorem 6.8. *Let*

$$\cdots \longrightarrow G_{n+1} \xrightarrow{\rho_n} \cdots \xrightarrow{\rho_0} G_0 = G$$

be the almost spontaneous sequence (see Definition 6.2) associated to $L_0 \subset G_0$ as defined on page 1010. If $\mathfrak{C} = \varinjlim C_n \otimes R_n$ stands for the formal closed subgroup of \widehat{G}_0 obtained by means of Proposition 6.4 and Proposition 6.5, then the standard sequence above is the the standard sequence of $\mathcal{N}_{\mathfrak{C}}^\infty(G) \rightarrow G$.

In order to provide a clear proof of Theorem 6.8, we need the following lemma.

Lemma 6.9. *Let $G' \rightarrow G$ and $H \rightarrow G$ be respectively an isomorphism and a closed immersion of \mathbf{FGSch}/R . Denote by $H' \rightarrow G'$ the closed immersion corresponding to $H \rightarrow G$ and let n be an integer.*

Assume that for some $\mu \in \mathbf{N}$, the standard sequence of $\mathcal{N}_H^\mu(G) \rightarrow G$ coincides with the spontaneous sequence associated to H when both are truncated at level $n + 1$. Then the same property holds for $\mathcal{N}_{H'}^\mu(G') \rightarrow G'$.

Proof of Theorem 6.8. We fix some $n \geq 0$ and show that the spontaneous sequence of $\mathfrak{C} \subset \widehat{G}$ truncated at level $n + 1$ is

$$G_{n+1} \xrightarrow{\rho_n} \cdots \xrightarrow{\rho_0} G_0 = G.$$

The goal is to apply Proposition 6.4. To avoid repetitions we let “ σ_{n+1} ” abbreviate “standard sequence truncated at level $n + 1$.” We also omit reference to G when possible.

Let μ be a positive integer satisfying the following properties.

- P1.** The σ_{n+1} of $\mathcal{N}_{\mathfrak{C}}^\infty$ coincides with the σ_{n+1} of $\mathcal{N}_{\mathfrak{C}}^\mu$;
- P2.** The spontaneous sequence of $L_0 \subset G_0$ truncated at level $n + 1$ coincides with the σ_{n+1} of $\mathcal{N}_{L_0}^\mu$;
- P3.** $\mu \geq n$.

That μ exists is a consequence of Corollary 5.18 and Theorem 6.3. From **P1**, the σ_{n+1} of $\mathcal{N}_{\mathfrak{C}}^\infty$ is the σ_{n+1} of $\mathcal{N}_{\mathfrak{C}}^\mu = \mathcal{N}_{C_\mu}^\mu$. From **P2** and Lemma 6.9, the σ_{n+1} of $\mathcal{N}_{C_\mu}^\mu$ is the truncation of the spontaneous sequence of $C_\mu \subset G_0$ at level $n + 1$. Due to Proposition 6.4 and **P3**, the spontaneous sequence associated to $C_\mu \subset G_0$, truncated at level $n + 1$, is just what we started with:

$$G_{n+1} \xrightarrow{\rho_n} \cdots \xrightarrow{\rho_0} G_0 = G. \quad \square$$

We then obtain our main converse of Theorem 5.14 as a consequence of Proposition 6.1, Definition 6.2 and Theorem 6.8.

Corollary 6.10. *Let*

$$\cdots \longrightarrow G_{n+1} \xrightarrow{\rho_n} \cdots \xrightarrow{\rho_0} G_0 = G$$

be a standard sequence (Definition 2.12). Denote the center of the Neron blowup ρ_n by $B_n \subset G_n \otimes k$. Assume that the four hypothesis in Proposition 6.1 hold:

- i) *The group G_0 is smooth over R , and B_0 is smooth over k ;*
- ii) *For every $n \geq 0$, the restriction $\rho_n \otimes k : B_{n+1} \rightarrow B_n$ is an isomorphism;*
- iii) *There exists an R -flat closed subgroup $L_0 \subset G_0$ with $L_0 \otimes k = B_0$;*
- iv) *For each representation V of B_0 , the first cohomology group $H^1(B_0; V)$ vanishes (linear reductivity).*

Then the above standard sequence is the standard sequence of $\mathcal{N}_{\mathfrak{C}}^\infty(G) \rightarrow G$, where \mathfrak{C} is a formal, R -flat, closed subgroup scheme of \widehat{G} .

7. Group schemes over R in differential Galois theory

We now wish to apply the theory so far developed to study differential Galois theory. Let $f : X \rightarrow S$ be a smooth morphism between locally noetherian schemes. We let $\mathcal{D}(X/S)$ be the ring of differential operators which in [16, IV₄, 16.8], respectively [6, Section 2], is denoted by $\text{diff}_{X/S}$, respectively $\text{diff}(\mathcal{O}_X, \mathcal{O}_X)$. We let

$$\mathcal{D}(X/S)\text{-mod},$$

stand for the category of (sheaves of) $\mathcal{D}(X/S)$ -modules whose underlying \mathcal{O}_X -module is coherent. In fact, in the present work, a $\mathcal{D}(X/S)$ -module will always mean an object of $\mathcal{D}(X/S)\text{-mod}$. The obvious action of $\mathcal{D}(X/S)$ on \mathcal{O}_X gives rise to an object of $\mathcal{D}(X/S)\text{-mod}$ which is denoted by $\mathbf{1}$; arrows in $\text{Hom}_{\mathcal{D}(X/S)}(\mathcal{E}, \mathcal{E}')$ will frequently be called horizontal morphisms between \mathcal{E} and \mathcal{E}' , and, in the particular case where $\mathcal{E} = \mathbf{1}$, these will be named horizontal sections of \mathcal{E}' . It is obvious that $\mathcal{D}(X/S)\text{-mod}$ is an abelian category and that the evident functor from $\mathcal{D}(X/S)\text{-mod}$ to coherent modules is exact and faithful. Furthermore, the tensor product of coherent modules induces a tensor product (denoted simply by $\otimes_{\mathcal{O}_X}$ or \otimes) in $\mathcal{D}(X/S)\text{-mod}$.

Remark 7.1. In order to render referencing more effective, we inform the reader that $\mathcal{D}(X/S)\text{-mod}$ is frequently denoted by $\mathbf{str}(X/S)$ (see [9, 26], [6, Section 2]). Also, as [6, Theorem 2.15] explains, if S is a \mathbf{Q} -scheme, $\mathcal{D}(X/S)\text{-mod}$ coincides with the category of \mathcal{O}_X -coherent modules endowed with an integrable connection in the sense of [22, Section 1].

Objects of $\mathcal{D}(X/S)\text{-mod}$ which underlie locally free \mathcal{O}_X -modules (vector bundles) are the objects of a full subcategory $\mathcal{D}(X/S)\text{-mod}^o$. Note that the natural “dualization” in the category of vector bundles allows us to define a dualization in $\mathcal{D}(X/S)\text{-mod}^o$; this will be denoted by a superscript $(-)^{\vee}$.

The proposition below is a basic result in the theory. (For proofs, the reader is directed to [22, Proposition 8.9] of [6, 2.16] for the first statement and to [21, page 40], [26, page 82] or [9, 5.1.1] for the second.)

Proposition 7.2. *If S is the spectrum of a field, any $\mathcal{M} \in \mathcal{D}(X/S)\text{-mod}$ is locally free as an \mathcal{O}_X -module. If S is the spectrum of R , any $\mathcal{M} \in \mathcal{D}(X/S)\text{-mod}$ which is free of π -torsion belongs to $\mathcal{D}(X/S)\text{-mod}^o$ (and conversely).*

We now fix some assumptions and notation which will be in force for the rest of this section. We write

$$S = \text{Spec } R,$$

and let

$$f : X \longrightarrow S,$$

be a smooth and geometrically connected (hence integral) morphism of finite type admitting a section $\xi : S \rightarrow X$. (The discrete valuation ring R is not assumed to be complete.) The base-change functor induced by $\text{Spec } k \rightarrow S$ will be denoted by a subscript “ k ”, and whenever convenient we write “ R ”, or “ k ” in place of “ S ” or “ $\text{Spec } k$ ”.

With these conventions, another cornerstone ensues. (For proofs, the reader should consult [9, Proposition 5.1.1] and [6, 2.16].)

Proposition 7.3. *The pull-back functors*

$$\xi^* : \mathcal{D}(X/R)\text{-mod} \longrightarrow \mathbf{mod}(R),$$

and

$$\xi_k^* : \mathcal{D}(X_k/k)\text{-mod} \longrightarrow \mathbf{mod}(k),$$

are exact and faithful.

In possession of these facts, we can put forward the main definitions of this section.

Definition 7.4.

- (1) Let \mathcal{M} be an object of $\mathcal{D}(X/S)\text{-mod}^o$. Write $\mathbf{T}^{a,b}\mathcal{M}$ for the $\mathcal{D}(X/S)$ -module $\mathcal{M}^{\otimes a} \otimes \mathcal{M}^{\vee \otimes b}$ and denote by $\langle \mathcal{M} \rangle_{\otimes}$ the full subcategory of $\mathcal{D}(X/R)\text{-mod}$ having as objects subquotients of direct sums $\mathbf{T}^{a_1,b_1}\mathcal{M} \oplus \dots \oplus \mathbf{T}^{a_m,b_m}\mathcal{M}$ for varying a_i, b_i ;
- (2) Let $\alpha : \mathcal{M}' \rightarrow \mathcal{M}$ be a monomorphism in $\mathcal{D}(X/R)\text{-mod}$ with both \mathcal{M} and \mathcal{M}' locally free as \mathcal{O}_X -modules. If $\text{Coker}(\alpha)$ is also locally free, we say, following [26, Definitions 10 and 23], that α is a *special monomorphism*. Call an object $\mathcal{M}'' \in \mathcal{D}(X/R)\text{-mod}^o$ a special sub-quotient of \mathcal{M} if there exists a *special monomorphism* $\mathcal{M}' \rightarrow \mathcal{M}$ and an epimorphism $\mathcal{M}' \rightarrow \mathcal{M}''$. The category of all special sub-quotients of various $\mathbf{T}^{a_1,b_1}\mathcal{M} \oplus \dots \oplus \mathbf{T}^{a_m,b_m}\mathcal{M}$ is denoted by $\langle \mathcal{M} \rangle_{\otimes}^s$.

The structure result which will enable us to see the theory of \mathcal{D} -modules through group theoretical lenses is the following (the proof, which is in Saavedra’s seminal work [24], is explained concisely in [9]):

Theorem 7.5. *Let $\mathcal{M} \in \mathcal{D}(X/R)\text{-mod}^o$. The R -point ξ induces an equivalence of abelian tensor categories*

$$\xi^* : \langle \mathcal{M} \rangle_{\otimes} \longrightarrow \text{Rep}_R(\text{Gal}'(\mathcal{M})),$$

where $\text{Gal}'(\mathcal{M})$ is a flat group scheme over R .

Definition 7.6. The group scheme $\text{Gal}'(\mathcal{M})$ is called the *full differential Galois group* of \mathcal{M} .

Definition 7.7. Let $\mathcal{M} \in \mathcal{D}(X/R)\text{-mod}^o$ and write

$$\rho : \text{Gal}'(\mathcal{M}) \longrightarrow \mathbf{GL}(\xi^* \mathcal{M})$$

for the associated representation of the full differential Galois group. The restricted differential Galois group of \mathcal{M} , denoted $\text{Gal}(\mathcal{M})$, is the group scheme Ψ_{ρ} of Definition 4.1.

Consider the diptych (Definition 4.1) of ρ :

$$\begin{array}{ccc} \Psi'_{\rho} & \longrightarrow & \text{Gal}(\mathcal{M}) \\ \uparrow & & \downarrow \\ \text{Gal}'(\mathcal{M}) & \xrightarrow{\rho} & \mathbf{GL}(\xi^* \mathcal{M}). \end{array}$$

From Proposition 4.10(2), Definition 7.4(1) and [9, Theorem 4.1.2] we know that the leftmost arrow above is an *isomorphism*. Of course, due to this, one may think that we have chosen an improper way to present things; the reader who wishes to complete the diagram by including the fundamental group scheme $\Pi(X/S, \xi)$ (see [9, Definition 5.1.4]) in the lower left corner is invited to do so at the expense of having to understand that $\text{Rep}_R(\Pi)$ is not what one might naively think it is. Since we focus on differential Galois groups and not fundamental group schemes, we leave $\Pi(X/S, \xi)$ as inspirational.

Then, as in Section 4 we have:

Proposition 7.8. *The following claims are true.*

- (1) *The arrow $\text{Gal}'(\mathcal{M}) \rightarrow \text{Gal}(\mathcal{M})$ induces an isomorphism on generic fibres and is a possibly infinite iteration of Neron blowups;*
- (2) *The functor ξ^* induces an equivalence*

$$\langle \mathcal{M} \rangle_{\otimes}^s \xrightarrow{\sim} \text{Rep}_R(\text{Gal}(\mathcal{M}))^o.$$

We are then ready to present the “empiric” version of Example 4.8.

Example 7.9. Assume that k is of characteristic zero and that R contains a copy of it. Let $A = R[x, 1/x]$ and define X to be $\text{Spec } A$. In this case, the structure of $\mathcal{D}(X/S)$ is quite simple, and to give the free \mathcal{O}_X -module $\mathcal{L} = \mathcal{O}_X \mathbf{e}$ the structure of a $\mathcal{D}(X/S)$ -module we only need to specify the action of $d/dx =: \partial$ (see 2.6 and 2.15 of [6] for details). We put

$$\partial \mathbf{e} = -\frac{\pi}{x} \mathbf{e}$$

and set out to compute $\text{Gal}(\mathcal{L})$ and $\text{Gal}'(\mathcal{L})$ (at some unspecified R -point).

In $\langle \mathcal{L} \rangle_{\otimes}$ we have the $\mathcal{D}(X/R)$ -module \mathcal{V} given by

$$\begin{aligned} \partial \mathbf{v}_1 &= \frac{1}{x}(\mathbf{v}_2 - \pi \mathbf{v}_1) \\ \partial \mathbf{v}_2 &= 0; \end{aligned}$$

it is the sub-object of $\mathbf{1} \oplus \mathcal{L}$ generated by $\mathbf{v}_1 = (1, \mathbf{e})$ and $\mathbf{v}_2 = (\pi, 0)$. (Needless to say, the construction of \mathcal{V} parallels that in Proposition 3.1.) Since $\pi(\mathbf{1} \oplus \mathcal{L}) \subset \mathcal{V}$, we conclude that $\langle \mathcal{V} \rangle_{\otimes} = \langle \mathcal{L} \rangle_{\otimes}$. We denote by Gal' the full differential Galois group of \mathcal{L} , which is the same as that of \mathcal{V} , at some R -point, so that we have, associated to \mathcal{L} , the representation $\rho : \text{Gal}' \rightarrow \mathbf{G}_{m,R}$. Let $\mathbf{G}'_{m,R}$ stand for the Neron blowup of $\mathbf{G}_{m,R}$ at the origin of the closed fibre. Since ρ_k is the trivial representation (it corresponds to \mathcal{L}_k), we derive a factorization

$$\begin{array}{ccc} \text{Gal}' & \xrightarrow{\rho'} & \mathbf{G}'_{m,R} \\ & \searrow \rho & \downarrow \gamma \\ & & \mathbf{G}_{m,R} \end{array}$$

of ρ . Under this factorization, the faithful representation of $\mathbf{G}'_{m,R}$ constructed in Proposition 3.1 is taken to \mathcal{V} . In particular, we have a faithful representation of $\mathbf{G}'_{m,R} \otimes k$ which is taken to $\mathcal{V}_k \in \langle \mathcal{L} \rangle_{\otimes}$. Now we note that \mathcal{V}_k is the module of the logarithm, so that $\langle \mathcal{V}_k \rangle_{\otimes} \xrightarrow{\sim} \text{Rep}_k(\mathbf{G}_{a,k})$. We arrive at a commutative diagram

$$\begin{array}{ccc} \text{Gal}'_k & \xrightarrow{\rho'_k} & \mathbf{G}'_{m,R} \otimes k \\ \downarrow & \nearrow & \\ \mathbf{G}_{a,k} & & \end{array}$$

Since the arrow

$$\text{Rep}_k(\mathbf{G}'_{m,R} \otimes k) \longrightarrow \text{Rep}_k(\mathbf{G}_{a,k})$$

takes a faithful representation of $\mathbf{G}'_{m,R} \otimes k$ to a faithful representation of $\mathbf{G}_{a,k}$, we conclude that $\mathbf{G}_{a,k} \rightarrow \mathbf{G}'_{m,R} \otimes k$ is a closed embedding. As $\mathbf{G}'_{m,R} \otimes k = \mathbf{G}_{a,k}$ and

k is of characteristic zero, $\mathbf{G}_{a,k} \rightarrow \mathbf{G}'_{m,R} \otimes k$ is an isomorphism. This guarantees that ρ' is an isomorphism [32, Lemma 1.3, page 551] so that the diptych of ρ (see Section 4) is

$$\begin{array}{ccc} \mathbf{G}'_{m,R} & \xrightarrow{\gamma} & \mathbf{G}_{m,R} \\ \rho' \uparrow & & \downarrow \text{id} \\ \text{Gal}' & \longrightarrow & \mathbf{G}_{m,R}. \end{array}$$

To summarise, $\text{Gal}' = \mathbf{G}'_{m,R}$ and $\text{Gal} = \mathbf{G}_{m,R}$. When reduced modulo π , this gives

$$\begin{array}{ccc} \mathbf{G}_{a,k} & \longrightarrow & \mathbf{G}_{m,k} \\ \sim \uparrow & & \downarrow \text{id} \\ \text{Gal}'_k & \xrightarrow{\rho_k} & \mathbf{G}_{m,k}, \end{array}$$

in which the image of ρ_k is the trivial group [31, 8.3, Corollary, page 65].

Remark 7.10. Note that this example shows that [2, Theorem 3.3.1.1, page 729] cannot be true. In fact, what in [2] is defined as $\text{Gal}(\mathcal{L})$, respectively $\text{Gal}(\mathcal{L}_k)$, is what we here call Gal' , respectively $\text{Im}(\rho_k)$, and we have showed that $\text{Gal}'_k \not\cong \text{Im}(\rho_k)$. It seems that the inconsistency in the proof lies in the definition of the arrow u in [2, 3.2.3.4, page 727].

Example 7.11. Assume k to be of characteristic zero and that R contains a copy of it. Let $X = \text{Spec } R[x]$ and define \mathcal{L} to be the free \mathcal{O}_X -module on the element \mathbf{e} . We then introduce on \mathcal{L} the structure of $\mathcal{D}(X/R)$ -module by fixing

$$\frac{\partial}{\partial x} \mathbf{e} = -\pi \mathbf{e}.$$

(The reader should consult [6, 2.15] to understand this construction.) Since this is just the differential equation of the “function” $\exp(-\pi x)$, we are led to consider the following identity:

$$\frac{\partial}{\partial x} \left(\sum_{v=0}^n \frac{(\pi x)^v}{v!} \mathbf{e} \right) = -\frac{x^n}{n!} \pi^{n+1} \mathbf{e}.$$

Consequently, we see that $\mathcal{L} \otimes R_n \simeq \mathbf{1} \otimes R_n$. Using Corollary 2.24, we conclude that $\text{Gal}'(\mathcal{L})$ is the automatic blowup of $\text{Gal}(\mathcal{L}) = \mathbf{G}_{m,R}$ at $\{e\}$.

8. Coincidence of Galois groups in the case of inflated \mathcal{D} -modules

We maintain the notation of Section 7 introduced after Proposition 7.2. In particular, $f : X \rightarrow S = \text{Spec } R$ is a smooth and geometrically connected (hence integral)

morphism admitting a section $\xi : S \rightarrow X$. We also assume that k is algebraically closed and that R contains a copy of it.

We now wish to study the differential Galois groups of certain objects in $\mathcal{D}(X/S)\text{-mod}$ of a very special kind: *inflations*. As X is certainly not of finite type over k – an assumption which is usually present in the literature, see for example [22, Section 1] or [6, Section] – we shall need a few adjustments to our theory.

8.1. Approximating R by smooth k -algebras of finite type

When R is the Zariski or etale local ring of a closed point on a smooth curve over k , it is possible to find a *filtered* family of k -subalgebras $\{R_\lambda\}$ of R enjoying the ensuing properties.

- i) The k -scheme $S_\lambda := \text{Spec } R_\lambda$ is smooth;
- ii) The natural arrow $S \rightarrow \varprojlim_\lambda S_\lambda$ is an isomorphism.

On the other hand, such a family of k -subalgebras invariably exists as can be seen from the arguments given by Artin on [3, pages 40 and 41]. (Unfortunately the *statements* of the results in [3, Section 4] do not immediately apply in our setting. Since it seems pointless to present a quick survey of the concerned theme, we simply leave the reader with the previous reference and the task of examining the mentioned pages.)

As explained in [16, IV₃ 8.8.2(ii) page 28 and IV₄, 17.7.8(ii) page 75], there exists some index α and a smooth morphism of finite type

$$f_\alpha : X_\alpha \longrightarrow S_\alpha,$$

giving back $f : X \rightarrow S$. In addition, [16, IV₃, 12.2.4, page 183] lets us assume that f_α is geometrically integral. In fact, many other properties of f are reflected starting from some model on, see [16, IV₃, 8.10.5] for example.

In order to ease communication in what comes, for $\lambda \geq \alpha$, we call X_λ, f_λ , etc., a *model of X, f* , etc., and we write models as well as “objects associated to these” using bold letters instead of putting subscripts.

Finally, two more conventions will prove useful: A certain X_μ is said to be *thinner* than X_λ if $\mu \geq \lambda$ if $\mu \geq \lambda$, and for each model $f : X \rightarrow S$ of $f : X \rightarrow S$, we write X_k for the fibre of f above the point

$$\text{Spec } k \xrightarrow{\text{closed point}} S \xrightarrow{\text{canonic}} S.$$

8.2. Inflating \mathcal{D} -modules

Let $f : X \rightarrow S$ be a model of $f : X \rightarrow S$ and let $\mathcal{M} \in \mathcal{D}(X/k)\text{-mod}$. Since on X we have a morphism of left \mathcal{O}_X -algebras

$$\mathcal{D}(X/S) \longrightarrow \mathcal{D}(X/k),$$

we can give \mathcal{M} the structure of a $\mathcal{D}(X/S)$ -module. This association, which is clearly functorial, is called the inflation functor and denoted by $\text{Inf}(\mathcal{M})$. Analogously, we define the inflation of \mathcal{M} to X , call it $\text{Inf}_X(\mathcal{M})$, as being the $\mathcal{D}(X/S)$ -module on X induced by $\text{Inf}(\mathcal{M})$ via the canonical morphism $X \rightarrow X$. The reader who is unfamiliar with the process of pulling-back \mathcal{D} -modules will find it made clear in [4, 2.1].

The following result will prove useful.

Lemma 8.1. *Let $\mathcal{M} \in \mathcal{D}(X/k)\text{-mod}$ and $\mathcal{N} \in \langle \mathcal{M} \rangle_{\otimes}$. Then, $\text{Inf}_X(\mathcal{N}) \in \langle \text{Inf}_X(\mathcal{M}) \rangle_{\otimes}^s$.*

Proof. This follows immediately from the fact that all \mathcal{C} in $\mathcal{D}(X/k)\text{-mod}$ underlie locally free \mathcal{O}_X -modules (see Proposition 7.2). □

8.3. The differential Galois group of an inflated \mathcal{D} -module

Theorem 8.2. *We assume in addition that $f : X \rightarrow S$ is projective. Let \mathcal{M} be an object of $\mathcal{D}(X/k)\text{-mod}$ for some model X of X , and let $\mathcal{M} = \text{Inf}_X(\mathcal{M})$. Then the morphism of group schemes over R*

$$\text{Gal}'(\mathcal{M}) \longrightarrow \text{Gal}(\mathcal{M})$$

is an isomorphism.

Example 8.3. Here is an example showing that the Theorem 8.2 *does not* hold if X is simply affine. Let $X = \text{Spec } \mathbf{C}[\pi, x]$ be the affine line over $\mathbf{C}[\pi]$, and $\mathcal{L} = \mathcal{O}_X \mathbf{e}$ the free \mathcal{O}_X -module on \mathbf{e} . We induce on \mathcal{L} the structure of a $\mathcal{D}(X/\mathbf{C})$ -module by

$$\begin{aligned} \frac{\partial}{\partial x} \mathbf{e} &= -\pi \mathbf{e} \\ \frac{\partial}{\partial \pi} \mathbf{e} &= -x \mathbf{e}. \end{aligned}$$

(The reader should consult [6, 2.15] to understand this construction.) Its inflation \mathcal{L} is then defined by

$$\frac{\partial}{\partial x} \mathbf{e} = -\pi \mathbf{e}.$$

As we saw in Example 7.11, $\text{Gal}'(\mathcal{L})$ is the automatic blowup of $\text{Gal}(\mathcal{L}) = \mathbf{G}_{m,R}$ at $\{e\}$.

Our proof of Theorem 8.2 depends on two known results (found in [12,27,33]). Since some work has to be done in order to adapt the substance of these theorems to our setting, we prefer to explain them in Section 9 and give a proof of Theorem 8.2 now.

Proof of Theorem 8.2. To lighten notation, we write G and G' in place of $\text{Gal}(\mathcal{M})$ and $\text{Gal}'(\mathcal{M})$, respectively. We also make use of the notation and conventions introduced on Section 8.1.

It is enough to prove that the functor

$$\text{Rep}_k(G_k) \longrightarrow \langle \mathcal{M}|_{X_k} \rangle_{\otimes},$$

is an equivalence (*cf.* Corollary 4.6(i) and diagram (4.2)). The proof is a consequence of the following claims.

Claim 1. Let \mathcal{N} and \mathcal{N}' be objects in $\mathcal{D}(X/k)\text{-mod}$ and let

$$\begin{array}{ccc} \text{Inf}_X(\mathcal{N})|_{X_k} & \xrightarrow{\theta} & \text{Inf}_X(\mathcal{N}')|_{X_k} \\ \parallel & & \parallel \\ \mathcal{N}|_{X_k} & & \mathcal{N}'|_{X_k}, \end{array}$$

be an arrow of $\mathcal{D}(X_k/k)$ -modules. Then there exists a morphism of $\mathcal{D}(X/S)$ -modules

$$\tilde{\theta} : \text{Inf}_X(\mathcal{N}) \longrightarrow \text{Inf}_X(\mathcal{N}'),$$

lifting θ .

Verification. We have an arrow of $\mathcal{D}(X_k/k)$ -modules

$$\theta : \mathcal{N}|_{X_k} \longrightarrow \mathcal{N}'|_{X_k},$$

which gives us an arrow of $\mathcal{D}(X_k/k)$ -modules

$$\sigma : \mathcal{O}_{X_k} \longrightarrow \mathcal{N}|_{X_k}^{\vee} \otimes \mathcal{N}'|_{X_k} =: \mathcal{E}|_{X_k}.$$

We will show that σ is the restriction of an arrow $\mathcal{O}_X \rightarrow \mathcal{E}$ of $\mathcal{D}(X/S)\text{-mod}$. Now, let

$$\iota : T \longrightarrow \mathcal{E}|_{X_k}$$

be the maximal trivial subobject; the arrow σ can therefore be written as a composition in $\mathcal{D}(X_k/k)\text{-mod}$

$$\mathcal{O}_{X_k} \xrightarrow{\tau} T \xrightarrow{\iota} \mathcal{E}|_{X_k}. \tag{8.1}$$

According to Theorem 9.1-(1), it is possible to find $\mathcal{T} \in \mathcal{D}(S/k)\text{-mod}$ and a morphism of $\mathcal{D}(X/k)$ -modules

$$\tilde{\iota} : f^*(\mathcal{T}) \longrightarrow \mathcal{E}$$

such that ι is the restriction to X_k of $\tilde{\iota}$. As f is proper, flat and geometrically integral, we have $f_*\mathcal{O}_X = \mathcal{O}_S$ [17, X, Proposition 1.2, page 202]; it then follows

that the functor f^* from vector bundles on S to vector bundles on X is full and faithful. As S is affine, we conclude that the morphism of \mathcal{O}_{X_k} -modules

$$f^*(\mathcal{O}_S)|_{X_k} = \mathcal{O}_{X_k} \xrightarrow{\tau} T = f^*(\mathcal{T})|_{X_k},$$

appearing in (8.1) is the restriction of a morphism

$$\tilde{\tau} = f^*(\delta) : f^*(\mathcal{O}_S) \longrightarrow f^*(\mathcal{T}).$$

Of course, δ need not be a morphism of $\mathcal{D}(S/k)$ -modules, but $f^*(\delta)$ is surely a morphism of $\mathcal{D}(X/S)$ -modules, that is, an arrow between inflations

$$\text{Inf}(\mathcal{O}_X) \longrightarrow \text{Inf}(f^*\mathcal{T}).$$

In conclusion, we have proved that σ is the restriction of $\tilde{\tau} \circ \tilde{\tau}$.

Claim 2. For each $V \in \langle \mathcal{M}|_{X_k} \rangle_{\otimes}$, there exist \mathcal{E} and \mathcal{E}' in $\langle \mathcal{M} \rangle_{\otimes}^s$ and an arrow of $\mathcal{D}(X/S)$ -**mod**

$$\tilde{\theta} : \mathcal{E} \longrightarrow \mathcal{E}' ,$$

such that

$$V \simeq \text{Coker}(\tilde{\theta}|_{X_k} : \mathcal{E}|_{X_k} \longrightarrow \mathcal{E}'|_{X_k}).$$

Verification. By definition, $\mathcal{M}|_{X_k} = \mathcal{M}|_{X_k}$. Hence, according to Theorem 9.1-(2) (applied to the dual of V), we can find \mathcal{N} and \mathcal{N}' in $\langle \mathcal{M} \rangle_{\otimes}$ fitting into an exact sequence of $\mathcal{D}(X/k)$ -modules:

$$\mathcal{N}|_{X_k} \xrightarrow{\theta} \mathcal{N}'|_{X_k} \longrightarrow V \longrightarrow 0.$$

Using Claim 1, θ is the restriction to X_k of an arrow in $\mathcal{D}(X/S)$ -**mod**

$$\tilde{\theta} : \text{Inf}_X(\mathcal{N}) \longrightarrow \text{Inf}_X(\mathcal{N}').$$

We then take $\mathcal{E} = \text{Inf}_X(\mathcal{N})$ and $\mathcal{E}' = \text{Inf}_X(\mathcal{N}')$, and the proof is finished since these do belong to $\langle \mathcal{M} \rangle_{\otimes}^s$ (see Lemma 8.1).

Claim 3. Denote by η the composition of functors

$$\text{Rep}_R(G) \longrightarrow \text{Rep}_R(G') \xrightarrow{\sim} \langle \mathcal{M} \rangle_{\otimes}.$$

For each $V \in \text{Rep}_k(G_k)$, there exists $N \in \text{Rep}_R(G)^o$ such that

- (a) V is a quotient of N_k ;
- (b) After eventually passing to a thinner model of X , there exists some $\mathcal{N} \in \langle \mathcal{M} \rangle_{\otimes}$ such that $\eta(N) = \text{Inf}_X(\mathcal{N})$.

Verification. According to [30, Proposition 3, page 41] we can “almost lift” V . Precisely, there exists $E \in \text{Rep}_R(G)^o$ and a surjection $E_k \rightarrow V$. By means of the equivalence

$$\eta : \text{Rep}_R(G)^o \xrightarrow{\sim} \langle \mathcal{M} \rangle_{\otimes}^s,$$

of Proposition 7.8, we can find a diagram in $\mathcal{D}(X/S)\text{-mod}$:

$$\begin{array}{ccc} \mathcal{F} & \longrightarrow & \mathcal{T} \\ \downarrow & & \\ \eta(E), & & \end{array}$$

where \mathcal{T} is some tensor power (see Definition 7.4-(1)) of $\mathcal{M} = \text{Inf}_X(\mathcal{M})$, the vertical arrow is an epimorphism (in $\mathcal{D}(X/S)\text{-mod}$), and the horizontal arrow is special (see Definition 7.4). In particular, $\mathcal{T} = \text{Inf}_X(\mathcal{T})$ for some tensor power \mathcal{T} of \mathcal{M} . According to Theorem 9.2, eventually passing to a thinner model of X , there exists

$$\mathcal{N} \in \langle \mathcal{M} : \mathcal{D}(X/k)\text{-mod} \rangle_{\otimes},$$

and an epimorphism

$$\text{Inf}_X(\mathcal{N}) \longrightarrow \mathcal{F}.$$

Using Lemma 8.1, we conclude that $\text{Inf}_X(\mathcal{N})$ in fact belongs to $\langle \mathcal{M} \rangle_{\otimes}^s$; the above equivalence then produces the desired N , viz. any object of $\in \text{Rep}_R(G)^o$ which is taken by η to $\text{Inf}_X(\mathcal{N})$. Indeed, (b) is verified by construction, and (a) follows from the fact that if $\eta(\theta) : \eta(N) \rightarrow \eta(E)$ is an epimorphism of $\mathcal{D}(X/S)$ -modules, then θ is an epimorphism in $\text{Rep}_R(G)$ (between objects of $\text{Rep}_R(G)^o$).

Claim 4. The functor

$$\eta_k : \text{Rep}_k(G_k) = \text{Rep}_R(G)_{(k)} \longrightarrow \langle \mathcal{M} \rangle_{\otimes, (k)},$$

is full.

Verification. Let $\varphi : \eta_k(V) \rightarrow \eta_k(V')$ be a morphism in $\mathcal{D}(X_k/k)\text{-mod}$. It then fits into a commutative diagram

$$\begin{array}{ccc} \eta(N) \otimes k & \xrightarrow{\theta} & \eta(N') \otimes k \\ \downarrow & & \uparrow \\ \eta_k(V) & \xrightarrow{\varphi} & \eta_k(V'), \end{array}$$

where $\eta(N) = \text{Inf}_X(\mathcal{N})$ and $\eta(N') = \text{Inf}_X(\mathcal{N}')$ are constructed from Claim 3. Claim 1 gives us a lift

$$\tilde{\theta} : \eta(N) \longrightarrow \eta(N')$$

of θ . Lemma 8.1 allied to the fact that \mathcal{N} and \mathcal{N}' belong to $\langle \mathcal{M} \rangle_{\otimes}$ shows that both $\text{Inf}_X(\mathcal{N})$ and $\text{Inf}_X(\mathcal{N}')$ lie in $\langle \mathcal{M} \rangle_{\otimes}^s$. Since η is an equivalence between $\text{Rep}_R(G)^o$

and $\langle \mathcal{M} \rangle_{\otimes}^s$, there exists $\sigma : N \rightarrow N'$ such that $\eta(\sigma) = \tilde{\theta}$. Since the vertical arrows in the above diagram also belong to the image of η_k , the proof of the claim is finished. \square

9. Adapting two known results to our setting

Our goal in this section is to adapt two known results (Theorem 9.1 and Theorem 9.2) to the setting of Theorem 8.2. The conventions and notation we follow are those in the beginning of Section 8. In particular, $f : X \rightarrow S$ is a smooth morphism with geometrically connected (hence integral) fibres and R contains a copy of the residue field k . We also adopt the hypothesis and terminology developed on Section 8.1, in particular S is a limit of affine smooth k -schemes. To handle the projectivity hypothesis on a model of $f : X \rightarrow S$ appearing in the next statement, the reader is referred to [16, IV₃, 8.10.5-(xiii), page 37].

Theorem 9.1. *Assume in addition that $f : X \rightarrow S$ is projective and let $\mathbf{f} : \mathbf{X} \rightarrow \mathbf{S}$ be a projective model of f . Let \mathcal{M} be an object of $\mathcal{D}(\mathbf{X}/k)\text{-mod}$ and write \mathbf{X}_k for the fibre of $\mathbf{f} : \mathbf{X} \rightarrow \mathbf{S}$ above the k -point of \mathbf{S} induced by the k -point of \mathbf{S} .*

- (1) *The maximal trivial subobject of $\mathcal{M}|_{\mathbf{X}_k}$ (this belongs to $\mathcal{D}(\mathbf{X}_k/k)\text{-mod}$) is the restriction of a subobject $\mathcal{T} \subset \mathcal{M}$. Moreover, \mathcal{T} is the pull-back to $\mathcal{D}(\mathbf{X}/k)\text{-mod}$ of an object of $\mathcal{D}(\mathbf{S}/k)\text{-mod}$;*
- (2) *If N belongs to $\langle \mathcal{M}|_{\mathbf{X}_k} \rangle_{\otimes}$, then there exists \mathcal{N} in $\langle \mathcal{M} \rangle_{\otimes}$ and a monic $N \rightarrow \mathcal{N}|_{\mathbf{X}_k}$.*

Proof. If $\Pi(-)$ stands for the affine group scheme associated to the category $\mathcal{D}(\mathbf{X}/k)\text{-mod}$, then the sequence of fundamental group schemes

$$\Pi(\mathbf{X}_k) \longrightarrow \Pi(\mathbf{X}) \longrightarrow \Pi(\mathbf{S}) \longrightarrow 1$$

is exact (cf. [33, Theorem 1.1] or [27, Theorem 1]). Using the characterisation of exactness presented in [13, Appendix], we immediately arrive at the desired conclusion. Moreover, the proof in loc.cit. shows that \mathcal{N} can be chosen in $\langle \mathcal{M} \rangle_{\otimes}$. \square

We now head to prove the following.

Theorem 9.2 (see [12, Theorem 5.10]). *Suppose that $f : X \rightarrow S$ is separated. Let $\mathbf{f} : \mathbf{X} \rightarrow \mathbf{S}$ be a model of $f : X \rightarrow S$ and \mathcal{M} an object of $\mathcal{D}(\mathbf{X}/k)\text{-mod}$. Write*

$$\mathcal{M} := \text{Inf}_{\mathbf{X}}(\mathcal{M})$$

and let

$$\mathcal{V} \longrightarrow \mathcal{M}$$

be a special monic (see Definition 7.4) in the category $\mathcal{D}(\mathbf{X}/\mathbf{S})\text{-mod}$. Then, eventually passing to a thinner model, there exists $\mathcal{S} \in \mathcal{D}(\mathbf{X}/k)\text{-mod}$ and an epimorphism

$$\text{Inf}_{\mathbf{X}}(\mathcal{S}) \longrightarrow \mathcal{V}.$$

Colloquially, each special $\mathcal{D}(X/S)$ -submodule of an inflation is also a quotient of an inflation. Moreover, \mathcal{S} can be chosen in $\langle \mathcal{M} \rangle_{\otimes}$.

The proof hinges on the concept of socle (see Lemma 9.3) and on a criterion which allows us to detect when a \mathcal{D} -submodule of an inflation actually comes from an inflation (Corollary 9.8).

Lemma 9.3. *Let \mathcal{L} and \mathcal{E} be objects of $\mathcal{D}(X/S)\text{-mod}$. Assume that \mathcal{E} , respectively \mathcal{L} , is locally free, respectively locally free of rank one, as an \mathcal{O}_X -module. Define the \mathcal{L} -socle of \mathcal{E} ,*

$$\text{Soc}(\mathcal{L}) \subset \mathcal{E},$$

as the sum of all subobjects of \mathcal{E} which are isomorphic to \mathcal{L} . The following properties are true.

- (1) The sub-object $\text{Soc}(\mathcal{L})$ of \mathcal{E} is special;
- (2) In $\mathcal{D}(X/S)\text{-mod}$ we have $\text{Soc}(\mathcal{L}) \simeq \mathcal{L}^{\oplus r}$ for some r .

Proof. Assume that \mathcal{L} is the trivial $\mathcal{D}(X/S)$ -module $\mathbf{1}$. Since $\langle \mathcal{E} \rangle_{\otimes}$ is equivalent to the category of representations of an affine and flat group scheme over R (see Theorem 7.5), the result becomes a simple exercise. The general case is treated by employing the $\mathcal{D}(X/S)$ -module $\mathcal{L}^{\vee} \otimes \mathcal{E}$. □

We now explain a condition which guarantees that a $\mathcal{D}(X/S)$ -submodule of some inflation is still an inflation. The results follow mainly from:

Theorem 9.4. *Let $T = \text{Spec } \Lambda$ be a k -scheme possessing a system of etale coordinates t_1, \dots, t_m relative to k , and $g : Y \rightarrow T$ a smooth morphism. Let $\mathcal{M} \in \mathcal{D}(Y/k)\text{-mod}$ and let $\mathcal{E} \rightarrow \text{Inf}(\mathcal{M})$ be a monomorphism in $\mathcal{D}(Y/T)\text{-mod}$. If*

$$\text{Hom}_{\mathcal{D}_{Y/T}}(\mathcal{E}, \text{Inf}(\mathcal{M})/\mathcal{E}) = 0,$$

then \mathcal{E} is in fact invariant under the action of $\mathcal{D}(Y/k)$ on \mathcal{M} , that is, it is a $\mathcal{D}(Y/k)$ -submodule.

Proof. Let $U = \text{Spec } A$ be an affine open subset of Y on which there are etale coordinates x_1, \dots, x_n relative to T . We then have etale coordinates t_1, \dots, t_m and x_1, \dots, x_n on U relative to k . Let $\partial^{[q,r]}$ be the differential operators constructed from [16, IV₄, 16.11.2, page 54] by means of the system of etale coordinates $t_1, \dots, t_m, x_1, \dots, x_n$; here $q \in \mathbf{N}^m$ and $r \in \mathbf{N}^n$. Note that $\partial^{[q,0]}(g^*(a)) = g^*(\partial_t^{[q]}(a))$, where $\partial_t^{[q]}$ is the differential operator $\partial_t^{[q]}$ constructed from the set of etale coordinates t_1, \dots, t_m . This means that we have lifted $\partial_t^{[q]}$, but the reader should note that if $\bar{x}_1, \dots, \bar{x}_n$ is another system of etale coordinates on U relative to T , then the differential operator obtained from $\bar{x}_1, \dots, \bar{x}_n$, call it $\bar{\partial}^{[q,0]}$, is not necessarily the same as $\partial^{[q,0]}$. However, $\partial^{[q,0]} - \bar{\partial}^{[q,0]}$ does annihilate Λ and induces therefore a section of $\mathcal{D}(U/T)$.

Let $\mu \in \mathbf{N}^m$ have zeroes all over except at one entry, where it has a 1. We define the additive map

$$\phi_\mu : \mathcal{E}(U) \longrightarrow \mathcal{M}(U)/\mathcal{E}(U), \quad e \longmapsto \partial^{[\mu,0]}(e) \pmod{\mathcal{E}(U)}.$$

It is easily verified that ϕ_μ is A -linear and that

$$\phi_\mu(\partial^{[0,r]}e) = \partial^{[0,r]}\phi_\mu(e).$$

Note that, if $\bar{x}_1, \dots, \bar{x}_n$ is another system of etale coordinates for U over T , employing the notation introduced above, the fact that $\partial^{[\mu,0]} - \bar{\partial}^{[\mu,0]}$ is a section of $\mathcal{D}(U/T)$ means that ϕ_μ is independent of the choice of the system of etale coordinates of U relative to T . This allows us to construct a morphism

$$\phi_\mu : \mathcal{E} \longrightarrow \mathcal{M}/\mathcal{E}$$

of $\mathcal{D}(Y/T)$ -modules. By assumption, $\phi_\mu = 0$; we conclude that \mathcal{E} is invariant under $\mathcal{D}^1(Y/k)$ (we adopt the notation of [16, IV₄, 16.8.3, page 40]).

We now assume that \mathcal{E} is invariant under $\mathcal{D}^\ell(Y/k)$. Pick any $\mu = (\mu_1, \dots, \mu_m) \in \mathbf{N}^m$ such that $\mu_1 + \dots + \mu_m = \ell + 1$. Define, as before, the map of additive groups:

$$\phi_\mu : \mathcal{E}(U) \longrightarrow \mathcal{M}(U)/\mathcal{E}(U), \quad e \longmapsto \partial^{[\mu,0]}(e) \pmod{\mathcal{E}(U)}.$$

If e is a section of \mathcal{E} over U and $\lambda = (\lambda_1, \dots, \lambda_m) \in \mathbf{N}^m$ is such that $\lambda_1 + \dots + \lambda_m \leq \ell$, then $\partial^{[\lambda,0]}(e)$ is, by hypothesis, also a section of \mathcal{E} . Then,

$$\begin{aligned} \phi_\mu(ae) &= \partial^{[\mu,0]}(ae) \pmod{\mathcal{E}(U)} \\ &= \sum_{\mu'+\mu''=\mu} \binom{\mu}{\mu'} \partial^{[\mu',0]}(a) \cdot \partial^{[\mu'',0]}(e) \pmod{\mathcal{E}(U)} \\ &= a \cdot \partial^{[\mu,0]}(e) \pmod{\mathcal{E}(U)} \\ &= a\phi_\mu(e) \pmod{\mathcal{E}(U)}. \end{aligned}$$

This shows that ϕ_μ is A -linear. Note also that [16, 16.11.2, page 54] gives

$$\phi_\mu(\partial^{[0,r]}e) = \partial^{[0,r]}(\phi_\mu(e)).$$

Finally, ϕ_μ is independent of the choice of the system of coordinates x_1, \dots, x_n of U relative to T . To verify this claim we adopt the same notations as before: for any other system $\bar{x}_1, \dots, \bar{x}_n$ of etale coordinates relative to T , we associate the differential operators $\bar{\partial}^{[q,r]}$, and note that $\bar{\partial}^{[\mu,0]} - \partial^{[\mu,0]}$ preserves $\mathcal{E}(U)$, so that ϕ_μ is independent of x_1, \dots, x_n . Hence, we obtain a morphism of $\mathcal{D}(Y/T)$ -modules

$$\phi_\mu : \mathcal{E} \longrightarrow \mathcal{M}/\mathcal{E},$$

which must therefore vanish. Consequently, since μ is arbitrarily chosen, \mathcal{E} is invariant under $\mathcal{D}^{\ell+1}(Y/k)$, and by induction we are done. \square

To deduce from Theorem 9.4 the result we need in order to prove Theorem 9.2, Corollary 9.8, we make a few minor observations.

Lemma 9.5. *Let $v : Y' \rightarrow Y$ be a schematically dominant [16, IV₃, 11.10.2, page 171] affine morphism of schemes. Let \mathcal{F} be a locally free \mathcal{O}_Y -module. Then $v^* : \Gamma(Y, \mathcal{F}) \rightarrow \Gamma(Y', v^*\mathcal{F})$ is injective.*

Lemma 9.6. *Assume in addition that $f : X \rightarrow S$ is separated. Let $\theta : \mathcal{E} \rightarrow \mathcal{M}$ be a morphism of coherent \mathcal{O}_X -modules and assume that on some finite open covering $\{U_i\}$ of X we can find retractions $r_i : \mathcal{M}|_{U_i} \rightarrow \mathcal{E}|_{U_i}$. (In particular, θ is a monomorphism of \mathcal{O}_X -modules.) Then, there exists a model \mathbf{X} of X , a morphism of coherent \mathcal{O}_X -modules $\theta : \mathcal{E} \rightarrow \mathcal{M}$, an open covering $\{U_i\}$ of \mathbf{X} , and retractions $r_i : \mathcal{M}|_{U_i} \rightarrow \mathcal{E}|_{U_i}$ which, when pulled back by the canonical arrow $X \rightarrow \mathbf{X}$, give back θ , $\{U_i\}$, and the $\{r_i\}$.*

Proof. All carefully explained in [16, IV₃], to which we refer in the following lines. From 8.10.5(v) (page 37), we can assume that all models are also separated. Using 8.5.2 (page 20) we obtain our $\theta : \mathcal{E} \rightarrow \mathcal{M}$. Using 8.2.11 (page 11) we find the covering $\{U_i\}$. Note that, employing 8.2.5 (page 9), the scheme U_i is the limit of the inverse images of U_i in the thinner models. To end, another application of 8.5.2 (page 20) gives the retractions. \square

Lemma 9.7. *Suppose in addition that $f : X \rightarrow S$ is separated. Let $\mathcal{M} \in \mathcal{D}(X/S)\text{-mod}$ and let \mathcal{M} stand for the $\mathcal{D}(X/S)$ -module induced by \mathcal{M} . Assume that \mathcal{M} is a vector bundle and let*

$$\theta : \mathcal{E} \longrightarrow \mathcal{M},$$

be a special monic of $\mathcal{D}(X/S)\text{-mod}$. Then there exists, possibly after passing to a thinner model of X , a morphism

$$\theta : \mathcal{E} \longrightarrow \mathcal{M},$$

in $\mathcal{D}(X/S)\text{-mod}$ giving back $\theta : \mathcal{E} \rightarrow \mathcal{M}$. In addition, the morphism of \mathcal{O}_X -modules θ can be chosen to be injective and to have a locally free cokernel.

Proof. Using Lemma 9.6, we can find a morphism of \mathcal{O}_X -coherent modules $\theta : \mathcal{E} \rightarrow \mathcal{M}$ giving back θ and possessing, locally, a retraction. Moreover, employing [16, IV₃, 8.5.5, page 23] it is licit to assume that \mathcal{E} and \mathcal{M} are locally free.

Passing to a thinner model, we can assume that θ comes from an arrow $\theta : \mathcal{E} \rightarrow \mathcal{M}$ of locally free \mathcal{O}_X -modules, see [16, IV₃, 8.5.2(i), page 20 and 8.5.5, page 23]. Moreover, Lemma 9.6 allows us to assume that locally θ possesses a retraction. We now need to show that \mathcal{E} can be given a structure of a $\mathcal{D}(X/S)$ -module, and for that we only need to prove that \mathcal{E} is locally invariant under the action of $\mathcal{D}(X/S)$.

So let $U \subset X$ be an affine open of X having the following properties: (a) the arrow $\theta(U) : \mathcal{E}(U) \rightarrow \mathcal{M}(U)$ possesses a retraction $r : \mathcal{M}(U) \rightarrow \mathcal{E}(U)$, (b) the

$\mathcal{O}(U)$ -module $\mathcal{M}(U)$ is free, and (c) on U we have etale coordinates x_1, \dots, x_n with respect to S . Moreover, we write $U = U \times_S S$.

In this setting, we conclude that, if $1 \otimes m \in \mathcal{O}(U) \otimes_{\mathcal{O}(U)} \mathcal{M}(U)$ actually lies in $\mathcal{O}(U) \otimes_{\mathcal{O}(U)} \mathcal{E}(U)$, then $1 \otimes \theta r(m) = 1 \otimes m$, which shows that $m \in \mathcal{E}(U)$ since $\mathcal{M}(U) \rightarrow \mathcal{O}(U) \otimes_{\mathcal{O}(U)} \mathcal{M}(U)$ is injective. Consequently, if $\partial^{[q]}$ is the differential operator on U constructed via the above system of coordinates [16, IV₄, 16.11.2, page 54] we see that $\partial^{[q]}(e) \in \mathcal{E}(U)$ for each $e \in \mathcal{E}(U)$ as $1 \otimes \partial^{[q]}(e)$ does belong to $\mathcal{O}(U) \otimes_{\mathcal{O}(U)} \mathcal{E}(U)$. □

Corollary 9.8. *We suppose that f is separated. Let $\mathcal{M} \in \mathcal{D}(X/k)\text{-mod}$ and consider a special monic $\theta : \mathcal{E} \rightarrow \text{Inf}_X(\mathcal{M})$ such that*

$$\text{Hom}_{\mathcal{D}(X/S)}(\mathcal{E}, \text{Inf}_X(\mathcal{M})/\mathcal{E}) = 0.$$

Then, replacing eventually X by some thinner model of X , there exist

$$\mathcal{E} \in \mathcal{D}(X/k)\text{-mod}$$

and an arrow $\theta : \mathcal{E} \rightarrow \mathcal{M}$ which is taken to θ by the functor $\text{Inf}_X(-)$.

Proof. Lemma 9.7 shows that $\theta : \mathcal{E} \rightarrow \text{Inf}_X(\mathcal{M})$ is the image of an arrow $\mathcal{E} \rightarrow \text{Inf}(\mathcal{M})$ between $\mathcal{D}(X/S)$ -modules. Moreover, θ is injective and its cokernel, call it \mathcal{C} , is locally free. We now show that

$$\text{Hom}_{\mathcal{D}(X/S)}(\mathcal{E}, \mathcal{C}) = 0.$$

Let $\varphi : \mathcal{E} \rightarrow \mathcal{C}$ be a morphism of $\mathcal{D}(X/S)$ -modules. Then, if $u : X \rightarrow X$ stands for the canonical morphism, we see that $u^*(\varphi) =: \varphi$ is a morphism of $\mathcal{D}(X/S)$ -modules from \mathcal{E} to the cokernel of θ . By assumption, $\varphi = 0$ and because of Lemma 9.5, $\varphi = 0$.

We can now apply Theorem 9.4 to show that θ is an arrow of

$$\mathcal{D}(X/k)\text{-mod}. \quad \square$$

We can now present the

Proof of Theorem 9.2. The idea is to apply Corollary 9.8 to the socle series obtained from Lemma 9.3. We begin by assuming that \mathcal{V} is of rank one as an \mathcal{O}_X -module.

Employing the terminology of Lemma 9.3, define $\text{Soc}_1(\mathcal{V})$ as the \mathcal{V} -socle of \mathcal{M} . If Soc_i is defined, put

$$\text{Soc}_{i+1}(\mathcal{V}) = \begin{array}{l} \text{inverse image in } \mathcal{M} \text{ of} \\ \text{the } \mathcal{V}\text{-socle of } \mathcal{M}/\text{Soc}_i(\mathcal{V}), \end{array}$$

so that

$$\text{Soc}_{i+1}(\mathcal{V})/\text{Soc}_i(\mathcal{V}) \simeq \mathcal{V}\text{-socle of } \mathcal{M}/\text{Soc}_i(\mathcal{V}).$$

Using that the socle is a special subobject (Lemma 9.3) we see that $\text{Soc}_i(\mathcal{V}) \subset \mathcal{M}$ is special too, so, for some $r \in \mathbf{N}$, we have

$$\text{Soc}_r(\mathcal{V}) = \text{Soc}_{r+1}(\mathcal{V}) = \dots$$

Due to the definition of the socle, we conclude that there are no submodules of

$$\mathcal{M}/\text{Soc}_r(\mathcal{V}),$$

isomorphic to \mathcal{V} . The assumption on the rank of \mathcal{V} and Proposition 7.2 then force all arrows

$$\mathcal{V} \longrightarrow \mathcal{M}/\text{Soc}_r(\mathcal{V}),$$

in $\mathcal{D}(X/S)\text{-mod}$ to be null. Using that

$$\text{Soc}_1(\mathcal{V}), \quad \frac{\text{Soc}_2(\mathcal{V})}{\text{Soc}_1(\mathcal{V})}, \quad \dots$$

are all isomorphic to direct sums of \mathcal{V} , we see that any arrow

$$\text{Soc}_r(\mathcal{V}) \longrightarrow \mathcal{M}/\text{Soc}_r(\mathcal{V}),$$

is null. By Corollary 9.8, the $\mathcal{D}(X/S)$ -module $\text{Soc}_r(\mathcal{V})$ is an inflation, *i.e.* after eventually passing to a thinner model, there exists $\mathcal{S} \in \mathcal{D}(X/k)\text{-mod}$ together with a monomorphism $\mathcal{S} \rightarrow \mathcal{M}$ of $\mathcal{D}(X/k)$ -modules such that

$$\text{Inf}_X(\mathcal{S}) \longrightarrow \text{Inf}_X(\mathcal{M}) = \mathcal{M}$$

is our special monic $\text{Soc}_r(\mathcal{V}) \rightarrow \mathcal{M}$. Since \mathcal{V} is a quotient of $\text{Soc}_r(\mathcal{V}) = \text{Inf}_X(\mathcal{S})$ and since \mathcal{S} is a subobject of \mathcal{M} , we are done.

In general, write $m = \text{rank}(\mathcal{V})$ and let $\text{Inf}_X(\mathcal{S}) \rightarrow \det(\mathcal{V})$ be an epimorphism. Then

$$\left\{ \wedge^{m-1} \mathcal{V} \right\}^\vee \otimes \det(\mathcal{V}) \simeq \mathcal{V},$$

which shows that \mathcal{V} is a quotient of $\mathcal{M}^\vee \otimes \text{Inf}_X(\mathcal{S})$. □

References

- [1] S. ANANTHARAMAN, *Schémas en groupes, espaces homogènes et espaces algébriques sur une base de dimension 1*, Bull. Soc. Math. France, Mém. **33** (1973), 5–79.
- [2] Y. ANDRÉ, *Différentielles non commutatives et Théorie de Galois différentielle ou aux différences*, Ann. Sci. École Norm. Sup. (4) **34** (2001), 685–739.
- [3] M. ARTIN, *Algebraic approximation of structures over complete local rings*, Inst. Hautes Études Sci. Publ. Math. **36** (1969) 23–58.
- [4] P. BERTHELOT, “ \mathcal{D} -modules arithmétiques. II: Descente par Frobenius”, Mém. Soc. Math. France, Mém. 81, 2000, 1–138.
- [5] P. BERTHELOT, *Cohomologie rigide et cohomologie rigide à supports propres I*, Université de Rennes, 1996.

- [6] P. BERTHELOT and A. OGUS, “Notes on Crystalline Cohomology”, Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1978.
- [7] N. BOURBAKI, “Algèbre, Chapter X”, Masson, Paris, 1980.
- [8] M. DEMAZURE and P. GABRIEL, “Groupes Algébriques”, Vol. 1, Masson & Cie, Éditeur, Paris; North-Holland Publishing Co., Amsterdam, 1970.
- [9] N. D. DUONG and P. H. HAI, *Tannakian duality over Dedekind ring and applications*, Math. Z. **288** (2018), 1103–1142.
- [10] M. DEMAZURE and A. GROTHENDIECK, *Propriétés générales des schémas en groupes*, In: “Documents mathématiques”, Vol. 7, 2011, Société Mathématique de France.
- [11] D. EISNEBUD and J. HARRIS, “The Geometry of Schemes”, Graduate Texts in Mathematics, Vol. 197, Springer-Verlag, New York, 2000.
- [12] H. ESNAULT and P. H. HAI, *The Gauss-Manin connection and Tannaka duality*, Int. Math. Res. Not. (2006), Art. ID 93978, 35 pp.
- [13] H. ESNAULT, P. H. HAI and X. SUN, *On Nori’s fundamental group scheme*, In: “Progr. in Math.”, Vol. 265, Birkhäuser, Basel, 2008, 377–398.
- [14] D. FERRAND, *Conducteur, descente et pincement*, Bull. Soc. Math. France **131** (2003), 553–585.
- [15] J. FRESNEL and M. VAN DER PUT, “Rigid Analytic Geometry and its Applications”, Progr. in Math., Vol. 218, Birkhäuser Boston, 2004.
- [16] A. GROTHENDIECK (with the collaboration of J. DIEUDONNÉ), *Éléments de Géométrie Algébrique*, Publ. Math. IHÉS **8**, **11** (1961); **17** (1963); **20** (1964); **24** (1965); **28** (1966); **32** (1967).
- [17] A. GROTHENDIECK, “Revêtements étales et groupe fondamental”, Documents Mathématiques, Société Mathématique de France, 2003.
- [18] P. J. HILTON and U. STAMMBACH, “A Course in Homological Algebra”, Second edition, Graduate Texts in Mathematics, Vol. 4, Springer-Verlag, New York, 1997.
- [19] J. C. JANTZEN, “Representations of Algebraic Groups”, Pure Appl. Math., Vol. 131, Academic Press, Inc., Boston, MA, 1987.
- [20] Q. LIU, “Algebraic Geometry and Arithmetic Curves”, Oxford Graduate Texts in Mathematics, Vol. 6, 2002.
- [21] N. KATZ, “Exponential Sums and Differential Equations”, Princeton University Press, 1990.
- [22] N. KATZ, *Nilpotent connections and the monodromy theorem: Applications of a result of Turrittin*, Inst. Hautes Études Sci. Publ. Math. **39** (1970), 175–232.
- [23] H. MATSUMURA, “Commutative Ring Theory”, Cambridge Studies in Advanced Mathematics, Vol. 8, (II ed.), Cambridge University Press, Cambridge, 1989.
- [24] N. SAAVEDRA RIVANO, “Catégories Tannakiennes”, SLN 265, Springer-Verlag, Berlin, 1972.
- [25] C. SABBAH, *The work of Andrey Bolibrukh on isomonodromic deformations*, In: “Differential Equations and Quantum Groups”, IRMA Lect. Math. Theor. Phys., Vol. 9, Eur. Math. Soc., Zürich, 2007, 9–25.
- [26] J. P. DOS SANTOS, *The behaviour of the differential Galois group on the generic and special fibres: A Tannakian approach*, J. Reine Angew. Math. **637** (2009), 63–98.
- [27] J. P. DOS SANTOS, *The homotopy exact sequence for the fundamental group scheme and infinitesimal equivalence relations*, Algebr. Geom. **2** (2015), 535–590.
- [28] T. SEKIGUCHI and N. SUWA, *A case of extension of group schemes over a discrete valuation ring*, Tsukuba J. Math. **14** (1990), 459–487.
- [29] J. P. SERRE, “Galois Cohomology”, Translated by Patrick Ion and revised by the author, Springer-Verlag, Berlin, 1997.
- [30] J. P. SERRE, *Groupe de Grothendieck des schémas en groupes réductifs déployés*, Inst. Hautes Études Sci. Publ. Math. **34** (1968), 37–52.
- [31] W. C. WATERHOUSE, “Introduction to Affine Group Schemes”, Graduate Texts in Mathematics, Vol. 66, Springer-Verlag, New York-Berlin, 1979.

- [32] W. C. WATERHOUSE and B. WEISFEILER, *One-dimensional affine group schemes*, J. Algebra **66** (1980), 550–568.
- [33] LEI ZHANG, *The homotopy sequence of the algebraic fundamental group*, Int. Math. Res. Not. **22** (2014), 6155–6174.

Institute of Mathematics
Vietnam Academy of Science and Technology
18 Hoang Quoc Viet
Cau Giay
Hanoi, Vietnam
nguyendaiduongqn@yahoo.com.vn
phung@math.ac.vn

Institut de Mathématiques de Jussieu
Paris Rive Gauche
4 place Jussieu
Case 247
75252 Paris Cedex 5, France
joao_pedro.dos_santos@yahoo.com